

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 4 : C07H 15/12, C12Q 1/70, 1/02 C12Q 1/06, C12P 21/00, 19/34 C12P 1/04, C12N 15/00, 7/00 C07K 13/00, 3/00, A61K 37/68 A61K 39/00, 45/02	A1	(11) International Publication Number: WO 89/ 01940 (43) International Publication Date: 9 March 1989 (09.03.89)
(21) International Application Number: PCT/US88/02940 (22) International Filing Date: 1 September 1988 (01.09.88) (31) Priority Application Numbers: 094,322 141,649 (32) Priority Dates: 4 September 1987 (04.09.87) 7 January 1988 (07.01.88) (33) Priority Country: US (60) Parent Applications or Grants (63) Related by Continuation US 094,322 (CIP) Filed on 4 September 1987 (04.09.87) US 141,649 (CIP) Filed on 7 January 1988 (07.01.88) (71) Applicant (for all designated States except US): BIOGEN, INC. [US/US]: 14 Cambridge Center, Cambridge, MA 02142 (US).	(72) Inventors; and (75) Inventors/Applicants (for US only): FISHER, Richard A. [US/US]: 1079 Beacon Street, Brookline, MA 02146 (US); GILL, BERT, Walter [US/US]: 107 Upland Road, Cambridge, MA 02140 (US); SATO, Vicki L. [US/US]: 43 Larch Road, Cambridge, MA 02138 (US); FLAVELL, Richard A. [GB/US]: 182 Reservoir Road, Killingworth, CT 06417 (US); MAR-AGANORE, John M. [US/US]: 84 Patrick Road, Tewksbury, MA 01876 (US); LIU, Theresa, R. [US/US]: 102 Emerson Road, Milton, MA 02186 (US). (74) Agents: HALEY, James, F., Jr. et al.: Fish & Neave, 875 Third Avenue, New York, NY 10022-6250 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), DK, FR (European patent), GB (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), US, US. Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: DNA SEQUENCES, RECOMBINANT DNA MOLECULES AND PROCESSES FOR PRODUCING SOLUBLE T4 PROTEINS

3 6 4 9

(57) Abstract

This invention relates to DNA sequences, recombinant DNA molecules and processes for producing soluble T4 protein. More particularly, this invention relates to DNA sequences that are characterized in that they code on expression in an appropriate unicellular host for soluble forms of T4, the receptor on the surface of T4⁺ lymphocytes, or derivatives thereof. In accordance with this invention, the DNA sequences, recombinant DNA molecules and processes of this invention may be employed to produce soluble T4 essentially free of other proteins of human origin. This soluble protein may then advantageously be used in the immunotherapeutic and diagnostic compositions and methods of this invention. The soluble T4-based immunotherapeutic compositions and methods of this invention are useful in treating immunodeficient patients suffering from diseases caused by infective agents whose primary targets are T4⁺ lymphocytes. According to a preferred embodiment, this invention relates to soluble T4-based compositions and methods which are useful in preventing, treating or detecting acquired immune deficiency syndrome, AIDS related complex and HIV infection.

Applicants: Gary Beaudry and
 Paul J. Maddon
 U.S. Serial No. 08/485,163
 Filed: June 7, 1995
 Exhibit 4

3650

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	GA Gabon	MR Mauritania
AU Australia	GB United Kingdom	MW Malawi
BB Barbados	HU Hungary	NL Netherlands
BE Belgium	IT Italy	NO Norway
BG Bulgaria	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CF Central African Republic	KR Republic of Korea	SE Sweden
CG Congo	LI Liechtenstein	SN Senegal
CH Switzerland	LK Sri Lanka	SU Soviet Union
CN Cameroon	LU Luxembourg	TD Chad
DE Germany, Federal Republic of	MC Monaco	TG Togo
DK Denmark	MG Madagascar	US United States of America
FI Finland	ML Mali	
FR France		

00000000

1

**DNA SEQUENCES, RECOMBINANT DNA MOLECULES
AND PROCESSES FOR PRODUCING SOLUBLE
T4 PROTEINS**

5

TECHNICAL FIELD OF INVENTION

This invention relates to DNA sequences, recombinant DNA molecules and processes for producing soluble T4 proteins. More particularly, this invention relates to DNA sequences that are characterized in
10 that they code on expression in an appropriate unicellular host for soluble forms of T4, the receptor on the surface of T4⁺ lymphocytes, or derivatives thereof. In accordance with this invention, the DNA sequences, recombinant DNA molecules and processes
15 of this invention may be employed to produce soluble T4 essentially free of other proteins of human origin. This soluble protein may then advantageously be used in the immunotherapeutic, prophylactic, and diagnostic compositions and methods of this invention.

20

The soluble T4 protein-based immunotherapeutic compositions and methods of this invention are useful in treating immunodeficient patients suffering from diseases caused by infective agents whose primary targets are T4⁺ lymphocytes. According to a
25 preferred embodiment, this invention relates to soluble T4 protein-based compositions and methods which are useful in preventing, treating or detecting

-2-

acquired immune deficiency syndrome, AIDS related complex and HIV infection.

BACKGROUND ART

The class of immune regulatory cells known as T cell lymphocytes can be divided into two broad functional classes, the first class comprising T helper or inducer cells -- which mediate T cell proliferation, lymphokine release and helper cell interactions for Ig release, and the second class comprising T cytotoxic or suppressor cells -- which participate in T cell-mediated killing and immune response suppression. In general, these two classes of lymphocytes are distinguished by expression of one of two surface glycoproteins: T4 (m.w. 55,000-62,000 daltons) which is expressed on T helper or inducer cells, probably as a monomeric protein, or T8 (m.w. 32,000 daltons) which is expressed on T cytotoxic or suppressor cells as a dimeric protein.

The primary structures of T4 and T8 have been deduced from their respective cDNA sequences [P. J. Maddon et al., "The Isolation and Nucleotide Sequence Of A cDNA Encoding The T Cell Surface Protein T4: A New Member Of The Immunoglobulin Gene Family", Cell, 42, pp. 93-104 (1985); D. R. Littman et al., "The Isolation And Sequence Of The Gene Encoding T8: A Molecule Defining Functional Classes Of T Lymphocytes", Cell, 40, pp. 237-46 (1985)]. Both predicted protein sequences define molecules with domains expected for surface antigens, including transmembrane and intracytoplasmic domains at the carboxyl end of the protein. In addition, both proteins contain an amino terminal region which shows striking homology to immunoglobulin and T cell receptor variable regions and which might function during target cell recognition [Maddon et al., supra].

89085513

-3-

In immunocompetent individuals, T4 lymphocytes interact with other specialized cell types of the immune system to confer immunity to or defense against infection [E. L. Reinherz and S. F. Schlossman, "The Differentiation Function Of Human T-Cells", Cell, 19, pp. 821-27 (1980)]. More specifically, T4 lymphocytes stimulate production of growth factors which are critical to a functional immune system. For example, they act to stimulate B cells, the descendants of hemopoietic stem cells, which promote the production of defensive antibodies. They also activate macrophages ("killer cells") to attack infected or otherwise abnormal host cells and they induce monocytes ("scavenger cells") to encompass and destroy invading microbes.

It has been found that the primary target of or receptor for certain infective agents is the T4 surface protein. These agents include, for example, viruses and retroviruses. When T4 lymphocytes are exposed to such agents, they are rendered nonfunctional. As a result, the host's complex immune defense system is destroyed and the host becomes susceptible to a wide range of opportunistic infections.

Such immunosuppression is seen in patients suffering from acquired immune deficiency syndrome ("AIDS"). AIDS is a disease characterized by severe or, typically, complete immunosuppression and attendant host susceptibility to a wide range of opportunistic infections and malignancies. In some cases, AIDS infection is accompanied by central nervous system disorders. Complete clinical manifestation of AIDS is usually preceded by AIDS related complex ("ARC"), a syndrome accompanied by symptoms such as persistent generalized lymphadenopathy, fever and weight loss. The human immunodeficiency virus ("HIV") retrovirus is thought to be

-4-

the etiological agent responsible for AIDS infection and its precursor, ARC [M. G. Sarngadharan et al., "Detection, Isolation And Continuous Production Of Cytopathic Retroviruses (HTLV-III) From Patients With AIDS And Pre-AIDS", Science, 224, pp. 497-508 (1984)].*

Between 85 and 100% of the AIDS/ARCS population test seropositive for HIV [G. N. Shaw et al., "Molecular Characterization Of Human T-Cell Leukemia (Lymphotropic) Virus Type III In The Acquired Immune Deficiency Syndrome", Science, 226, pp. 1165-70 (1984)]. The number of adults in the United States infected with HIV has been estimated to be between 1 and 2.5 million [D. Barnes, "Strategies For An AIDS Vaccine", Science, 233, pp. 1149-53 (1986); M. Rees, "The Sombre View Of AIDS", Nature, 326, pp. 343-45 (1987)]. These estimates include 64,900 individuals who do not belong to an identified group at risk for AIDS [S. L. Sivak and G. P. Wormser, "How Common Is HTLV-III Infection In The United States?", New Eng. J. Med., 313, p. 1352 (1985)]. The apparent annual rate of diagnosis for those infected with HIV virus is between 1 and 2% -- a rate which may increase significantly in future years.

The genome of retroviruses, such as HIV, contains three regions encoding structural proteins. The gag region encodes the core proteins of the virion. The pol region encodes the virion RNA-dependent DNA polymerase (reverse transcriptase). The

30

* In this application, human immunodeficiency virus ("HIV"), the generic term adopted by the human retrovirus subcommittee of the International Committee On Taxonomy Of Viruses to refer to independent isolates from AIDS patients, including human T cell lymphotropic virus type III ("HTLV-III"), lymphadenopathy-associated virus ("LAV"), human immunodeficiency virus type 1 ("HIV-1") and AIDS-associated retrovirus ("ARV") will be used.

3654

89085519

env region encodes the major glycoprotein found in the membrane envelope of the virus and in the cytoplasmic membrane of infected cells. The capacity of the virus to attach to target cell receptors and to
5 cause fusion of cell membranes are two HIV properties controlled by the env gene. These properties are believed to play a fundamental role in the pathogenesis of the virus.

HIV env proteins arise from a precursor
10 polypeptide that, in mature form, is cleaved into a large heavily glycosylated exterior membrane protein of about 481 amino acids -- gp120 -- and a smaller transmembrane protein of about 345 amino acids which may be glycosylated -- gp41 [L. Ratner et al.,
15 "Complete Nucleotide Sequence Of The AIDS Virus, HTLV-III", Nature, 313, pp. 277-84 (1985)].

The host range of the HIV virus is associated with cells which bear the surface glycoprotein T4. Such cells include T4 lymphocytes and
20 brain cells [P. J. Maddon et al., "The T4 Gene Encodes The AIDS Virus Receptor And Is Expressed In The Immune System And The Brain", Cell, 47, pp. 333-48 (1986)]. Upon infection of a host by HIV virus, the T4 lymphocytes are rendered non-functional. The
25 progression of AIDS/ARCS syndromes can be correlated with the depletion of T4⁺ lymphocytes, which display the T4 surface glycoprotein. This T cell depletion, with ensuing immunological compromise, may be attributable to both recurrent cycles of infection and
30 lytic growth from cell-mediated spread of the virus. In addition, clinical observations suggest that the HIV virus is directly responsible for the central nervous system disorders seen in many AIDS patients.

The tropism of the HIV virus for T4⁺ cells
35 is believed to be attributed to the role of the T4 cell surface glycoprotein as the membrane-anchored virus receptor. Because T4 behaves as the HIV virus

-6-

receptor, its extracellular sequence probably plays a direct role in binding HIV. More specifically, it is believed that HIV envelope selectively binds to the T4 epitope(s), using this interaction to initiate entry into the host cell [A. G. Dalgelish et al., "The CD4 (T4) Antigen Is An Essential Component Of The Receptor For The AIDS Retrovirus", Nature, 312, pp. 763-67 (1984); D. Klatzmann et al., "T-Lymphocyte T4 Molecule Behaves As The Receptor For Human Retrovirus LAV", Nature, 312, pp. 767-68 (1984)]. Accordingly, cellular expression of T4 is believed to be sufficient for HIV binding, with the T4 protein serving as a receptor for the HIV virus.

The T4 tropism of the HIV virus has been demonstrated in vitro. When HIV virus isolated from AIDS patients is cultured together with T helper lymphocytes preselected for surface T4, the lymphocytes are efficiently infected, display cytopathic effects, including multinuclear syncytia formation and are killed by lytic growth [D. Klatzmann et al., "Selective Tropism Of Lymphadenopathy Associated Virus (LAV) For Helper-Inducer T Lymphocytes", Science, 225, pp. 59-63 (1984); F. Wong-Staal and R. C. Gallo, "Human T-Lymphotropic Retroviruses", Nature, 317, pp. 395-403 (1985)]. It has been demonstrated that a cloned cDNA version of human T4, when expressed on the surface of transfected cells from non-T cell lineages, including murine and fibroblastoid cells, endows those cells with the ability to bind HIV [P. J. Maddon et al., "The T4 Gene Encodes The AIDS Virus Receptor And Is Expressed In The Immune System And The Brain", Cell, 47, pp. 333-48 (1986)].

During the course of HIV infection, the host mounts both a humoral and a cellular immune response to the virus. These responses include the appearance of antibodies which bind to a number of viral products and which exhibit neutralizing effect

-7-

- or antibody dependent cellular cytotoxic functions [M. Guroff-Robert et al., "HTLV-III-Neutralizing Antibodies In Patients With AIDS And AIDS-Related Complex", Nature, 316, pp. 72-74 (1985); D. D. F. Barin et al., "Virus Envelope Protein Of HTLV-III Represents Major Target Antigen For Antibodies In AIDS Patients", Science, 228, pp. 1094-96 (1985); A. H. Rook et al., "Sera From HTLV-III/LAV Antibody Positive Individuals Mediate Antibody Dependent Cellular Cytotoxicity Against HTLV-III/LAV Infected T Cells", J. Immunol., 138, pp. 1064-68 (1987)].
- Epitopes of the HIV envelope have been identified as important determinants in eliciting a neutralizing antibody response. And, determinants in antibody dependent cellular cytotoxicity ("ADCC") activity include HIV env and, possibly, gag epitopes.

- In the absence to date of effective treatments for AIDS, many efforts have centered on prevention of the disease. Such preventative measures include HIV antibody screening for all blood, organ and semen donors and education of AIDS high-risk groups regarding transmission of the disease.

- Experimental or early-stage clinical treatment of AIDS and ARCS conditions have included the administration of antiviral drugs, such as HPA-23, phosphonoformate, suramin, ribavirin, azidothymidine ("AZT") and dideoxycytidine, which apparently interfere with replication of the virus through reverse transcriptase inhibition. Although each of these drugs exhibits activity against HIV in vitro, only AZT has demonstrated potential benefits in clinical trials. AZT administration in effective amounts, however, has been accompanied by undesirable and debilitating side effects, such as bone marrow depression. It is likely, therefore, that hematologic toxicity will be a major rate limiting factor in the long term use of AZT.

3657

89085519

-8-

Other proposed methods for treating AIDS have focused on the development of agents having activity against steps in the viral replicative cycle other than reverse transcription. Such methods include the administration of interferons or the application of hybridoma technology. Most of these treatment strategies are expected to require the co-administration of immunomodulators, such as interleukin-2.

To date, the need exists for the development of effective immunotherapeutic agents and methods for the treatment of AIDS, ARCS, HIV infection and other immunodeficiencies caused by T lymphocyte depletion or abnormalities.

DISCLOSURE OF THE INVENTION

The present invention solves the problems referred to above by providing, in large amounts, soluble T4 and soluble derivatives thereof that act as receptors for infective agents whose primary target is the T4 surface protein of T4⁺ lymphocytes. Advantageously, this invention also provides soluble T4 essentially free of other proteins of human origin and in a form that is not contaminated by viruses, such as HIV or hepatitis B virus.

As will be appreciated from the disclosure to follow, the DNA sequences and recombinant DNA molecules of this invention are capable of directing, in an appropriate host, the production of soluble T4 or derivatives thereof. The polypeptides of this invention are useful, either as produced in the host or after further derivatization or modification, in a variety of immunotherapeutic compositions and methods for treating immunodeficient patients suffering from diseases caused by infective agents whose primary targets are T4⁺ lymphocytes. According to various embodiments of this invention, such compo-

-9-

sitions and methods relate to a soluble receptor for HIV, soluble T4 proteins and polypeptides and antibodies thereto. The soluble T4 proteins and polypeptides of this invention include monovalent, as well
5 as polyvalent forms.

The compositions and methods of this invention, which are based upon soluble T4 proteins, polypeptides or peptides and antibodies thereto, are particularly useful for the prevention, treatment or
10 detection of the HIV-related infections AIDS and ARC. More specifically, the soluble T4-based compositions and methods of this invention employ soluble T4-like polypeptides -- polypeptides which advantageously interfere with the T4/HIV interaction
15 by blocking or competitive binding mechanisms which inhibit HIV infection of cells expressing the T4 surface protein. These soluble T4-like polypeptides inhibit adhesion between T4⁺ lymphocytes and infective agents which target T4⁺ lymphocytes and inhibit
20 interaction between T4⁺ lymphocytes and antigen presenting cells and targets of T4⁺ lymphocytes mediated killing. By acting as soluble virus receptors, the compositions of this invention may be used as anti-viral therapeutics to inhibit HIV binding to T4⁺
25 cells and virally induced syncytium formation at the level of receptor binding.

This invention accomplishes these goals by providing DNA sequences coding on expression in an appropriate unicellular host for soluble T4 proteins*
30 and soluble derivatives thereof.

* As used in this application, "soluble T4 protein", "soluble T4" and "soluble T4-like polypeptides" include all proteins, polypeptides and peptides which
35 are natural or recombinant soluble T4 proteins, or soluble derivatives thereof, and which are characterized by the immunotherapeutic (anti-retroviral)

(footnote continued on following page)

3659

89085519

This invention also provides recombinant DNA molecules containing these DNA sequences and unicellular hosts transformed with them. Those hosts permit the production of large quantities of the novel soluble T4 proteins, polypeptides, peptides and derivatives of this invention for use in a wide variety of therapeutic, prophylactic and diagnostic compositions and methods

The DNA sequences of this invention are selected from the group consisting of:

(a) the DNA inserts of p199-7, pBG377, pBG380, pBG381, p203-5, pBG391, pBG392, pBG393, pBG394, pBG395, pBG396, pBG397, p211-11, p214-10 and p215-7;

(b) DNA sequences which hybridize to one or more of the foregoing DNA inserts and which code on expression for a soluble T4-like polypeptide; and

(c) DNA sequences which code on expression for a soluble T4-like polypeptide coded for on expression by any of the foregoing DNA inserts and sequences.

According to an alternate embodiment, this invention also relates to a DNA sequence comprising the DNA insert of p170-2, said sequence coding on expression for a T4-like polypeptide. And, this invention also relates to recombinant DNA molecules and processes for producing T4 protein using that DNA sequence.

(footnote continued from preceding page)

or immunogenic activity of soluble T4 protein. They include soluble T4-like compounds from a variety of sources, such as soluble T4 protein derived from natural sources, recombinant soluble T4 protein and synthetic or semi-synthetic soluble T4 protein.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is an autoradi graph depicting the purification of T4 protein from U937 cells by immunoaffinity chromatography.

5 Figure 2 depicts autoradiograph and Western blot data demonstrating that immunoaffinity-purified, solubilized native T4 protein binds to HIV envelope protein.

10 Figure 3 depicts the nucleotide sequence and the derived amino acid sequence of T4 cDNA obtained from PBL clone λ 203-4. In this figure, the amino acids are represented by single letter codes as follows:

15	Phe: F	Leu: L	Ile: I	Met: M
	Val: V	Ser: S	Pro: P	Thr: T
	Ala: A	Tyr: Y	His: H	Gln: Q
	Asn: N	Lys: K	Asp: D	Glu: E
	Cys: C	Trp: W	Arg: R	Gly: G

20 * = position at which a stop codon is present.

In Figure 3, the T4 protein translation start (AA₂₃) is located at the methionine at nucleotides 201-203 and the mature N-terminus is located at the lysine (AA₃) at nucleotides 276-278.

25 Figure 4 is a schematic outline of the construction of cDNA clones pBG312.T4 (also called p171-1) and p170-2.

Figure 5 is a schematic outline of the construction of plasmid pEC100.

30 Figure 6 depicts amino acid comparisons at positions 3, 64 and 231 of various T4 cDNA clones.

Figures 7A and 7B depict the protein domain structure of purified, solubilized T4 protein and recombinant soluble T4 mutants.

35 Figures 8A-8D are schematic outlines of constructions of various intermediate plasmids and other plasmids used to express recombinant soluble T4 ("rsT4") of this invention.

-12-

Figure 9A is a schematic outline of the construction of plasmid p199-7.

Figures 9B and 9C are schematic outlines of the construction of plasmid p203-5.

5 Figure 10 depicts the synthetic oligo-nucleotide linkers employed in various constructions according to this invention.

Figure 11 depicts the nucleotide sequence of the entire plasmid defined by p199-7 ($P_{mutet.rsT4}$) and its rsT4.2 insert and the amino acid sequence deduced from the rsT4 sequence. This includes the ClaI-ClaI cassette which defines the Met perfect rsT4.2 coding sequence.

10 Figure 12 depicts a protein blot analysis of an induction of rsT4.2 expression from SG936/p199-7.

Figure 13 is a schematic outline of the construction of plasmid pBG368.

Figures 14A-14C are schematic outlines of constructions of various plasmids of this invention.

Figure 15 depicts the nucleotide sequence of plasmid pBG391.

Figure 16 depicts the nucleotide sequence of plasmid pBG392. In this figure, the T4 protein translation start (AA_{23}) is located at the methionine at nucleotides 1207-1209 and the mature N-terminus is located at the lysine (AA_3) at nucleotide 1281-84.

Figure 17 is a schematic outline of constructions of various plasmids of this invention.

Figure 18 depicts the synthetic oligonucleotide linkers employed in various constructions according to this invention.

Figure 19 depicts the nucleotide sequence of plasmid pBG394.

Figure 20 depicts the nucleotide sequence of plasmid pBG396.

Figure 21 depicts the nucleotide sequence of plasmid pBG393.

Figure 22 depicts the nucleotide sequence of plasmid pBG395.

5 Figure 23 is a Coomassie stained gel of rsT4.2 purified from the conditioned medium of the pBG380 transfected CHO cell line BG380G of plasmid p196-10.

10 Figure 24 is a schematic outline of the construction of plasmid p196-10.

Figure 25 is a schematic outline of the construction of plasmid pBG394.

Figure 26 is a schematic outline of the construction of plasmid p211-11.

15 Figure 27 is a schematic outline of the construction of plasmid p215-7.

Figure 28 is a schematic outline of the construction of plasmid p218-8.

20 Figure 29A is a Coomassie stained gel of rsT4.113.1 purified from the conditioned medium of pBG211-11 transfected E.coli.

Figure 29B is an autoradiograph depicting a Western blot analysis of rsT4.113.1 expressed in E.coli.

25 Figure 30, panels (a)-(c) depict the purification of rsT4.113.1 from E.coli transformants.

Figure 31, panels (a)-(c) depict the refolding of purified rsT4.113.1.

30 Figure 32 is an autoradiograph depicting the immunoprecipitation of ³⁵S-metabolically labelled CHO cell lines producing recombinant soluble T4.

Figure 33 depicts an immunoblot analysis of COS 7 cell lines producing recombinant soluble T4.

35 Figure 34 depicts in graphic form the results of a competition assay between rsT4.113.1 and rsT4.3 for binding to OKT4A or OKT4.

-14-

Figures 35-37 depict in graphic form the results of competition assays between rsT4.111 and rsT4.3 for binding to, respectively, OKT4A, Leu-3A and OKT4.

5 Figure 38 depicts in graphic form an ELISA assay for rsT4.113.1 from E.coli transformants.

Figure 39 depicts in graphic form the results of a p24 radioimmunoassay using recombinant soluble T4 according to this invention.

10 Figures 40 and 41 depict the results of syncytia inhibition assays using recombinant soluble T4 proteins according to this invention.

Figure 42 is a schematic outline of the construction of plasmid pBiv.1.

15 Figure 43 depicts the bivalent recombinant soluble T4 protein produced by pBiv.1.

DETAILED DESCRIPTION OF THE INVENTION

We isolated the DNA sequences of this invention from two libraries: a λ gt cDNA library derived the T cell tumor line REX and a λ gt10 cDNA library derived from peripheral blood lymphocytes. However, we could also have employed libraries prepared from other cells that express T4. These include, for example, H9 and U937. We also used a human genomic bank to isolate various fragments of the T4 gene.

For screening these libraries, we used a series of chemically synthesized anti-sense oligonucleotide DNA probes based upon the T4 protein sequence set forth in Maddon et al. (1985), supra.

For screening, we hybridized our oligonucleotide probes to our cDNA libraries utilizing a plaque hybridization screening assay. We selected clones hybridizing to several of our probes. And, after isolating and subcloning the cDNA inserts of the selected clones into plasmids, we determined

their nucleotide sequences and compared the amino acid sequences deduced from those nucleotide sequences to the amino acid sequences referred to in Maddon et al. (1985), supra. As a result of these comparisons, we determined that all of our selected clones were characterized by cDNA inserts coding for amino acid sequences of human T4.

We have depicted in Figure 3 the nucleotide sequence of full-length T4 cDNA obtained from deposited clone p170-2 and the amino acid sequence deduced therefrom. That cDNA sequence was subsequently subjected to in vitro site-directed mutagenesis and restriction fragment substitution so that its cDNA sequence was identical to that of Maddon et al.

After modifying our T4 cDNA sequence to be identical to that of Maddon et al., we truncated samples of it in various positions to remove the coding regions for the transmembrane and intracytoplasmic domains. The remaining cDNA sequences encoded a soluble T4 which retained the extracellular region believed to be responsible for HIV binding.

We then constructed various clones characterized by such cDNA inserts coding for human soluble T4. Those cDNA sequences may be used in a variety of ways in accordance with this invention. More particularly, those sequences or portions of them, or synthetic or semi-synthetic copies of them, may be used as DNA probes to screen other human or animal cDNA or genomic libraries to select by hybridization other DNA sequences that are related to soluble T4. Typically, conventional hybridization conditions, e.g., about 20° to 27°C below T_m, are employed in such selections. However, less stringent conditions may be necessary when the library is being screened with a probe from a different species than that from

-16-

which the library is derived, e.g., the screening of a mouse library with a human probe.

Such cDNA inserts, portions of them, or synthetic or semi-synthetic copies of them, may also be used as starting materials to prepare various mutations. Such mutations may be either degenerate, i.e., the mutation does not change the amino acid sequence encoded by the mutated codon, or non-degenerate, i.e., the mutation changes the amino acid sequence encoded by the mutated codon. Both types of mutations may be advantageous in producing or using soluble T4's according to this invention. For example, these mutations may permit higher levels of production or easier purification of soluble T4 or higher T4 activity.

For all of these reasons, the DNA sequences of this invention are selected from the group consisting of:

- (a) the DNA inserts of p199-7, pBG377, pBG380, pBG381, p203-5, pBG391, pBG392, pBG393, pBG394, pBG395, pBG396, pBG397, p211-11, p214-10 and p215-7;
- (b) DNA sequences which hybridize to one or more of the foregoing DNA inserts and which code on expression for a soluble T4-like polypeptide; and
- (c) DNA sequences which code on expression for a soluble T4-like polypeptide coded for on expression by any of the foregoing DNA inserts and sequences.

Preferably, the DNA sequences of this invention code for a polypeptide selected from the group consisting of a polypeptide of the formula AA₂₃-AA₃₆₂ of Figure 3, a polypeptide of the formula AA₁-362 of Figure 3, a polypeptide of the formula Met-AA₁-362 of Figure 3, a polypeptide of the formula AA₁-374 of Figure 3, a polypeptide of the formula Met-AA₁-374 of Figure 3, a polypeptide of the formula AA₁-377 of Figure 3, a polypeptide of the formula

Met-AA₁₋₃₇₇ of Figure 3, a polypeptide of the formula
AA_{-23-AA₃₇₄} of Figure 3, a polypeptide of the formula
AA_{-23-AA₃₇₇} of Figure 3, or portions thereof.

DNA sequences according to this invention
5 also preferably code for a polypeptide selected from
the group consisting of a polypeptide of the formula
AA_{-23-AA₁₈₂} of Figure 16, a polypeptide of the
formula AA_{1-AA₁₈₂} of Figure 16, a polypeptide of
the formula Met-AA₁₋₁₈₂ of Figure 16, a polypeptide
10 of the formula AA_{-23-AA₁₈₂} of Figure 16, followed by
the amino acids asparagine-leucine-glutamine-histidine-
serine-leucine, a polypeptide of the formula
AA_{1-AA₁₈₂} of Figure 16, followed by the amino acids
asparagine-leucine-glutamine-histidine-serine-leucine,
15 a polypeptide of the formula Met-AA₁₋₁₈₂ of Figure 16,
followed by the amino acids asparagine-leucine-
glutamine-histidine-serine-leucine, a polypeptide of
the formula AA_{-23-AA₁₁₃} of Figure 16, a polypeptide
of the formula AA_{1-AA₁₁₃} of Figure 16, a polypeptide
20 of the formula Met-AA₁₋₁₁₃ of Figure 16, a polypeptide
of the formula AA_{-23-AA₁₁₁} of Figure 16, a polypeptide
of the formula AA_{1-AA₁₁₁} of Figure 16, a polypeptide
of the formula Met-AA₁₋₁₁₁ of Figure 16, a polypep-
tide of the formula AA_{-23-AA₁₃₁} of Figure 16, a poly-
25 peptide of the formula AA_{1-AA₁₃₁} of Figure 16, a
polypeptide of the formula Met-AA₁₋₁₃₁ of Figure 16,
a polypeptide of the formula AA_{-23-AA₁₄₅} of Figure 16,
a polypeptide of the formula AA_{1-AA₁₄₅} of Figure 16,
a polypeptide of the formula Met-AA₁₋₁₄₅ of Figure 16,
30 a polypeptide of the formula AA_{-23-AA₁₆₆} of Figure 16,
a polypeptide of the formula AA_{1-AA₁₆₆} of Figure 16,
a polypeptide of the formula Met-AA₁₋₁₆₆ of Figure 16,
or portions thereof.

Additionally, DNA sequences of this inven-
35 tion code for a polypeptide selected from the group
consisting of a polypeptide of the formula AA_{-23-AA₃₆₂}
of mature T4 protein, a polypeptide of the formula
3667

89085519

-18-

AA₁₋₃₆₂ of mature T4 protein, a polypeptide of the formula Met-AA₁₋₃₆₂ of mature T4 protein, a polypeptide of the formula AA₁₋₃₇₄ of mature T4 protein, a polypeptide of the formula Met-AA₁₋₃₇₄ of mature T4 protein, a polypeptide of the formula AA₁₋₃₇₇ of mature T4 protein, a polypeptide of the formula Met-AA₁₋₃₇₇ of mature T4 protein, a polypeptide of the formula AA₂₃₋₃₇₄ of mature T4 protein, a polypeptide of the formula AA₂₃₋₃₇₇ of mature T4 protein, or portions thereof.

DNA sequences according to this invention also code for a polypeptide selected from the group consisting of a polypeptide of the formula AA₂₃₋₁₈₂ of mature T4 protein, a polypeptide of the formula AA₁₋₁₈₂ of mature T4 protein, a polypeptide of the formula Met-AA₁₋₁₈₂ of mature T4 protein, a polypeptide of the formula AA₂₃₋₁₈₂ of mature T4 protein, followed by the amino acids asparagine-leucine-glutamine-histidine-serine-leucine, a polypeptide of the formula AA₁₋₁₈₂ of mature T4 protein, followed by the amino acids asparagine-leucine-glutamine-histidine-serine-leucine, a polypeptide of the formula Met-AA₁₋₁₈₂ of mature T4 protein, followed by the amino acids asparagine-leucine-glutamine-histidine-serine-leucine, a polypeptide of the formula AA₂₃₋₁₁₃ of mature T4 protein, a polypeptide of the formula AA₁₋₁₁₃ of mature T4 protein, a polypeptide of the formula Met-AA₁₋₁₁₃ of mature T4 protein, a polypeptide of the formula AA₂₃₋₁₁₁ of mature T4 protein, a polypeptide of the formula AA₁₋₁₁₁ of mature T4 protein, a polypeptide of the formula Met-AA₁₋₁₁₁ of mature T4 protein, a polypeptide of the formula AA₂₃₋₁₃₁ of mature T4 protein, a polypeptide of the formula AA₁₋₁₃₁ of mature T4 protein, a polypeptide of the formula Met-AA₁₋₁₃₁ of mature T4 protein, a polypeptide of the formula AA₂₃₋₁₄₅ of mature T4 protein, a polypeptide of

89085519

3668

the formula AA_1-AA_{145} of mature T4 protein, a polypeptide of the formula $Met-AA_{1-145}$ of mature T4 protein, a polypeptide of the formula $AA_{23}-AA_{166}$ of mature T4 protein, a polypeptide of the formula AA_1-AA_{166} of mature T4 protein, a polypeptide of the formula $Met-AA_{1-166}$ of mature T4 protein, or portions thereof.

The amino terminal amino acid of mature T4 protein isolated from T cells begins at lysine, the third amino acid of the sequence depicted in Figure 16. Accordingly, soluble T4 proteins also include polypeptides of the formula AA_3-AA_{377} of Figure 16, or portions thereof. Such polypeptides include polypeptides selected from the group consisting of a polypeptide of the formula AA_3 to AA_{362} of Figure 16, a polypeptide of the formula AA_3 to AA_{374} of Figure 16, a polypeptide of the formula AA_3-AA_{182} of Figure 16, a polypeptide of the formula AA_3-AA_{113} of Figure 16, a polypeptide of the formula AA_3-AA_{131} of Figure 16, a polypeptide of the formula AA_3-AA_{145} of Figure 16, a polypeptide of the formula AA_3-AA_{166} of Figure 16, and a polypeptide of the formula AA_3-AA_{111} of Figure 16. Soluble T4 proteins also include the above-recited polypeptides preceded by an N-terminal methionine group.

Soluble T4 protein constructs according to this invention may also be produced by truncating the full length T4 protein sequence at various positions to remove the coding regions for the transmembrane and intracytoplasmic domains, while retaining the extracellular region believed to be responsible for HIV binding. More particularly, soluble T4 polypeptides may be produced by conventional techniques of oligonucleotide directed mutagenesis; restriction digestion, followed by insertion of linkers; or chewing back full length T4 protein with enzymes.

-20-

Alternatively, soluble T4 polypeptides may be chemically synthesized by conventional peptide synthesis techniques, such as solid phase synthesis [R. B. Merrifield, "Solid Phase Peptide Synthesis. I. The Synthesis Of A Tetrapeptide", J. Am. Chem. Soc., 83, pp. 2149-54 (1963)].

The DNA sequences of this invention code for soluble proteins and derivatives that are believed to bind to Major Histocompatibility Complex antigens and envelope glycoprotein of certain retroviruses, such as HIV. Preferably, they also inhibit syncytium formation, believed to be the mode of intracellular HIV virus spread. And, they may inhibit interaction between T4⁺ lymphocytes and antigen-presenting cells and targets of T4⁺ cell mediated killing. Most preferably, they also inhibit adhesion between T4⁺ lymphocytes and infective agents, such as the HIV virus, whose primary targets are T4⁺ lymphocytes.

The DNA sequences of this invention are also useful for producing soluble T4 or its derivatives coded for on expression by them in unicellular hosts transformed with those DNA sequences. As well known in the art, for expression of the DNA sequences of this invention, the DNA sequence should be operatively linked to an expression control sequence in an appropriate expression vector and employed in that expression vector to transform an appropriate unicellular host.

Such operative linking of a DNA sequence of this invention to an expression control sequence, of course, includes the provision of a translation start signal in the correct reading frame upstream of the DNA sequence. If the particular DNA sequence of this invention being expressed does not begin with a methionine, the start signal will result in an additional amino acid -- methionine -- being located at the N-terminus of the product. While

-21-

such methionyl-containing product may be employed directly in the compositions and methods of this invention, it is usually more desirable to remove the methionine before use. Methods are available in the art to remove such N-terminal methionines from polypeptides expressed with them. For example, certain hosts and fermentation conditions permit removal of substantially all of the N-terminal methionine in vivo. Other hosts require in vitro removal of the N-terminal methionine. However, such in vivo and in vitro methods are well known in the art.

A wide variety of host/expression vector combinations may be employed in expressing the DNA sequences of this invention. Useful expression vectors, for example, may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences, such as various known derivatives of SV40 and known bacterial plasmids, e.g., plasmids from E.coli including col E1, pCR1, pBR322, pMB9 and their derivatives, wider host range plasmids, e.g., RP4, phage DNAs, e.g., the numerous derivatives of phage λ , e.g., NM989, and other DNA phages, e.g., M13 and filamentous single stranded DNA phages, yeast plasmids, such as the 2 μ plasmid or derivatives thereof, and vectors derived from combinations of plasmids and phage DNAs, such as plasmids which have been modified to employ phage DNA or other expression control sequences. For animal cell expression, we prefer to use plasmid pBG368, a derivative of pBG312 [R. Cate et al., "Isolation Of The Bovine And Human Genes For Mullerian Inhibiting Substance And Expression Of The Human Gene In Animal Cells", Cell, 45, pp. 685-98 (1986)] which contains the major late promoter of adenovirus 2.

In addition, any of a wide variety of expression control sequences -- sequences that con-

3671

89085513

-22-

trol the expression of a DNA sequence when operatively linked to it -- may be used in these vectors to express the DNA sequence of this invention. Such useful expression control sequences, include, for example, the early and late promoters of SV40 or the adenovirus, the lac system, the trp system, the TAC or TRC system, the major operator and promoter regions of phage λ , the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase, e.g., Pho5, the promoters of the yeast α -mating factors, the polyhedron promoter of the baculovirus system and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof. For animal cell expression, we prefer to use an expression control sequence derived from the major late promoter of adenovirus 2.

A wide variety of unicellular host cells are also useful in expressing the DNA sequences of this invention. These hosts may include well known eukaryotic and prokaryotic hosts, such as strains of E.coli, Pseudomonas, Bacillus, Streptomyces, fungi, such as yeasts, and animal cells, such as CHO and mouse cells, African green monkey cells, such as COS 1, COS 7, BSC 1, BSC 40, and BMT 10, insect cells, and human cells and plant cells in tissue culture. For animal cell expression, we prefer CHO cells and COS 7 cells.

It should of course be understood that not all vectors and expression control sequences will function equally well to express the DNA sequences of this invention. Neither will all hosts function equally well with the same expression system. However, one of skill in the art may make a selection among these vectors, expression control sequences, and hosts without undue experimentation and without

-23-

departing from the scope of this invention. For example, in selecting a vector, the host must be considered because the vector must replicate in it. The vector's copy number, the ability to control that copy number, and the expression of any other proteins encoded by the vector, such as antibiotic markers, should also be considered.

In selecting an expression control sequence, a variety of factors should also be considered. These include, for example, the relative strength of the system, its controllability, and its compatibility with the particular DNA sequence of this invention, particularly as regards potential secondary structures. Unicellular hosts should be selected by consideration of their compatibility with the chosen vector, the toxicity of the product coded for on expression by the DNA sequences of this invention to them, their secretion characteristics, their ability to fold proteins correctly, their fermentation requirements, and the ease of purification of the products coded on expression by the DNA sequences of this invention.

Within these parameters, one of skill in the art may select various vector/expression control system/host combinations that will express the DNA sequences of this invention on fermentation or in large scale animal culture, e.g., CHO cells or COS 7 cells.

The polypeptides produced on expression of the DNA sequences of this invention may be isolated from the fermentation or animal cell cultures and purified using any of a variety of conventional methods. One of skill in the art may select the most appropriate isolation and purification techniques without departing from the scope of this invention.

The polypeptides produced on expression of the DNA sequences of this invention are essentially

3673

89085519

-24-

free of other proteins of human origin. Thus, they are different than T4 protein purified from human lymphocytes.

The polypeptides of this invention are
5 useful in immunotherapeutic compositions and methods. For example, the polypeptides of this invention are active in inhibiting infection by agents whose primary targets are T4⁺ lymphocytes by interfering with their interaction with those target lymphocytes. More
10 preferably, the polypeptides of this invention may be employed to saturate the T4 receptor sites of T4-targeted infective agents. Thus, they exert anti-viral activity by competitive binding with cell surface T4 receptor sites. This effect is plainly
15 of great utility in diseases, such as AIDS, ARC and HIV infection. Accordingly, the polypeptides and methods of this invention may be used to treat humans having AIDS, ARC, HIV infection or antibodies to HIV. In addition, these polypeptides and methods
20 may be used for treating AIDS-like diseases caused by retroviruses, such as simian immunodeficiency viruses, in mammals, including humans.

According to one embodiment of this invention, antibodies to soluble T4 proteins and polypeptides
25 may be used in the treatment, prevention, or diagnosis of AIDS, ARC and HIV infection.

The polypeptides of this invention may also be used in combination with other therapeutics used in the treatment of AIDS, ARC and HIV infection.
30 For example, soluble T4 polypeptides may be used in combination with anti-retroviral agents that block reverse transcriptase, such as AZT, HPA-23, phosphonoformate, suramin, ribavirin and dideoxycytidine. Additionally, these polypeptides may be used
35 with anti-viral agents such as interferons, including alpha interferon, beta interferon and gamma interferon, or glucosidase inhibitors, such as

castanospermine. Such combination therapies advantageously utilize lower dosages of those agents, thus avoiding possible toxicity.

And, the polypeptides of this invention
5 may be used in plasmapheresis techniques or in blood bags for selective removal of viral contaminants from blood. According to this embodiment of the invention, soluble T4 polypeptides may be coupled to a solid support, comprising, for example, plastic or
10 glass beads, or a filter, which is incorporated into a plasmapheresis unit.

Additionally, the compositions of this invention may be employed as immunosuppressants useful in preventing or treating graft-vs-host disease,
15 autoimmune diseases and allograft rejection.

The compositions of this invention typically comprise an immunotherapeutic effective amount of a polypeptide of this invention and a pharmaceutically acceptable carrier. Therapeutic methods of
20 this invention comprise the step of treating patients in a pharmaceutically acceptable manner with those compositions.

The compositions of this invention for use in these therapies may be in a variety of forms.
25 These include, for example, solid, semi-solid and liquid dosage forms, such as tablets, pills, powders, liquid solutions or suspensions, liposomes, suppositories, injectable and infusable solutions. The preferred form depends on the intended mode of administration and therapeutic application. The compositions also preferably include conventional pharmaceutically acceptable carriers and adjuvants which
30 are known to those of skill in the art.

Generally, the pharmaceutical compositions
35 of the present invention may be formulated and administered using methods and compositions similar to those used for other pharmaceutically important poly-

-26-

peptides (e.g., alpha-interferon). Thus, the polypeptides may be stored in lyophilized form, reconstituted with sterile water just prior to administration, and administered by the usual routes of administration such as parenteral, subcutaneous, intravenous, intramuscular or intralesional routes. An effective dosage may be in the range of from 0.5 to 5.0 mg/kg body weight/day, it being recognized that lower and higher doses may also be useful.

This invention also relates to soluble receptors and their use in diagnosing or treating viral agents which target or bind to those receptors. Such soluble receptors may be used as decoys to absorb viral agents and to halt the spread of viral infection. Alternatively, virus-killing agents may be attached to the soluble protein receptors, providing a direct mode of delivery of those agents to the virus.

More particularly, the polypeptides of this invention are useful in diagnostic compositions and methods to detect or monitor the course of HIV infection. Advantageously, these polypeptides are useful in diagnosing variants of the HIV virus, regardless of origin of the infecting HIV agent.

For example, soluble T4 proteins and polypeptides according to this invention, which have a high affinity for HIV, may be advantageously used to increase the sensitivity of HIV assay systems now based upon monoclonal or polyclonal antibodies.

More specifically, soluble T4 proteins and polypeptides may be used to pretreat test plasma to concentrate any HIV present, even in small amounts, so that it is more easily recognized by the antibody. And soluble T4 proteins and polypeptides may be used to purify the HIV envelope protein gp120.

Alternatively, the soluble T4 proteins and polypeptides of this invention may be used to replace

-27-

anti-HIV antibodies now used in various assays. These soluble T4 proteins and polypeptides are preferable to anti-HIV antibodies for two reasons. First, soluble T4, exhibits an affinity for HIV of approximately 10^{-9} , a level which exceeds the 10^{-7} to 10^{-8} values of anti-HIV antibodies. And, while anti-HIV antibodies are more likely to be specific for different HIV isolates, strain variations would not affect a soluble T4 protein-based assay, since all HIV isolates must be capable of interacting with the T4 receptor as a prerequisite to infectivity.

For example, a soluble T4 protein or polypeptide may be linked to an indicator, such as an enzyme, and used in an ELISA assay. Here, soluble T4 advantageously acts as a measure of both HIV in a test sample and any free HIV envelope gp120 protein.

And, polyvalent forms of soluble T4 proteins or polypeptides may be produced, for example, by chemical coupling or genetic fusion techniques, thus increasing even further the avidity of soluble T4 for HIV.

In order that this invention may be better understood, the following examples are set forth. These examples are for purposes of illustration only, and are not to be construed as limiting the scope of the invention in any manner.

EXAMPLES

Purification Of Native Solubilized T4

We purified native T4 from the T4⁺-promonocytic cell line U937 derived from a histocytic lymphoma to approximately 50% purity using immunoaffinity chromatography as follows.

We grew U937 cells [a gift from Dr. Scott Hammer, New England Deaconess Hospital] to 10^6 cells/ml in RPMI 1640, 10% FCS, harvested and washed them in 1X PBS. We then lysed the cell pellet

3677

89085519

-28-

in 20 mM Tris-HCl (pH 7.7), 0.5% NP-40 (a non-ionic detergent), 0.2% NaDOC, 0.2 mM EGTA, 0.2 mM PMSF and 5 µg/ml BPTI at 4×10^7 cells/ml. Because this purification was carried out in the presence of a non-ionic detergent, T4, which is normally membrane-bound via its hydrophobic transmembrane domain, was isolated as a solubilized protein. We spun the lysate in a GS 3 rotor for 10 min at 10,000 rpm and stored the supernatant at -70°C.

Subsequently, we preabsorbed the clarified cell extract with mouse IgG-Sepharose, followed by protein A Sepharose and then passed the flowthrough through an immunoaffinity column comprising immobilized 19Thy anti-T4 monoclonal antibody on Affigel-10 [a gift from Dr. Ellis Reinherz, Dana Farber Cancer Institute, Boston, Massachusetts]. We washed the column extensively and eluted the bound material with 50 mM glycine-HCl (pH 2.5), 0.15 M NaCl, 0.5% NP-40, 5 µg/ml BPTI and 0.2 mM EGTA.

We then separated 10 µl aliquots of each elution fraction on a 10% SDS-PAGE under reducing conditions, with the bands being visualized by silver staining. As shown in Figure 1, a major silver-stained band of 55 Kd was visible. We then carried out two assays on the 55 Kd protein and sequenced the amino terminus of the protein to confirm its identity as native solubilized T4.

Sequencing Of Native Solubilized T4

We determined the N-terminal amino acid sequence of our solubilized native T4 which we isolated from a detergent extract of U937 cells by immunoaffinity chromatography as described above.

Techniques for determining the amino acid sequences of various proteins and peptides derived from them are well known in the art. We chose automated Edman degradation to determine the amino

-29-

terminus of our solubilized native T4. More specifically, we gel purified and electroeluted approximately 5 µg of the solubilized native T4 and then subjected it to automated Edman degradation using a gas phase sequencer (Applied Biosystems 470A). We then identified the PTH-amino acids produced at each cycle of the Edman chemistry by high pressure liquid chromatography, on-line with the sequencer, in a PTH-amino acid analyzer (Applied Biosystems 120A). Direct analysis of the protein provided amino terminal sequence information which, when compared to the amino acid sequence deduced from the cDNA sequence of human T4 [Maddon et al. (1985), supra], identified the purified protein as human T4.

15 Radioimmunoassay Of Native Solubilized T4

To determine that our purification process enriched for T4, we assayed fractions from the immunoaffinity elution step in a T4-specific sandwich radioimmunoassay, based upon the ELISA assay of P. E. Rao et al., in Cellular Immunology, 80, pp. 310-19 (1983). We coated each well of a Removawell strip (Dynatech Labs, Alexandria, Virginia) with 50 µl of 10 µl/ml OKT4 antibody (ATCC #CRL 8002) or MOPC195 (a background binding control) in 0.05 M sodium bicarbonate buffer (pH 9.4) at 4°C overnight. We washed the wells and then filled them with 1% FCS in PBS to saturate the protein binding capacity of the plastic. After removing the 1% FCS solution, we added test samples, in 50 µl aliquots, to the wells. We then incubated the samples for 4 hours at room temperature. Subsequently, we removed the samples and washed the wells four times with 0.05% Tween-20 in PBS. We then added ¹²⁵I-labelled 19Thy antibody (50,000-100,000 cpm per well) and incubated the wells at 4°C overnight. We then washed the wells four

3679

89085519

-30-

times and separated each well for bound ^{125}I detection in a Beckman gamma detect r.

As shown in Figure 1, in which values were plotted following subtraction for background, the peak fraction of solubilized native T4 protein detected by radioimmunoassay coincided with elution of the 55 Kd protein seen by silver staining.

Western Blot Assay For T4

Although many antibodies have been developed for detecting T4 antigen, none are useful for protein blot analysis (Dr. Ellis Reinherz, personal communication). In order to develop antibodies useful for Western blot detection of soluble T4 to follow the purification of T4 and recombinant soluble T4, we raised polyclonal, hyperimmune anti-T4 antisera in rabbits against three synthetic T4 oligopeptides. These oligopeptides are represented in Figure 3 as follows:

	<u>Oligopeptide</u>	<u>Amino Acid Coordinates</u>
20	JB-1	44-63
	JB-2	133-156
	JB-3	325-343

We had previously synthesized these peptides using conventional phosphoamide DNA synthesis techniques. See, e.g., Tetrahedron Letters, 22, pp. 1859-62 (1981). We synthesized the peptides on an Applied Biosystems 380A DNA Synthesizer and purified them by gel electrophoresis.

(i) Coupling Of T4 Peptides To BTG

We coupled each of these peptides to the carrier protein bovine thyroglobulin ("BTG") [Sigma, St. Louis, Missouri] according to a modification of procedures set forth in J. Rothbard et al., J. Exp. Med., 160, pp. 208-21 (1984) and R. C. Kennedy et al., "Antiserum To A Synthetic Peptide Recognizes The

89085519

-31-

HTLV-III Envelope Glycoprotein", Science, 231, pp. 1556-59 (1986).

More specifically, we mixed 10 mg of BTG diluted in 1 ml of PBS with 1.3 mg of m-maleimido-benzoyl-N-hydroxysuccinimide ester ("MBS") in 0.5 ml of dimethylformamide ("DMF"). We mixed the reaction mixture well and reacted it for about 1 hour at 25°C. Subsequently, we loaded the mixture onto a Sephadex G25 gel filtration column (Pharmacia, Sweden) which had been pre-equilibrated with 0.1 M PBS (pH 6.0). We then collected a total of thirty 2 ml aliquot elution fractions and read the absorbance of each fraction at 280 nm (" A_{280} "). We then pooled the three peak fractions (15, 16 and 17) to create the activated carrier.

We dissolved 10 mg of NaBH_4 in 2.5 ml of 0.1 M sodium borate solution to produce a sodium borohydride solution. Subsequently, we diluted approximately 8 mg of each of synthetic T4 peptides JB-1, JB-2 and JB-3 with 1 ml of 0.1 M borate buffer and then mixed each solution with 200 μl of the sodium borohydride solution, incubating the mixture on ice for 5 minutes. We then warmed each peptide solution to 25°C, brought each solution to pH 1.0 with 1 N HCl (during which frothing occurred) and then brought each solution to pH 7.0 with 1 N NaOH (after the frothing had stopped).

We then coupled each peptide to BTG by adding 1.2 ml of the peptide solution to 6 ml of the activated carrier solution. We allowed the coupling reaction to proceed overnight by incubating the reaction mixture at room temperature.

(ii) Inoculation Of Test Animals

We dissolved each of the BTG-coupled peptides prepared above in sterile Freund's complete adjuvant, to a final concentration of 1 $\mu\text{g}/\text{ml}$ coupled

-32-

peptide in PBS. Subsequently, we inoculated each of three rabbits (New Zealand white) by intramuscular injection of 500 µg of one of the coupled peptides into each rabbit. We inoculated a fourth rabbit
5 (New Zealand white) in the same manner with a mixture of the three coupled peptides. All rabbits were prebled prior to boosting to establish an average baseline for each response to be measured. The rabbits were boosted at 6 weeks with 500 µg coupled
10 peptide in incomplete Freund's adjuvant.

Serum was collected from each rabbit monthly for 4 months after immunization. The serum was then assayed for antipeptide titer.

15 (iii) ELISA With Antipeptide Sera
Against Peptide Coated Plates

In this assay, we determined that antiserum raised in an animal by each of peptides JB-1, JB-2 and JB-3 binds to that peptide. Accordingly, those peptides are immunogenic and elicit a response in
20 test animals.

To carry out the assay, we coated Immulon-2 (Dynatech Labs, Alexandria, Virginia) microtiter plates with 50 µl per well of 50 µg/ml uncoupled peptide in PBS and incubated the plates overnight at
25 4°C. Plates coated with peptide 46R*, which served as controls, were treated identically. We then washed the plates 4 times with PBS-Tween (0.5%) and 4 times with water. The plates were blotted dry by gentle tapping over paper towels. After blotting the plates,

30

* Peptide 46 corresponds to amino acids ("AA") 728-751 of the env gene of the HIV genome. The amino acid numbering corresponds to that set forth for the env gene in L. Ratner et al., "Complete Nucleotide
35 Sequence Of The AIDS Virus, HTLV-III", Nature, 313, pp. 277-84 (1985). Peptide 46 has the sequence:

LP I P R G P D R P E G I E E E G G E R D R D R.

3682

89085519

-33-

we added 200 μ l. of a 5% FCS/PBS solution to each well and incubated the plates for 1 hour at room temperature.

We then assayed serum samples from the rabbits on the pre-coated plates prepared as described above. We assayed the antibody response to the immunogen peptide at an initial dilution of 1:100, followed by serial 10-fold dilutions in 5% FCS/PBS.

After a 2 hour incubation period at room temperature, we washed the plates and blotted them dry as described above. We then added 50 μ l of a 1:1500 dilution of horseradish peroxidase ("HRP")-conjugated goat anti-rabbit-IgG [Cooper Biomedical, Malvern, Pennsylvania] in 5% FCS/PBS to each well and incubated the plates at room temperature for 1 hour. We washed the plates with PBS-Tween 0.5%. We then added 50 μ l of 0.42 mM TMB. We stopped the enzyme reactions with 50 μ l of 2 M H_2SO_4 . We then analyzed the plates spectrophotometrically at 450 nm using a microtiter plate reader [Dynatech Labs, Alexandria, Virginia].

We observed that antiserum against each of peptides JB-1, JB-2 and JB-3 binds to the corresponding peptide. We also observed that antiserum against a mixture of peptides JB-1, JB-2 and JB-3 binds to peptides JB-1 and JB-3 under the conditions set forth above. The titers of each of the four antisera tested against the peptides in the solid-phase ELISA are shown below, where "ND" represents values not determined:

Peptide	Approximate Titer Against:		
	JB-1	JB-2	JB-3
JB-1	>1/50,000	0	ND
JB-2	0	1/50,000	ND
JB-3	0	0	1/10,000
JB-1 + JB-2 + JB-3	1/4,000	ND	1/7,000

-34-

Ig fractions from two of the three anti-peptide sera raised against individual peptides, anti-JB-1 and anti-JB-2, recognized the 55 Kd T4 antigen band of native solubilized T4 in a Western blot analysis of protein eluted from the 19Thy (anti-T4) monoclonal antibody affinity column described above. As in the case of the radioimmunoassay of native solubilized T4, the detection of the 55 Kd protein coincides with its apparent elution from the affinity column. This provides further evidence that our T4 purification procedure enriched for solubilized T4.

Thus, these polyclonal sera are useful in the detection of nanogram quantities of T4 (both native and recombinant forms) by Western analysis.

Binding of Cell-Free T4 To HIV Envelope

We then tested our purified solubilized native T4 isolated from U937 cells for its ability to bind to the HIV envelope protein gp160/gp120. To carry out this direct binding assay, we incubated ³⁵S-labelled gp160/gp120 detergent cell extract derived from a recombinant cell line 7d2 (a gift from Drs. Mark Kowalski and William Haseltine, Dana-Farber Cancer Institute) with samples of solubilized native T4, each of which had been preincubated with one type of monoclonal antibody.

More specifically, we mixed 5 µl of solubilized T4 in a microfuge tube with 5 µg (about 3 µl) of OKT4 (ATCC #CRL 8002), a monoclonal antibody recognizing an epitope on T4 which does not interfere with HIV binding [J. A. Hoxie et al., J. Immunol., 136, pp. 361-63 (1986)] or with 5 µg of OKT4A (Ortho Diagnostics #7142), a monoclonal antibody that interferes with HIV binding to T4 positive cells [J. Steven McDougal et al., J. Immunol., 137, pp. 2937-2944 (1986)]. Alternatively, we mixed 50 µl of solubilized

-35-

T4 with 5 µg of αHTLV III gp120 (Dupont #NEN-9284). We then incubated the mixtures on ice for 1 hour.

Subsequently, we added 150 µl of ³⁵S-labelled gp160/gp120 cell extract or ³⁵S-labelled control cell extract (precleared with protein-A Sepharose) to the preincubated solubilized T4/monoclonal antibody mixtures and rocked the tubes overnight at 4°C. We then precipitated the T4/gp160/gp120 immune complexes by adding 30 µl of protein-A Sepharose to each tube and rocking for 2 hours at 4°C to allow the protein-A Sepharose to bind to the antibody complexes. Subsequently, we spun down the beads in an Eppendorf microfuge and after extensive washings, we eluted with 40 µl SDS sample buffer at 65°C for 10 minutes. We then loaded 20 µl of the eluted material on a 7.5% SDS-PAGE gel which was run under reducing conditions.

Figure 2 depicts autoradiograph and Western blot results of the T4/gp160/gp120 coimmunoprecipitations. In Figure 2, lanes 1-5 were autoradiographed after treatment with 40% sodium salicylate and lanes 6-7 were developed on a Western blot with rabbit antisera JB-2.

As shown in Figure 2, gp160/gp120 protein was coimmunoprecipitated in the presence of T4 with OKT4 (lane 5) but not in the presence of T4 with OKT4A (lane 4). Lane 3 shows the positive control for gp160/gp120 using αHTLV III gp120 monoclonal antibody. Neither negative control with ³⁵S-labelled control extract (lane 1) or protein-A Sepharose alone (lane 2) showed bands migrating in the position of gp160/gp120. Based upon the bands that developed on the Western blot, the amount of T4 precipitated with either OKT4 (lane 6) or OKT4A (lane 7) appeared to be similar.

This demonstrates that purified, solubilized native T4, which is naturally membrane bound, can

3685

89085510

-36-

still interact with the HIV glycoprotein in solution. Accordingly, we believe that cell free soluble T4 is useful in preventing the binding interaction between HIV and the T4 receptor of T4⁺ lymphocytes. By competing with cell surface T4 for binding to the HIV envelope protein gp120, soluble T4 is useful in blocking HIV infection.

Synthesis Of Oligonucleotide DNA Probes

The nucleotide sequence and a deduced amino acid sequence for a cDNA that purportedly encodes the entire human T4 protein have been reported [Maddon et al., (1985), supra]. The deduced primary structure of the T4 protein reveals that it can be divided into domains as demonstrated below:

15	<u>Structure/Proposed Location</u>	<u>Amino Acid Coordinates</u>
	Hydrophobic/Secretory Signal	-23 to -1
	Homology to V-Regions/ Extracellular	+1 to +94
20	Homology to J-Regions/ Extracellular	+95 to +109
	Glycosylated Region/ Extracellular	+110 to +374
25	Hydrophobic/Transmembrane Sequence	+375 to +395
	Very Hydrophilic/ Intracytoplasmic	+396 to +435

Based on the sequence for the above-listed domains, we chemically synthesized antisense oligonucleotide DNA probes using conventional phosphoamide DNA synthesis techniques. See, e.g., Tetrahedron Letters, 22, pp. 1859-62 (1981). We synthesized the probes on an Applied Biosystems 380A DNA synthesizer and purified them by gel electrophoresis.

89085519

-37-

Furthermore, we synthesized the probes such that they were complementary to the DNA sequences which code for the amino acid sequence, i.e., the probes were antisense, to enable them to recognize and hybridize to the corresponding sequences in DNA, as well as in mRNA. The nucleotide sequences of the eleven selected regions of the T4 protein [corresponding to the nucleotide numbering set forth in Maddon et al., (1985), *supra*] were the following:

	<u>Oligonucleotide</u>	<u>Nucleotide Coordinates</u>
10	1	145-171
	2	742-765
	3	1414-1440
15	6	427-453
	7	1303-1329
	8	1012-1038
	9	97-118
	10	10-36
20	11	1698-1724
	12	397-423
	14	261-287

Before using our DNA probes for screening, we 5' end-labelled each of the single-stranded DNA probes with ^{32}P using $[\gamma\text{-}^{32}\text{P}]\text{-ATP}$ and T4 polynucleotide kinase, substantially as described by A. M. Maxam and W. Gilbert, "A New Method For Sequencing DNA", Proc. Natl. Acad. Sci. USA, 74, pp. 560-64 (1977).

30 Construction of λ gt10 Peripheral Blood Lymphocytes cDNA Library

To prepare our Peripheral Blood Lymphocytes (PBL) cDNA library, we processed PBL, from a single leukophoresis donor, through one round of absorption

3687

89085519

-38-

to remove monocytes. We then stimulated the non-adherent cells with IFN- γ 1000 μ /ml and 10 μ g/ml PHA for 24 hours. We isolated RNA from these cells using phenol extraction [Maniatis et al., Molecular Cloning,
5 p. 187 (Cold Spring Harbor Laboratory) (1982)] and prepared poly A⁺ mRNA by one round of oligo dT cellulose chromatography. We ethanol precipitated the RNA, dried it in a speed vac and resuspended the RNA in 10 μ l H₂O (0.5 μ g/ μ l). We treated the RNA for 10
10 min at room temperature in CH₃COOH (5 mM final concentration) and β -mercaptoethanol (0.26 M). We then added the methyl mercury treated RNA to 0.1 M Tris-HCl (pH 8.3) at 43°C, 0.01 M Mg, 0.01 M DTT, 2 mM Vanadyl complex, 5 μ g oligo dT₁₂₋₁₈, 20 mM KCl, 1 mM dCTP,
15 dGTP, dTTP, 0.5 mM dATP, 2 μ Ci[α -³²P]dATP and 30 U 1.5 μ l AMV reverse transcriptase (Seikagaku America) in a total volume of 50 μ l. We incubated the mixture for 3 minutes at room temperature and then for 3 hours at 44°C, after which time we stopped the reaction by
20 the addition of 2.5 μ l of 0.5 M EDTA.

We extracted the reaction mixture with an equal volume of phenol:chloroform (1:1) and precipitated the aqueous layer two times with 0.2 volume of 10 M NH₄AC and 2.5 volumes EtOH and dried it under
25 vacuum. The yield of cDNA was 1.5 μ g.

We synthesized the second strand according to the methods of Okayama and Berg [Mol. Cell. Biol., 2, p. 161 (1982)] and Gubler and Hoffman [Gene, 25, pp. 263-69 (1983)], except that we used the DNA polymerase I large fragment in the synthesis.
30

We blunt ended the double-stranded cDNA by resuspending the DNA in 80 μ l TA buffer (0.033 M Tris Acetate (pH 7.8); 0.066 M KAcetate; 0.01 M MgAcetate; 0.001M DTT; 50 μ g/ml BSA), 5 μ g RNase A, 4 units RNase
35 H, 50 μ M β NAD, 8 units E.coli ligase, 0.3125 mM dATP, dCTP, dGTP, and dTTP, 12 units T₄ polymerase and incubated the reaction mixture for 90 min at

3688

89085519

-39-

37°C, added 1/20 volume of 0.5M EDTA, and extracted with phenol:chloroform. We chromatographed the aqueous layer on a G150 Sephadex column in 0.01M Tris-HCl (pH 7.5), 0.1 M NaCl, 0.001 M EDTA and
5 collected the lead peak containing the double-stranded cDNA and ethanol precipitated it. Yield: 0.605 µg cDNA.

We ligated the double-stranded cDNA to linker 35/36:

10 5'AATTCTGAGCTCGAGCGCGGCCTGC3'
3' GCTCTGAGCTCGCGCCGGCG5'

using standard procedures. We then size selected the cDNA for 800 bp and longer fragments on a S500 Sephacryl column, and ligated it to EcoRI-digested
15 bacteriophage lambda vector gt10 (a gift of Dr. Ellis Reinherz). We packaged aliquots of the ligation reaction in Gigapak (Stratagene) according to the manufacturer's protocol. We used the packaged phage to infect E.coli BNN102 cells and plated the
20 cells for amplification. The resulting library contained 1.125×10^6 independent recombinants.

We also screened a PBL cDNA library in the bacteriophage lambda vector gt10 (a gift of Dr. Ellis Reinherz), which was synthesized from mRNA from a
25 T4⁺ tumor cell line named REX, which expresses T4 protein at high levels [O. Acuto et al., "The Human T Cell Receptor: Appearance In Ontogeny And Biochemical Relationship Of Lambda and Beta Subunits on IL-2 Dependent Clones And T Cell Tumors", Cell,
30 34, pp. 717-26 (1983)].

Screening Of The Libraries

We then used three of our ³²P-labelled synthetic oligonucleotide antisense probes, probes 3, 6 and 9, to screen in parallel our two λgt10 cDNA
35 libraries using the plaque hybridization screening technique described in R. Cate et al., "Isolation Of

3689

89085519

The Bovine And Human Genes For Mullerian Inhibiting Substance And Expression Of The Human Gene In Animal Cells", Cell, 45, pp. 685-98 (1986), with minor modifications. We modified the Cate et al. procedure by hybridizing without tetramethyl ammonium chloride to accommodate our use of unique probes, rather than mixtures, to probe the plaque filters.

We used the three probes, which had been previously 5' end-labelled with [γ -³²P]-ATP according to the method of A. Maxam and W. Gilbert, Meth. Enzymol., 68, pp. 499-80 (1979) to screen in parallel the PBL cDNA library and the REX cDNA library discussed above.

From our screening of the PBL library, we isolated a nearly full length soluble T4 cDNA clone -- λ 203-4 (or λ gt10.PBL.T4) -- containing a 3.064 kb insert which could be cleaved from the λ gt10 vector with EcoRI.

From our screening of the REX cell library, we isolated an incomplete T4 cDNA clone containing a 1,200 bp cDNA insert. We then further characterized the DNA from these clones by DNA sequencing analysis.

We also screened a bacteriophage lambda human genomic library, constructed in the vector EMBL3 by Dr. Mark Pasek (Biogen Inc., Cambridge, Massachusetts) [N. Murray in Lambda 2, eds. R. Hendrix, J. Roberts, F. Stahl, R. Weisberg, pp. 3935-422 (1983)]. The library contains DNA fragments, created by partial restriction of chromosomal DNA from the human lymphoblastoid cell line GM1416,48, XXXX (Human Genetic Mutant Cell Repository, Camden, New Jersey) with Sau3a, ligated onto EMBL3 arms which had been subjected to cleavage with BamHI according to the procedures outlined in Maniatis et al., (1982), supra. Plating of the phage library, lysis, and transfer of the phage DNA onto nitrocellulose were performed as described by W. D. Benton and R. W. David, "Screening

-41-

f Lambda gt Recombinant Clones By Hybridization To Single Plaques In Situ", Science, 196, p. 180 (1977) and Maniatis et al. (1982). Hybridization conditions were those described by Cate et al. (1986), supra, except that tetramethylammonium chloride (TMACl) was omitted from the washing buffer.

Approximately 2 million plaques were screened in parallel hybridizations with probe 1 and probe 3 discussed above. One phage, called CM47, which hybridized with probe 3 in the primary screenings, was subjected to DNA sequence analysis to determine the existence and position of an intron between the coding sequences for the predicted extracellular and transmembrane domains. No phage clones containing T4 sequences were found screening with probe 1, probably because it includes a sequence interrupted by an intron [D. R. Littman and S. N. Gettner, Nature, 325, pp. 453-55 (1987); and our observations].

Partial sequence analysis of CM47 shows that an intron interrupts the sequence corresponding to the codon for valine (amino acid 363) of the deduced primary sequence for T4 (Figure 3 -- in which introns are indicated by a solid line). This intron defines a potential site for introducing a stop codon in order to express a soluble form of T4. Another intron found within the coding sequence for T4 interrupts the codon for arginine (amino acid 295) and a third intron in CM47 is found between the codons for arginine (amino acid 402) and arginine (amino acid 403) (Figure 3).

Sequencing Of cDNA Clones

We then subcloned EcoRI digested DNA from clone λ 203-4 into animal expression vector pBG312 [R. Cate et al., supra] to facilitate sequence analysis. More specifically, as depicted in Figure 4,

3691

89085519

-42-

we then digested λ gt10.PBL.T4 with EcoRI to excise the 3.064 kbp EcoRI-EcoRI fragment containing the full length T4 cDNA. This cDNA sequence, including the entire coding region for soluble T4 and for full length T4 was deposited in p170-2. We used T4 ligase to ligate the fragment into animal expression vector pBG312 [supra] which had been previously cut with EcoRI, to form pBG312.T4 and p170-2 (Figure 4). We then determined the nucleotide sequence of the EcoRI fragment of pBG312.T4 using Maxam Gilbert technology [A. M. Maxam and W. Gilbert, "A New Method For Sequencing DNA", Proc. Natl. Acad. Sci. USA, 74, pp. 560-64 (1977)] (see Figure 3, which depicts the PBL cDNA sequence in comparison to that reported by Maddon et al., (1985), supra). This analysis showed that the 3.064 kbp PBL full length complementary DNA copy of T4 cDNA contained the coding sequence for T4, approximately 200 bp of 5' noncoding sequence and approximately 1500 bp of 3' noncoding sequence.

We then cut pBG312.T4 with PstI and removed the resulting 3' protruding ends with Klenow and isolated an approximately 2.5 kbp fragment. We then inserted the fragment into the polylinker of pBG312 (which had been previously restricted at the SmaI site) to form plasmid p170-2, which contains the full length PBL T4 cDNA sequence (see Figure 3).

As depicted in Figure 3, the PBL T4 cDNA contains a nucleotide sequence almost identical to the approximately 1,700 bp sequence reported by Maddon et al., (1985), supra. The PBL T4 cDNA, however, contains three nucleotide substitutions that, in the translation product of this cDNA, would produce a protein containing three amino acid substitutions compared to the sequence reported by Maddon et al. As shown in Figure 3, these differences are at amino acid position 3, where the asparagine of Maddon et al. is replaced with lysine; position 64,

00005510

-43-

where the tryptophan of Maddon et al. is replaced with arginine and at position 231, where the phenylalanine of Maddon et al. is replaced with serine. The asparagine reported at position 3 of Maddon et al. instead of lysine was the result of a sequencing error (Dr. Richard Axel, personal communication). The significance of the amino acid replacements at positions 64 and 231, which may represent allelic polymorphism [T. C. Fuller et al., Human Immunology, 9, pp. 89-102 (1982); W. Stohl and H. G. Kunkel, Scand. J. Immunol., 20, pp. 273-78 (1984); N. Amino et al., Lancet, 2, pp. 94-95 (1984); and M. Sato et al., J. Immunol., 132, pp. 1071-73 (1984)], is not known.

DNA sequence analysis [Maxam and Gilbert, supra] of the insert in pEC100 of the REX clone suggests that it represents the product of a splicing error, because 5' noncoding sequence appears to have been spliced with coding sequence beginning with the GGT codon for glycine (amino acid 49) (see Figure 3 and Figure 5). The T4 coding sequence in pEC100* from glycine (amino acid 49) to isoleucine (amino acid 435) is identical to the sequence of Maddon et al., (1985), supra.

In comparison, our earlier N-terminal protein sequence analysis of native T4 protein purified from U937 cells shows a T4 expression product with asparagine as amino acid 3. These differences are also set forth in Figure 6, which also depicts comparisons at corresponding positions of the partial clone from the REX cell line λ gt10 library; our

* We constructed pEC100 by digesting the incomplete T4 cDNA clone from the REX library with EcoRI and isolating the 1,200 bp cDNA insert. We then ligated it to pUC12 (Boehringer Mannheim, Indianapolis, Indiana) which had been previously cut with EcoRI to form pEC100.

-44-

genomic clone from a λ EMBL3 library; mouse T4 sequences [Tourvieille et al., Science, 234, p. 610 (1986)] and sheep T4 sequences [Classon et al., Immunogenetics, 23, p. 129 (1986)].

5 Construction of Soluble T4 Mutants

We then employed the technique of in vitro site-directed mutagenesis and restriction fragment substitution to modify the T4 cDNA coding sequence of p170-2 in sequential steps to be identical to
10 that reported by Maddon et al., (1985), supra. We first used oligonucleotide-directed mutagenesis to modify the amino acids at positions 3 and 64. Next, we employed restriction fragment substitution with a fragment including the serine 231 codon of a partial
15 T4 cDNA isolated from a T4 positive lymphocyte cell line [O. Acuto et al., Cell, 34, pp. 717-26 (1983)] library in λ gt11 (a gift from Dr. Ellis Reinherz), to modify the amino acid at position 231. We then truncated our modified T4 cDNA sequence to remove
20 the coding regions for the transmembrane and intracytoplasmic domains. Subsequently, we constructed three different soluble T4 mutants from our full length T4 clone PBL T4 by linker insertion between restriction sites in order to increase the probability
25 of empirically finding a stable, secretable T4 molecule. The structure of each of these mutants is depicted in Figure 7A.

Line A of Figure 7A represents a hydropathy analysis of our full length soluble T4 carried out
30 using a computer program called Pepplot (University of Wisconsin Genetics Computer Group) according to J. Kyte and R. F. Doolittle, J. Mol. Biol., 157, pp. 105-32 (1982). Line B depicts the protein domain structure of full length T4 [Maddon et al., (1985)
35 supra] in which "S" represents the secretory signal sequence, "V" represents the immunoglobulin-like

variable region sequence, "J" represents the immunoglobulin-like joining region sequence, "U" represents the unique, extracellular region sequence, "TM" represents the transmembrane sequence and "C" represents the cytoplasmic region sequence. In line B, the transmembrane amino acid sequence and some flanking sequence is written below the TM domain. Line C depicts the protein domain structure of recombinant soluble T4 mutants rsT4.1 in pBG377, rsT4.2 in pBG380 and rsT4.3 in pBG381. Line D represents the protein domain structure of E.coli rsT4 gene (Met-perfect construct) (p199-7) which is deleted for the T4 N-terminal signal sequence (S).

We constructed the first three soluble T4 mutant gene fragments by truncating our full length soluble T4 cDNA at positions corresponding to either intron/exon boundaries or to protein domain boundaries defined by hydropathy analysis predictions. More specifically, we introduced synthetic linkers into the unique AvaI site that is 5' to the transmembrane/extracellular domain boundary to produce an in-frame translational stop codon, thus constructing T4 genes that lack the transmembrane and cytoplasmic domains of the full length T4 sequence.

For example, mutant rsT4.1 in pBG377 was truncated by the insertion of a stop codon following amino acid 362, lysine, which corresponds to the position of an intron separating the extracellular and transmembrane domain exons. The positions both of this intron and of the adjacent intron that splits the transmembrane and cytoplasmic domains were determined by DNA sequence analysis of chromosomal T4 clones isolated from the λ EMBL3 genomic library described above. Although the significance of the intron positions flanking the T4 transmembrane domain is not known, the determination of the genetic structure could provide important information for design-

-46-

ing rsT4 mutants, since exons frequently define functional domains [W. Gilbert, "Why Genes In Pieces?", Nature, 271, p. 501 (1978)].

We then constructed mutant rsT4.2 in pBG380
5 by truncating the T4 cDNA at the boundary of the transmembrane and extracellular domains at amino acid 374. And, we constructed mutant rsT4.3 in pBG381 by truncating the T4 cDNA at amino acid 377, three amino acids downstream from the transmembrane/
10 extracellular domain boundary and within the transmembrane domain.

We also employed the technique of oligonucleotide site directed mutagenesis, according to D. Strauss et al., "Active Site Of Triosephosphate
15 Isomerase: In Vitro Mutagenesis And Characterization Of An Altered Enzyme", Proc. Natl. Acad. Sci. USA, 82, pp. 2272-76 (1985), to construct a fourth soluble T4 mutant from our full length T4 clone PBL T4. The structure of this mutant is depicted in Figure 7A,
20 line D, which represents the protein domain structure of E.coli rsT4 gene (Met-perfect rsT4.2) construct, deposited in pl99-7, which is deleted for the T4 N-terminal signal sequence (S).

We also constructed various other soluble
25 T4 deletion mutants to determine which smaller fragments of the T4 sequence provide a protein which binds to HIV. These constructions were based on our belief that only the amino terminal sequence of T4 is required for binding to HIV. This belief, in turn,
30 was based upon observations that the monoclonal antibody OKT4A blocks infection of T4 positive cells by HIV and it appears to recognize an epitope in the amino portion of T4 [Fuller et al., supra]. Such fragments of T4, which lack glycosylation and which
35 are capable of binding HIV and blocking infection, may be produced in E.coli or chemically synthesized.

-47-

The structure of each of these deletion mutants is depicted in Figure 7B. In that figure, line A depicts the protein domain structure of full length T4 [Maddon et al., (1985), supra; Figure 7A].

5 In line B, the protein structure of recombinant soluble T4 mutants are depicted as follows: rsT4.7 in p203-5, rsT4.7 in pBG392, rsT4.8 in pBG393, rsT4.9 in pBG394, rsT4.10 in pBG395, rsT4.11 in pBG397, rsT4.12 in pBG396, rsT4.111 in pBG215-7, rsT4.113.1
10 in pBG211-11 and rsT4.113.2 in pBG214-10.

We constructed soluble T4 derivatives p203-5, pBG392, pBG393, pBG394 and pBG396 by truncating our rsT4.2 gene after the StuI sites at amino acids 183 and 264 of rsT4.2. More specifically, we
15 constructed derivative rsT4.7 in p203-5 and in pBG392 by truncating the rsT4.2 cDNA at amino acid 182. And, we constructed each of derivatives rsT4.9 in pBG394 and rsT4.12 in pBG396 by truncating the rsT4.2 cDNA at amino acids 113, and 166, respectively. One
20 may also construct each of derivatives rsT4.10 in pBG395 and rsT4.11 in pBG397 by truncating the rsT4.2 cDNA at amino acids 131 and 145, respectively.

Expression of T4 and Soluble T4 Polypeptides In Bacterial Cells

25 The cDNA sequences of this invention can be used to transform eukaryotic and prokaryotic host cells by techniques well known in the art to produce recombinant soluble T4 polypeptides in clinically and commercially useful amounts.

30 For example, we constructed expression vector p199-7, as shown in Figure 9A, as follows.

We preceded the construction depicted in Figure 9A by the construction of various intermediate plasmids, as depicted in Figures 8A-8D. Those con-
35 structions were carried out using conventional

3697

89085519

-48-

recombinant techniques. The linkers employed in those constructions are set forth in Figure 10.

As depicted in Figures 8A and 8B, starting with p170-2, which contains our full length T4 DNA sequence, coding for T4 characterized by three different amino acids than that of Maddon et al., (1985), supra, we produced various constructs which direct the expression of soluble T4. Some of these constructs are characterized in that one or more of those amino acid differences have been changed to correspond to the respective amino acids of Maddon et al. In this figure, as well as in the other figures, amino acid changes are reflected by an arrow.

Plasmid p192-6 contains the Met perfect rsT4.2 sequence derived by oligonucleotide site-directed mutagenesis which removed the entire T4 N-terminal signal sequence as shown in Figure 8C. And, to provide a convenient means of transferring the rsT4.2 Met perfect sequence into E.coli expression vectors, the steps described in Figure 8D were carried out to produce p195-8, a plasmid containing the Met perfect rsT4.2 sequence flanked by ClaI restriction sites. The ClaI-ClaI cassette of p195-8 optimizes the distance between the 5' ClaI site and the initiating Met codon. In Figure 8D, ST8 rop⁻ is a tetracycline resistance encoding pAT153-based plasmid containing the rop⁻ mutation that permits high plasmid copy number, a promoter and ribosome binding site from bacteriophage gene 32 and the gene 32 transcription termination sequence.

Cleavage of p195-8 with ClaI produced the fragment used to assemble p199-7, a construction which directs the expression of Met perfect rsT4.2 under the control of the P_L promoter (Figure 9A). As the first step, to construct a vector from which rsT4.2 expression is under control of the P_L promoter,

-49-

we constructed the vector p197-1² from p1034 (plmuGCSF) (Figure 9A).

We then cut p1034 with EcoRI and BamHI to excise the GCSF cDNA insert and a portion of the phage mu ribosome binding site sequence -- which we subsequently reconstructed with oligonucleotides. The synthetic linkers used were linkers 57-60 (Figure 10).

We then ligated the synthetic linker into the EcoRI/BamHI-cut p1034 to form p197-12. One could, instead, replace these steps by starting with any suitable E.coli expression vector containing a ClaI site appropriately placed between the promoter and terminator sequences. We cut p197-12 with ClaI and inserted a ClaI-ClaI cassette containing the cDNA sequence of rsT4.3 in pBG381 and phage transcription terminator derived from p1034. The sequence of this cassette is depicted in Figure 11. The resulting plasmid, p199-7, contains the rsT4.2 "Met perfect" gene in that vector.

Alternatively, one could derive the Met perfect rsT4.2 sequence from plasmid pBG380, deposited in connection with this application, and gap out the signal sequence to create p192-6.

We tested for expression of p199-7 as follows. SG936, an E.coli lon htp_r double mutant [ATCC 39624] [S. Goff and A. Goldberg, "ATP-Dependent Protein Degradation In E.coli", in Maximizing Gene Expression, W. Reznikoff and L. Gold (eds.) (1986)], was transformed with p199-7 by conventional procedures [Maniatis et al. (1982)] to form SG936/p199-7, a transformant containing a plasmid with the Met-perfect rsT4.2 gene behind the P_L promoter. Transformants were selected on LB agar plates containing 10 mcg/ml tetracycline (tet). After streaking out several single colonies for single colony isolation, one was chosen at random for testing induction f

3699

89085519

rsT4.2 synthesis. We picked a single colony from an LB-agar tet⁺ plate into 20 ml Luria Broth (LB) and 10 mcg/ml tet in a 125 ml shake flask and grew it overnight in a shaking air incubator (New Brunswick Scientific, New Jersey) at 30°C.

We then initiated an induction culture by adding 0.5 ml of the overnight culture to 50 ml LB and tet in a 500 ml flask which was grown at 30°C in a shaking air incubator. When the culture reached an OD(600) of 0.4, we transferred it to a 42°C water-bath and shook it gently for approximately 20 minutes. After heat induction at 42°C, the flask was transferred to a 39°C air incubator (New Brunswick Scientific, New Jersey) where it was shaken vigorously at 250 rpm. We withdrew samples just after the 42°C heat shock, and at hourly time points for 4 hours, and then after overnight growth. The samples were measured for growth by OD(600) and analyzed following SDS-PAGE for the pattern of protein synthesis by Coomassie blue protein staining and by Western blot analysis with our rabbit anti-peptide antibody probes (described above). Based on the relative molecular weight and protein blot analysis, the expression of rsT4.2 was induced from SG936/p199-7 following heat induction at 42°C (Figure 12).

We transformed p199-7 into a P_Lmu.tet expression vector, an E.coli expression vector, at the unique ClaI site (see Figure 11). The nucleotide and amino acid sequences of p199-7 are shown in Figure 11.

The expression of soluble T4 from p199-7 in E.coli was measured by Western blot analysis of whole cell extracts following SDS-PAGE using the rabbit polyclonal anti-peptide JB-1 or anti-peptide JB-2 antibodies as probes (Figure 12).

We also constructed expression vector p203-5, as shown in Figure 9B, as follows.

We started with p197-7, which has the same sequence as the P_{Lum} vector p197-1 (see Figure 9A), except that there is a single nucleotide deletion in the 5' noncoding region following the P_L promoter. That deletion, which is a deletion of nucleotide #40 -- adenine -- of p197-12 (see Figure 11), resulted from a deletion in the region that was constructed from linkers 57-60 (see Figure 10). p197-7 contains the rsT4.2 gene comprising 374 amino acids. Alternatively, one could also use p197-7 as a starting plasmid.

We cut p197-7 with ClaI. We also cut p195-8 (see Figures 8D and 9A) with ClaI to remove the ClaI - ClaI cassette containing the cDNA sequence of rsT4.2. Subsequently, we inserted the ClaI-ClaI cassette into p197-7 to produce p198-2.

We then digested p198-2 with StuI to remove 80 amino acids (amino acid 185 to amino acid 264) of the mature T4 protein coding sequence. Unexpected methylation, however, prevented cutting at the second StuI site, so that only the StuI site at amino acid 184 was cleaved. Following ligation, the plasmid DNA was transformed into E.coli and we examined several plasmid clones for the deletion using standard procedures. None of those plasmids contained the expected StuI deletion.

Subsequent DNA sequence analysis of one of these plasmids, called p203-5, showed that two guanine residues (see amino acids 183 and 184; nucleotides 818 and 819 of Figure 3) of the StuI recognition sequence had been deleted following cleavage due to exonuclease digestion caused by the use of exonuclease-contaminated StuI enzyme. This dinucleotide deletion produced a translation frameshift following amino acid 182 (glutamine) and introduced a stop codon six amino acid codons downstream from the frameshift (Figure 9C). The unexpected

-52-

methylation of the second StuI site together with the deletion that resulted in a few stop codon produced a gene encoding a shortened form of recombinant soluble T4, called rsT4.7. The rsT4.7 sequence
5 encodes a 182 amino acid N-terminal segment of the mature T4 sequence followed by, at the C-terminus, six amino acids -- asparagine-leucine-glutamine-histidine-serine-leucine -- of non-T4 sequence and finally by a TAA stop codon.

10 The expression of soluble T4 from p203-5 in E.coli was measured by Western blot analysis as previously described.

Expression of T4 and Soluble T4
Polypeptides In Animal Cells

15 We inserted both soluble T4 genes and the unmodified gene encoding membrane-bound T4 into animal expression vector pBG368. More specifically, we inserted each of the soluble gene constructs into pBG368 under the transcriptional control of the
20 adenovirus late promoter, to give plasmids pBG377, pBG380 and pBG381. We also made two pBG312-based constructions, called pBG378 and pBG379, which direct the expression of recombinant full length T4 protein. pBG378 and pBG379 code for the same full
25 length T4 protein but in pBG379, a portion of the 3' untranslated sequence has been removed. Subsequently, to test for expression of recombinant soluble T4 and recombinant full length T4, we cotransfected Chinese hamster ovary ("CHO") cells with one of each of
30 those plasmids and with the plasmid pAdD26.

We first constructed pBG368 as follows. As depicted in Figure 13, we cut animal cell expression vector pBG312 [R. Cate et al., "Isolation Of The Bovine And Human Genes For Mullerian Inhibiting
35 Substance And Expression Of The Human Gene In Animal Cells", Cell, 45, pp. 685-98 (1986)] with EcoRI and

BglIII to delete one of each of the two EcoRI and the two BglIII restriction sites (the EcoRI site at position 0 and the BglIII site located at approximately position 99). The resulting plasmid, pBG368, retained
5 an EcoRI site in the cloning region and a BglIII site after the cloning region. This left a single EcoRI site and a single BglIII site in the polylinker for cloning purposes.

More specifically, we deleted one EcoRI
10 site and one BglIII site by sequential partial digestion of pBG312 with restriction enzymes EcoRI and BglIII, respectively. We filled in with Klenow and 4 nucleotides then religated to produce pBG368, which contains unique restriction sites for EcoRI and BglIII
15 enzymes.

Once transient expression of soluble T4 was verified, we constructed stable cell lines that continuously expressed soluble T4. To do this, we employed the stable cell expression host, the
20 dihydrofolate reductase deletion mutant (DHFR⁻) Chinese hamster ovary cell line [F. Kao et al., "Genetics Of Somatic Mammalian Cells X Complementation Analysis of Glycine-Requiring Mutants", Proc. Natl. Acad. Sci., 64, pp. 1284-91 (1969); L. Chasin and
25 G. Urlab "Isolation Of Chinese Hamster Cell Mutants Deficient In Dihydrofolate Reductase Activity", Proc. Natl. Acad. Sci., 77, pp. 4216-80 (1980)].

Using this system, we cotransfected each T4 gene construct with pAdD26 [R. J. Kaufman and
30 P. A. Sharp, "Amplification And Expression Of Sequences Cotransfected With a Modular Dihydrofolate Reductase Complementary DNA Gene", J. Mol. Biol., 159, pp. 661-21 (1982) containing the mouse DHFR gene. Before carrying out the co-transfections, we
35 linearized all plasmids by restriction enzyme cleavage and, prior to transfection, we mixed each plasmid with pAdD26 so that the molar ratio of pAdD26 to T4

-54-

was 1:10. This maximized the number of T4 gene copies per transfectant.

Within the cell, the plasmids were ligated together to form polymers that can become integrated into host chromosomal sequences by illegitimate recombination [J. Haynes and C. Weissmann, "Constitutive, Long-Term Production Of Human Interferons By Hamster Cells Containing Multiple Copies Of a Cloned Interferon Gene", Nucl. Acids Res., 11, pp. 687-706 (1983); S. J. Scahill et al., "Expression And Characterization Of The Product Of A Human Immune Interferon cDNA Gene In Chinese Hamster Ovary Cells", Proc. Natl. Acad. Sci. USA, 80, pp. 4654-58 (1983)]. We selected transfectants that express the mouse DHFR gene in culture medium lacking nucleotides. We then subjected these transfectants to a series of increasing concentrations of methotrexate, a toxic folate analogue that binds DHFR, to select for cells levels of DHFR.

Resistance to methotrexate by increased expression of DHFR is frequently the result of DHFR gene amplification, which can include the reiteration of large chromosomal segments, called amplified units [R. J. Kaufman and P. A. Sharp, "Amplification And Expression Of Loss Of Dihydrofolate Reductase Genes In A Chinese Hamster Ovary Cell Line", Molec. Cell. Biol., 1, pp. 1069-76 (1981)]. Therefore, cointegration of DHFR and rsT4 sequences permitted the amplification of rsT4 genes. Stably transfected cell lines were isolated by cloning in selective growth medium, then screened for T4 expression with a T4 antigen (RIA) [D. Klatzmann et al., Nature, 312, pp. 767-68 (1984)] and by immunoprecipitation from conditioned medium after [³⁵S] cysteine ("³⁵S-Cys") metabolic labelling.

We also inserted the soluble T4 derivative rsT4.7 gene into an animal cell expression plasmid as follows.

-55-

As set forth in Figure 14C, we cut plasmid pBG381 (Figure 14A) with EcoRI and NheI. We then cut p186-6 with EcoRI and NheI to remove the 786 base pair fragment. We ligated that fragment into the
5 digested pBG381 to form plasmid pBG391. The T4 sequence in pBG391 is identical to both that of Maddon et al. (1985) supra at positions 64 (tryptophan) and 231 (phenylalanine) and to that of pBG381. However, at position 3, the asparagine reported by
10 Maddon et al. and present in pBG381 is replaced with lysine. The nucleotide sequence of pBG391 is depicted in Figure 15.

We then digested p203-5 with NheI and OxaNI to remove the 483 base pair fragment. We
15 inserted that fragment into NheI/OxaNI-digested pBG391 to form plasmid pBG392, the animal cell expression construct of rsT4.7. The T4 sequence in rsT4.7 contains amino acids identical to that of Maddon et al.'s full length sequence at amino acid
20 positions 64 (tryptophan) and 231 (phenylalanine). However, at position 3, the asparagine reported by Maddon et al. is replaced with lysine. The nucleotide sequence of pBG392 is depicted in Figure 16.

In Figure 14D, we have depicted the construction of other animal cell expression constructs
25 containing sequences encoding the deletions rsT4.9 in pBG394, and rsT4.12 in pBG396. Those constructions were carried out using conventional recombinant techniques. The linkers employed in those constructions
30 are set forth in Figure 18. The nucleotide sequences of pBG394 and pBG396 are shown in Figures 19 and 20.

Plasmid pBG393, shown in Figure 17, contains rsT4.8, the perfect form of rsT4.7. pBG393 contains 182 amino acids of the mature T4 sequence,
35 without the additional non-T4 6 amino acids at the C-terminus following amino acid 182. The nucleotide
3705 sequence of BG393 is shown in Figure 21.

89085519

-56-

Other animal cell expression plasmids according to this invention may be constructed as depicted in Figure 17. These include rsT4.10 in pBG395 and rsT4.11 in pBG397 (see Figure 18 for specific linkers).

The nucleotide sequence of BG395 is shown in Figure 22.

Purification Of Recombinant Soluble T4

Recombinant soluble T4 construct pBG380 expressed in DHFR⁻ CHO cells was grown to confluency in a α -Modified Eagles Medium (Gibco) supplemented with 10% fetal calf serum, 1 mM glutamine and the antibiotics penicillin and streptomycin (100 μ g/ml of each). The cells were grown at 37°C in two 21 Cell Factory Systems (Nunc). We then washed the confluent cells free of fetal calf serum with α -Modified Eagles Medium without fetal calf serum and cultured the cells in α -Modified Eagles Medium at 37°C for 4 days. Subsequently, we harvested the conditioned media, filtered it through a Millipore Millidisk 0.22 μ hydrophilic filter cartridge (Millipore #MCGL 305-01) and concentrated the secreted proteins on a fast-S ion exchange column (S-Sepharose Fast Flow, Pharmacia #17-0511-01) in 20 mM MES buffer (pH 5.5).

We then eluted the bound proteins with 20 mM Tris-HCl (pH 7.7) and 0.3 M NaCl. The elution pool was subsequently diluted with 2 volumes of 20 mM Tris-HCl (pH 7.7) and it was then loaded on a column comprising immobilized 19Thy anti-T4 monoclonal antibody coupled to Affigel-10 [a gift of Dr. Ellis Reinherz, Dana Farber Cancer Institute, Boston, Massachusetts]. We washed the column extensively and eluted the bound material as 0.5 ml fractions with 50 mM glycine-HCl (pH 2.5), 150 mM NaCl, 0.1 mM EGTA and 5 μ g/ml bovine pancreatic trypsin inhibitor, Aprotinin (Sigma #A1153). We used Western blots

-57-

developed with rabbit antisera raised against peptide
JB-2 to follow the purification. We employed silver
stained gels to follow binding and elution of rsT4.2
during the chromatography. Figure 23 depicts a
5 Coomassie stained gel of purified rsT4.2.

Gel sizing-column chromatography analysis
of the purified rsT4.2 from the pBG380 transfected
CHO cell line, BG380G, suggests that rsT4 is monmeric
under physiologic pH and salt concentration.

10 Sequencing Of Recombinant Soluble T4 Protein

We then determined the N-terminal amino
acid sequence of a recombinant soluble T4, specifi-
cally rsT4.2, molecule purified from the conditioned
15 medium of the pBG380 transfected CHO cell line BG80G,
as described above, by automated Edman degradation
in an Applied Biosystems 470A gas phase sequenator
[R. B. Pepinsky et al., J. Biol Chem., 261,
pp. 4239-46 (1986)].

20 The amino terminal sequence matched the
sequence which we had previously determined for
solubilized native T4 isolated from U937 cells, supra.
The amino terminal sequences of native solubilized
T4 (sT4) and purified rsT4 protein are $\Delta 2$ proteins,
25 as compared to the amino terminal sequence predicted
by Maddon et al., (1985), supra, with the mature amino
terminus located at position 3 of that sequence. The
amino terminal sequences of solubilized native T4
(sT4), recombinant soluble T4 (rsT4.2) secreted by
30 CHO transfectant BG380G containing pBG380 and the
protein sequence deduced by Maddon et al. (1985),
supra are as follows:

sT4:	X-K-V-V-L-X-K-K-X-D-T-V-E-L-T-X-T-A-S-E-
rsT4.2:	N-K-V-V-L-G-K-K-G-D-T-V-E-L-T-X-T-A-S-E-

3707

89085519

-58-

Maddon
et al.

Q-G-N-K-V-V-L-G-K-K-G-D-T-V-E-L-T-C-T-A-S-E

In the above sequences, the amino acids
are represented by single letter codes as follows:

5 Phe: F Leu: L Ile: I Met: M
 Val: V Ser: S Pro: P Thr: T
 Ala: A Tyr: Y His: H Gln: Q
 Asn: N Lys: K Asp: D Glu: E
 Cys: C Trp: W Arg: R Gly: G

10 X: not determined or ambiguous.

We also constructed pBG211-11, a plasmid
coding for the N-terminal 113 amino acids of soluble
T4 protein. This construct, which codes for a pro-
tein characterized by a single disulfide bridge,
15 between the cysteines at amino acid positions 18 and
86, is conveniently expressed in E.coli.

To construct p211-11, as depicted in Fig-
ure 24, we first cut p195-8 (see Figures 8D and 9A)
with ClaI to remove the ClaI-ClaI cassette contain-
20 ing the cDNA sequence of rsT4.2. We then digested
pAT153y3SH16ΔAmp, the tryptophan operon promoter
plasmid from the gamma interferon producing E.coli
strain BN374 with ClaI, and deleted the cDNA coding
for gamma interferon. Subsequently, we inserted
25 the ClaI-ClaI cassette into the ClaI-cut E.coli
plasmid in front of the tryptophan operon promoter
and ligated to produce p196-10.

As shown in Figure 25, we then subjected
pBG380 to oligonucleotide-directed mutagenesis to
30 insert three tandem translational stop codons follow-
ing the T4 cDNA sequence coding for amino acids -23
to 113 in pBG380, to produce pBG394.

We then constructed p211-11 from fragments
of each of p196-10, pBG394 and p1034 as depicted in
35 Figure 26. The first fragment including the vector
sequences, was produced by restricting p196-10 with


3708

89085519

-59-

HindIII and ClaI to remove the T4 coding sequence from amino acids 61 through 374 of rsT4.2 and including vector sequence following the 3' end of the rsT4 gene. The second fragment, a HindIII - BglII segment including the codons for T4 amino acids 61-113 of rsT4.9 immediately followed by a triplet of stop codons in tandem, was isolated by HindIII/BglII digestion of pBG394. The third fragment, a BamHI - ClaI fragment containing a bacteriophage T4 transcriptional termination signal [H. N. Kirsch and B. Allet, "Nucleotide Sequences Involved In Bacteriophage T4 Gene 32 Translational Self-Regulation", Proc. Natl. Acad. Sci. USA, 79, pp. 4937-41 (1982)], was isolated by BamHI/ClaI digestion of p1034. We then ligated these three fragments to produce p211-11, a T4 construct coding for a 113 amino acid soluble form of T4 protein, with asparagine at amino acid position 3 (i.e., rsT4.113.1).

We then subjected p211-11 to oligonucleotide site-directed mutagenesis (Figure 27) to change the amino acid at position 3 from asparagine to lysine using the oligonucleotide T4-66:



This produced plasmid p214-10, a fully corrected 113 amino acid soluble T4 vector coding for a 113 amino acid soluble form of T4 protein, with lysine at amino acid position 3 (i.e., rsT4.113.2). As shown in Figure 27, we subjected p214-10 to oligonucleotide site-directed mutagenesis to delete glutamine and glycine at, respectively,

3709

89085519

-60-

amino acid positions 1 and 2 of the T4 sequence using the oligonucleotide T4AID-87:

C
↓

5' GTA TCG ATT TGG
ATG ATG AAA AAA
GTA GTA 3'.

This produced p215-7, a 111 amino acid soluble T4 construct, including the trp promoter, which directs the expression of a 111 amino acid soluble form of T4 protein, with lysine at amino acid position 3 (i.e., rsT4.111).

We next constructed p218-8, a 111 amino acid construct which directs the expression of a 111 amino acid soluble form of T4 protein, with lysine at amino acid position 3 (i.e., rsT4.111) under the control of the P_L promoter, as depicted in Figure 28.

More specifically, we cut p197-12 (Figure 9A) with ClaI to remove the 101 bp fragment containing linker and terminator sequences. We also cut p215-7 with ClaI to remove the ClaI - ClaI cassette containing the cDNA sequence of rsT4.111 and the ϕ T4 transcriptional terminator sequence [Kirsch and Allet, supra]. Subsequently, we inserted the ClaI - ClaI cassette into the ClaI-cut p197-12 to produce p218-8.

In order to express rsT4.113.1, we transformed E.coli A89 with p211-11 by conventional techniques [Maniatis et al. (1982), supra] to form E.coli A89/p211-11. E.coli A89 is a tetracycline sensitive derivative of E.coli SG936. We isolated E.coli A89 from E.coli SG936 according to the method of S. R. Maloy and W. D. Nunn, "Selection For Loss Of Tetracycline Resistance By Escherichia coli", J. Bact., 145, pp. 110-12 (1981), which is based upon the ability of the lipophilic chelating agent fusaric acid to selectively inhibit resistant strains.

3710

89085519

-61-

More specifically, we plated *E. coli* SG936 on medium containing, per liter, 5 g tryptone, 5 g yeast extract, 10 g NaCl, 10 g $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, 50 mg chlortetracycline-HCl, 12 mg fusaric acid, 0.1 mM ZnCl_2 and 15 g agar. Colonies which grew at 30°C (putative tetracycline-sensitive strains) were retested for tetracycline sensitivity on L-agar plates containing 5 µg/ml tetracycline. One tetracycline-sensitive strain, designated A89, was then shown to be unable to grow on LB agar at 42°C, thus verifying the presence of the *htpR* mutation.

Transformants were selected by tetracycline resistance. We picked a single colony into 20 ml of minimal medium plus 0.2% casamino acids plus tryptophan (100 µg/ml) plus tetracycline (10 µg/ml) in a 100 ml shake flask placed in a shaking air incubator at 30°C and allowed the cells to grow up overnight. The following morning, we inoculated 40 ml of minimal medium plus 0.2% casamino acids plus tryptophan (100 µg/ml) plus tetracycline (10 µg/ml) with the overnight culture at $\text{OD}_{600} = 0.05$ in a 500 ml flask. The cells were grown to midlog phase and then induced by pelleting, washing once in minimal medium and then resuspending in minimal medium plus 0.2% casamino acids plus tetracycline (10 µg/ml), in the absence of tryptophan. We removed 0.6 OD_{600} of cells after 0, 1, 2, 3 and 4 hours incubation and after growth overnight.

The aliquots were centrifuged and cell pellets were subjected to lysis by boiling in Laemmli gel loading buffer. After centrifugation to remove cell debris, half of each sample was subjected to SDS-PAGE, followed by Western blot analysis with our rabbit antipeptide antibody probes or by Coomassie blue protein staining (Figures 29A and 29B).

3711

89085519

-62-

Purification Of rsT4.113.1

We then purified rsT4.113.1 from the E.coli transformant by means of two essentially quantitative steps involving anion-exchange and gel-filtration chromatographies performed under reducing and denaturing conditions.

More specifically, we suspended 14 g of wet cells from a 4 L shake-flask fermentation in 100 ml of a 20mM Tris (pH 7.5) buffer containing 20 µg/ml DNase, 20 µg/ml RNase and 1 mM phenylmethylsulfonylfluoride ("PMSF"). The suspension was applied to a French Press at 1000 psi in two passages and then centrifuged in an SA 600 rotor at 18,000 g for 15 min at 4°C. The resulting pellet was solubilized in 20 ml of a 20 mM Tris (pH 7.5) buffer containing 7 M urea and 10 mM 2-mercaptoethanol. We then subjected the suspension to ultracentrifugation at 85,000 g for 90 min at 4°C. The supernatant was diluted by the addition of 80 ml of 20 mM Tris (pH 7.5) buffer containing 7 M urea and 10 mM 2-mercaptoethanol and 40 ml of the sample was applied to a 3 x 4 cm Q-Sepharose fast-flow column (Sigma, St. Louis, Missouri) which had been pre-equilibrated in the same buffer. The column was developed with a gradient in 400 ml total volume of increasing NaCl from 0 to 0.3 M in the same Tris/urea/2-mercaptoethanol buffer. Column fractions were monitored for absorbance at 280 nm and for protein content by SDS-PAGE (15% acrylamide). The fractions were also analyzed by Western blots. Figure 30, panel (a) is a chromatogram displaying the purification of rsT4.113.1 by ion-exchange chromatography. In that figure, peaks containing rsT4.113.1 are identified. The rsT4.113.1 was found to elute early in the NaCl gradient and to be well-resolved from low-molecular weight contaminants.

-63-

In order to separate rsT4.113.1 from high-molecular weight contaminants, we carried out gel-filtration chromatography on an rsT4.113.1-containing pool for final purification of the protein to near homogeneity (>95% purity). More specifically, we prepared a pool containing 20 mg of protein in 50 ml and then concentrated to 10 ml in a stirred-cell ultrafiltration unit (Amicon, Danvers, MA.) using a PM-30 membrane (Amicon). Subsequently, 5.0 ml of the concentrate was applied to a 1.5 x 95 cm S-300 column (Sigma) equilibrated and developed in the same Tris/urea/2-mercaptoethanol buffer. We monitored the column fractions for absorbance at 280 nm and for protein content by SDS-PAGE. The fractions were also analyzed by Western blots. A pool containing rsT4.113.1 (approximately 4 mg) in 15 ml was thus prepared. Figure 30, panel (b) is a chromatogram displaying the purification of rsT4.113.1 by gel-filtration separation of the rsT4.113.1 pool. In that figure, peaks containing rsT4.113.1 are identified.

Figure 30, panel (c) is an SDS-PAGE analysis depicting the purification of the rsT4 derivative throughout the centrifugation and chromatography steps. In Figure 30, panel (c), the lanes depicted are:

lane A:	molecular weight standards
lane B:	cell extracts
lane C:	cell pellet following solubilization of cell extract in non-denaturing conditions
lane D:	supernatant following solubilization of cell extract in non-denaturing buffer
lane E:	supernatant following ultracentrifugation step

3713

89085519

-64-

lane F: Q-Sepharose pool
lane G: S-300 gel-filtration pool.

Refolding Of Purified rsT4.113.1

We refolded the purified rsT4.113.1 by
5 dilution and dialysis steps to non-denaturing and
oxidized conditions. More specifically, refolding
of the protein at a concentration of 0.5 OD (280)/ml
was achieved by stepwise dialysis against 500 volumes
10 of 3 M urea, 20 mM Tris (pH 7.5); 500 volumes of 1 M
urea, 0.1 M ammonium acetate (pH 6.8) and, finally,
the same volume of a phosphate-buffered saline solu-
tion. Throughout the refolding procedure, samples
of the protein were monitored for relative content
by spectral analysis and by high-performance liquid
15 chromatography ("HPLC") performed on a 150A liquid
chromatographic system (Applied Biosystems, Inc.,
Foster City, California). An octasilyl column
(Aquapore RP-300, 0.46 x 3.0 cm) was equilibrated in
80% 0.1% trifluoroacetic acid ("TFA")/water (sol-
20 vent A) and 20% 0.085% TFA/70% acetonitrile (sol-
vent B) and developed with a linear gradient of
increasing acetonitrile concentration from 20% to
80% (solvent B) over 45 min at a flow rate of
0.5 ml/min.

25 As shown in Figure 31, panel (a), protein
in 7 M urea, 10 mM 2-mercaptoethanol and 20 mM
Tris(pH 7.5) eluted from the HPLC column at 49%
acetonitrile in the gradient. In subsequent steps,
from 1 M urea/1 mM ammonium acetate (pH 6.8) [Fig-
30 ure 31, panel (b)] to phosphate buffered saline
[Figure 31, panel (c)], an increasing percentage of
rsT4.113.1 was found to elute earlier in the HPLC
gradient -- at 47% acetonitrile. The identity of
the earlier eluting peak as oxidized product was
35 verified by reduction of rsT4.113.1 in non-chaotropic

-65-

solutions and application of sample thus treated to HPLC under the same conditions.

The elution of oxidized rsT4.113.1 prior to reduced protein on HPLC suggests that formation of the single disulfide bridge decreases relative hydrophobicity of the protein [J. L. Browning et al., Anal. Biochem., 155, pp. 123-28 (1986)]. Spectral analysis of rsT4.113.1 was performed throughout the course of refolding in order to monitor relative yield of soluble protein in the procedure. The refolding method allowed approximately 20% recovery of rsT4.113.1. HPLC analysis indicated a less than 15% contaminant of reduced protein in the preparation (Figure 30, panel (c), lane G).

15 Sequencing Of Renatured rsT4.113

We then carried out amino acid analysis of rsT4.113.1 by automated Edman degradation in an Applied Biosystems 470A gas phase sequenator equipped with a 900 A data system. Phenylthiohydantion amino acids generated during the course of the degradative chemistry were analyzed on-line using an Applied Biosystems 120A PTH-analyzer equipped with a PTH-C18 2.1 x 220 mm column. Protein (10 µg) for sequence analysis was applied to SDS-PAGE (15% acrylamide) and electroblotted on an Immobilon membrane (Millipore Corp., Bedford, Massachusetts) as described by P. Matsudaira, J. Biol. Chem., 262, pp. 10035-38 (1987).

Amino acid analysis of protein samples was performed by hydrolysis of protein in 6 N HCl, in vacuo, for 24 h at 110°C. The hydrolysates were then applied to a Beckman 6300 Analyzer equipped with post-column detection by ninhydrin. Western blot analysis of the SDS-PAGE gels was carried out by standard techniques using rabbit antisera JB-1.

3715

89085519

-66-

Sequence analysis revealed an amino terminal sequence of: Met-Gln-Gly-Asn-Lys-Val-Val ...

5 The purified rsT4.113.1 protein was found to contain stoichiometric quantities of the amino-terminal methionine placed in the protein construct for expression in E.coli and an intact polypeptide chain consistent with a sequence derived from the plasmid construction. Recovery of phenylthiohydantoinyl-methionine at the first cycle of the degradative chemistry was 60% consistent with routine initial yields obtained in the automated Edman. This observation excludes the possibility that a significant percentage of the rsT4.113.1 lacked the initiation methionine, i.e., the NH₂-methionine was not removed by expression of rsT4.113.1 in E.coli, or that sequence analysis was impaired by the presence of glutamine at the first cycle of the degradative chemistry. Sequence analysis was performed for 40 cycles and no evidence of lysine carbamylation was observed. Amino acid analysis displayed a close correlation of actual and theoretical values for amino acids, thus indicating the marked absence of proteolytic degradation in the course of expression, or purification, or both.

25 Immunoprecipitation Of CHO Cell Lines Producing Soluble T4

We tested the conditioned media from ³⁵S-Cys metabolically labelled CHO cells transfected with one of the T4 mutant constructs pBG377, pBG380, pBG381, the full length recombinant T4 construct pBG379, of this invention or vector only, to determine whether any produced a molecule recognized by the anti-T4 monoclonal antibody 19 Thy. To carry out this test, we incubated about 10⁷ CHO cells transfected with either pBG380, pBG381, pBG377, pBG379 or pBG312, for 35 5 hours at 37°C with 180 µCi/ml ³⁵S-labelled cysteine

89085519

-67-

[DuPont, New England Nuclear] in 4 ml RPMI cys⁻ medium (Gibc). After labelling of the cells, 1 ml of filtered, conditioned media was made 0.5 mM with phenyl-methyl-sulphonyl fluoride and immunoprecipitated

5 with OKT4 and protein A Sepharose [P. H. Sayre and E. L. Reinherz, Eur. J. Immunol., 15, pp. 291-95 (1985)]. Subsequently, we incubated media from the ³⁵S-labelled cells with OKT4 (ATCC #CRL 8002). We then immuno-precipitated with protein A Sepharose

10 and subjected the immuno-precipitates to SDS-PAGE under reducing conditions on 10% polyacrylamide gels [U. K. Laemmli, Nature, 227, pp. 680-85 (1980)]. Autoradiography was carried out with X-Omat X-ray film (Eastman Kodak).

15 As shown in lanes 3-5 of Figure 32, both pBG380 (rsT4.2) and pBG381 (rsT4.3) directed the synthesis of a secreted, immune, ³⁵S-labelled T4 protein that was recognized by the OKT4 anti-T4 antibody. The immunoprecipitated truncated molecules migrated as 49 Kd proteins, a result consistent with their predicted molecular weights. In contrast, no soluble T4 antigen could be detected in the conditioned media of cell lines stably transfected with pBG377 (rsT4.1) or pBG379 (rflT4).

20 Immunoprecipitation analysis of cellular extracts of cell lines transfected with pBG377 suggests that the rsT4.1 gene may be misfolded, which could account for a block in its secretion [M. J. Gething et al., Cell, 46, pp. 939-50 (1986)].

25 In Figure 32, the lanes represent the following: Lane 1: immunoprecipitation from conditioned medium of CHO cells stably co-transfected with vectors pBG312 and pAdD26. Lane 2: blank. Lanes 3 and 4: immunoprecipitation from conditioned

30 medium of CHO cells stably co-transfected with pBG380 (rsT4.2) and pAdD26. Lanes 5 and 6: immunoprecipitation from conditioned medium of CHO cells stably

35

3717

89085519

-68-

co-transfected with pBG381 (rsT4.3) and pAdD26.
Lane 7: immunoprecipitation from conditioned medium
of CHO cells stably co-transfected with recombinant
full length T4 (pBG379) and pAdD26. In Figure 32,
5 the arrow indicates the predicted position of the
soluble T4 from pBG380 or pBG381 relative to the
migration of standard molecular weight markers.

Immunoprecipitation Of COS 7 Cell Lines
Producing Recombinant Soluble T4

10 We expressed recombinant soluble T4
derivatives pBG392, pBG393 and pBG394 in COS 7 cells
by electroporation, essentially as described by
G. Chu et al., "Electroporation For The Efficient
Transfection Of Mammalian Cells With DNA", Nuc.
15 Acids Res., 15, pp. 1311-26 (1987). More specifi-
cally, we introduced 20 µg closed circular plasmid
DNA and 380 µg of carrier (sonicated salmon sperm
DNA) into 3×10^7 COS 7 cells. The cells were
electroporated using a Gene Pulser (Biorad) set at
20 300 volts. Subsequently, we incubated the COS 7
cells in Dulbecco's Modified Eagle's Medium supple-
mented with 10% fetal calf serum for 24 hours. We
then harvested the conditioned media, filtered it
through a Millipore Millidisk 0.22 µ hydrophilic
25 filter cartridge (Millipore #MCGL 305-01) and
concentrated the secreted proteins on a fast-S ion
exchange column (S-Sepharose Fast Flow, Pharmacia
#17-0511-01) in 20 mM MES buffer (pH 5.5).

We then eluted the bound proteins with
30 20 mM Tris-HCl (pH 7.7) and 0.3 M NaCl. The elution
pool was subsequently diluted with 2 volumes of 20 mM
Tris-HCl (pH 7.7) and it was then loaded on a column
comprising either 19Thy anti-T4 monoclonal antibody
and protein A Sepharose or OKT4A and protein A
35 Sepharose. We washed the column extensively and
eluted the bound material as 0.5 ml fractions with

3718

89085519

-69-

50 mM glycine-HCl (pH 2.5), 150 mM NaCl, 0.1 mM EGTA and 5 µg/ml Bovine pancreatic trypsin inhibitor, Aprotinin (Sigma, #A1153). The immunoprecipitates were subjected to SDS PAGE (10% gel) followed by immunoblotting against rabbit antisera raised against peptide JB-1. We employed silver stained gels to follow binding and elution of rsT4 during chromatography.

Figure 33 depicts an immunoblot analysis of transiently expressed pBG392 (rsT4.7) [lanes 10, 11]; pBG393 (rsT4.8) [lanes 4, 7, 8] and pBG394 (rsT4.9) [lane 5]. The standards are 50 ng purified rsT4.3 (lane 1); 150 ng purified rsT4.3 (lane 2) and 250 ng purified rsT4.3 (lane 3). The arrow indicates the expected position of migration of a protein with the relative molecular weight of rsT4.7: 21,000 daltons. The sample that was to be loaded into lane 4 was lost and lanes 6 and 9 are blank.

As shown in lanes 10 and 11 of Figure 35, pBG392 (rsT4.7) directed the synthesis of a secreted, immune protein that was recognized by the anti-T4 antibodies OKT4A and 19Thy. Lanes 4, 7 and 8 also demonstrate that pBG393 (rsT4.8) directed the synthesis of a secreted, immune protein that was recognized by OKT4A and 19Thy. This analysis illustrates that rsT4.7 contains the OKT4A epitope. It also suggests that the binding region for HIV envelope binding resides in the amino 182 terminal residues of T4.

In contrast, no soluble T4 could be detected in the media of cell lines transfected with pBG394 (rsT4.9) [see lane 5]. Immunoprecipitation analysis of cellular extracts of cell lines transfected with pBG397, however, showed that rsT4.9 was recognized by OKT4A. We believe that rsT4.9, a 113 amino acid construct, binds the HIV virus and that it represents a second generation soluble T4, one with only two

3719

89085510

-70-

cysteines and one of three disulfide bridges. Accordingly, rsT4.9 is easily produced in E.coli or yeast systems.

5 Similarly, although no soluble T4 could be detected in the media of cell lines transfected with pBG396 (rsT4.12), analysis of cellular extracts of those cell lines showed that rsT4.12 was recognized by OKT4A. Thus, rsT4.12 may also bind HIV virus.

10 Radioimmunoassay And Epitope Analysis Of rsT4.113

In order to determine if the 113 fragment of rsT4 contained structural determinants for binding to OKT4A, Leu-3A and OKT4, we then carried out radioimmunoassay and epitope analysis of rsT4.113
15 using a competitive inhibition radioimmunoassay [C. J. Newby et al., "Solid-Phase Radioimmune Assays" in Handbook Of Experimental Immunology, D. M. Weir (Ed.), 1, pp. 34.1-34.8 (1986)]. As OKT4A and Leu-3A block infectivity of HIV in vitro [Dalglish et al.,
20 supra] and binding of T4 to gp120/160 [McDougal et al., supra], this analysis served as a first approximation as to whether or not rsT4.113 contained structural elements for interaction with HIV.

We first coated U-bottom 96 well microtiter
25 plates (Falcon) with 50 µl/well goat-anti-mouse IgG (Hyclone Typing Kit, Logan, Utah) in PBS (pH 7.0) to a concentration of 50 µg/ml and incubated the plates overnight at 4°C. We then rinsed the plates with 1X PBS and blotted them dry. The plates were then
30 blocked by the addition of 100 µl/well of a 1X PBS solution containing 5% bovine serum albumin for 1 hour at room temperature. We rinsed the plates with PBS, blotted dry and then spotted them with 50 µl of one of three antibody solutions containing
35 either OKT4 (10 µg/ml in block buffer); OKT4A (500 ng/ml in block buffer) or Leu-3A (Becton-

-71-

Dickinson) (500 ng/ml in block buffer). We let the plates stand for 2 hours at room temperature. We then washed the plates 3 times with a PBS/0.05% Tween-80 solution and 2 times with 1X PBS and blotted them dry.

5 In a separate plate, we titrated competitor samples of unlabeled rsT4.113.1 from 20 µg/ml and serially diluted twice (including no competitor control), with final volumes in each well of 25 µl.

10 The positive control for this assay was competition with unlabeled rsT4.3 (375 amino acids). We then added 25 µl of ¹²⁵I-rsT4.3 containing 10,000 cpm/25 µl (prepared according to A. E. Bolton and W. M. Hunter, Radioimmunoassay And Related Methods,

15 Chapter 2c). Subsequently, we spotted the entire 50 µl content of each well onto the assay plate containing each of the three antibody solutions and incubated for 2 h at room temperature. We then washed the plates 3 times with a PBS/0.5% Tween-80

20 solution and 2 times with 1X PBS, blotted them dry and then counted the wells in a Beckman gamma counter for radioactivity.

As shown in Figure 34, rsT4.113.1 competes with ¹²⁵I-rsT4.3 for absorption to an OKT4A solid

25 phase in a dose-dependent manner. Additionally, rsT4.113.1 competes with ¹²⁵I-rsT4.3 for absorption to a Leu-3A solid phase in a dose-dependent manner. By comparison to unlabeled rsT4.3, rsT4.113.1 exhibits a molar affinity for those antibodies within a factor

30 of 3. In the 0.4 to 25 µg/ml concentration range tested, rsT4.113 did not compete with radiolabelled rsT4.3 for binding to OKT4. In a similar assay, we observed that rsT4.111 also competes with ¹²⁵I-rsT4.3 for binding to OKT4A and Leu-3A, but not to OKT4

35 [Figures 35-37].

Based on these results, we believe that the epitopes for OKT4A and Leu-3A are contained within

3721

89085519

-72-

the amino-terminal 113 amino acids of T4. We also believe that the epitope for OKT4 binding is localized within the carboxy terminal of the T4 polypeptide.

Accordingly, we believe that the gp120-binding domain is localized within the amino terminal 113 or 111 amino acids of the T4 protein. Based on this belief, we synthesized various synthetic oligopeptides which contain sequence within that structural domain. These oligopeptides are represented in Figure 3 as follows:

	<u>Oligopeptide</u>	<u>Amino Acid Coordinates</u>
	JB-1	44-63
	rsT4 #6	18-29
	rsT4 #7	5-56
15	rsT4 #8	84-97
	rsT4 #9	30-63

We synthesized these peptides using conventional phosphoamide DNA synthesis techniques [Tetrahedron Letters, 22, pp. 1859-62 (1981)]. We synthesized the peptides on an Applied Biosystems 380A DNA Synthesizer and purified them by gel electrophoresis.

ELISA Assay For rsT4.113

We also carried out an ELISA assay for rsT4.113.1 produced by p211-11-transformed E.coli. Throughout this assay, dilutions were made in blocking solution and, between each step, we washed the plates with PBS/0.05% Tween-20. More specifically, we coated wells of Immulon 2 (Dynatech, Chantilly, Virginia) plates with .005 OD (280 nm)/ml of OKT4 (IgG2b) in 0.05 M bicarbonate buffer to a volume of 50 µl/well and incubated the plates overnight at 4°C. We then blocked the plates with 5% bovine serum albumin in PBS, 200 µl/well, and incubated for 30 minutes at room temperature.

Subsequently, we added 50 µl of 50 ng/ml rsT4.3 to each well, incubating overnight at 4°C.

3722

89085519

-73-

We then added 50 μ l/well of a mixture containing rsT4.113.1 and 10 ng/ml of OKT4A and incubated for 2 1/2 hours at room temperature. Using a Hyclone Kit (Hyclone), we then carried out the following steps. First, we added 1 drop of rabbit anti-mouse IgG2a to each well and incubated the plates for 1 hour at room temperature. We then added 100 μ l of peroxidase-labeled anti-rabbit IgG, diluted 1:4000 with blocking buffer to each well, and incubated for 1 hour at room temperature.

We prepared a substrate reagent as follows. We diluted substrate reagent 1:10 in distilled water and added two O-phenyl-ethylene-diamine ("OPD") chromophore tablets per 10 ml of substrate. We let the mixture dissolve thoroughly by mixing with a vortex. Alternatively, a TMB peroxidase substrate system (Kirkegaard & Perry Catalogue #50-76-00) may be used. Subsequently, we added 100 μ l of the chromophore solution to each well, incubated for 10-15 minutes at room temperature and then stopped the color development with 100 μ l of 1N H_2SO_4 . We then measured OD at 490 nm, using an ELISA plate reader.

The results of the assay are demonstrated in Figure 38.

We then subjected the soluble T4 proteins produced by the T4 constructs of this invention to various functional assays.

Assays Of The Antiviral Activity Of Soluble T4

The antiviral activity of soluble T4 according to this invention was evaluated using modifications of various in vitro systems used to study antiviral agents and neutralizing antibodies [D. D. Ho et al., "Recombinant Human Interferon Alpha (A) Suppresses HTLV-III Replication In Vitro", Lancet, pp. 602-04 (1985); K. Hartshorn et al.,

89085519

-74-

"Synergistic Inhibition Of HTLV-III Replication
In Vitro By Phosphonofornate And Recombinant Inter-
feron Alpha-A", Antimicrob Ag Chemoth, 30, pp. 189-91
(1986)].

5 For each of these assays, we prepared graded
concentrations of soluble T4 and preincubated them
with an H9 derived IIIB isolate of HIV [a gift from
Drs. M. Popovic and R. Gallo, National Cancer
Institute, Bethesda, Maryland]. The isolate was
10 maintained as a chronically infected culture in H9
cells. Cell-free HIV stocks were obtained from
supernatant fluids of HTLV-III infected H9 cultures
(culture conditions: 1×10^6 cells/ml with 75% viable
cells). We prepared serial 10 fold dilutions of
15 recombinant soluble T4 ranging from 10 picograms/ml
to 10 micrograms/ml and incubated them with fifty
50% tissue culture infectious doses (TCID₅₀) of HIV
for 1 hour at 37°C, in RPMI-1640 supplemented with
20% heat inactivated fetal calf serum (FCS). We
20 then added 150 µl of H9 cells to a final concentra-
tion of 0.5×10^6 cells/ml which were not HIV-infected
to the wells containing aliquots of the recombinant
soluble T4/HIV mixture.

We adjusted each virus inoculum to a con-
25 centration of 250 TCID₅₀/ml. We preincubated 100 µl
of the virus inoculum with 200 µl recombinant solu-
ble T4 or 100 µl immunoglobulin prepared in tripli-
cate serial 2-fold dilutions for 1 hour at 37°C
prior to inoculation onto $1.5 - 2 \times 10^6$ H9 cells in
30 5 ml RPMI 1640 supplemented fetal calf serum (20%),
HEPES (10mM), penicillin (250 U/ml), streptomycin
(250 µg/ml) and L-glutamine (2mM). On days 5, 6, 7,
10 and 14, we examined each culture for characteris-
tic cytopathic effects ("CPE"). Neutralization was
35 defined as the inhibition of syncytia formation com-
pared with controls.

3724

89085519

-75-

The positive control used was HIV ser positive neutralizing serum, as described in D. D. Ho et al., "Human Immunodeficiency Virus Neutralizing Antibodies Recognize Several Conserved Domains On The Envelope Glycoproteins", J. Virol., 61, pp. 2024-28 (1987). The negative controls used were HIV seronegative serum only and buffer only.

Cytopathic Effect Assay (CPE)

In this assay, following conventional protocols for cytopathic effect assays [Klatzmann et al. (1984), supra and Wong-Staal and Gallo (1985), supra], we microscopically examined the H9 cells for evidence of cytopathic effects of HIV.

The CPE was scored on a four point scale from 1+ to 4+, with 4+ representing the highest degree of CPE.

By day 14, wells containing recombinant soluble T4 according to this invention (rsT4.2, derived from the pBG380 transfected CHO cell line BG380) at 10 µg/ml showed no evidence of CPE, while the negative control showed 1+ to 3+ CPE.

p24 Radioimmunoassay

We then tested soluble T4 as an inhibitor of viral replication in an HIV virus replication assay according to D. D. Ho et al., J. Virol., 61, pp. 2024-28 (1987) and J. Sodroski et al., Nature, 322, pp. 470-74 (1986). We carried out the assay essentially as described, except that the cultures were propagated in microtiter wells containing 200 µl. In this assay, we evaluated the ability of the soluble T4 polypeptides of this invention to block HIV replication, as measured by HIV p24 antigen production. We sampled supernatants twice weekly for HIV p24 antigen as described below.

3725

89085519

-76-

We obtained an assay kit [HTLV-III p24 Radioimmuno assay System, Catalogue No. NEK-040, NEK-040A, Biotechnology Systems, New Research Products, Dupont] which contains affinity purified ^{125}I labelled HIV p24 antigen, a rabbit anti-p24 antibody and a second goat anti-rabbit antibody which is used to precipitate antigen-antibody complexes. We carried out the assay according to the protocol included with the kit. Accordingly, we mixed a sample to be assayed or one of a series of amounts of unlabelled p24 antigen with a fixed amount of ^{125}I labelled p24 and a fixed limited amount of rabbit anti-p24 antibody. We incubated the samples overnight at room temperature and then added a goat anti-rabbit immunoglobulin preparation for 5 minutes at 40°C . We centrifuged the samples in a microfuge and aspirated the supernatant fluid. Pelleted ^{125}I labelled p24 was quantitated for each sample by gamma counting and a standard curve for the ^{125}I p24 displaced by the known amounts of antigen added to standard tubes was constructed. We then calculated the ^{125}I labelled p24 displaced by the antigen present in the unknown samples by interpolation using the standard curve constructed from the known amounts of p24 antigen contained in the standard samples. The results are shown in the table below.

3726

89085519

-77-

p24 ASSAY OF F⁻V REPLICATION INHIBITION

	<u>Day</u>	<u>rsT4.2</u> <u>(μg/ml)</u>	<u>Patient</u> <u>Serum</u>	<u>Average</u> <u>CPM</u>	<u>% Bound/</u> <u>Unbound</u>
5	7	-	Negative	344	8.5
		-	Positive	2,237	112.4
		0.5*	-	551	19.9
		5.0**	-	1,766	86.6
10	10	-	Negative	230	2.2
		-	Positive	2,459	124.6
		0.5*	-	322	7.3
		5.0**	-	1,980	96.3
15	14	-	Negative	221	1.8
		-	Positive	2,284	115.0
		0.5*	-	246	3.1
		5.0**	-	1,988	98.7

These results demonstrate that soluble T4 according to this invention at a concentration of 5 μ g/ml completely inhibits virus replication as measured in this standard 14 day assay. These results are also depicted in Figure 39 in graphic form. In Figure 39, values were calculated from a standard curve of p24 according to assay kit instructions.

* This concentration was initially believed to be 1.0 μ g/ml, based upon our preliminary approximation that 1 unit of absorbance at 280 nm (" A_{280} "), was equivalent to 1 mg of rsT4.2. Absorbance at 280 nm is a commonly used first approximation of protein concentration. Upon amino acid analysis of the protein, however, we found that it had a higher extinction coefficient than originally approximated, with 1 A_{280} unit of rsT4.2 being equivalent to 0.5 mg of the protein.

** This concentration was initially believed to be 10 μ g/ml, based upon our preliminary approximation that 1 unit of absorbance at 280 nm (" A_{280} "), was equivalent to 1 mg of rsT4.2. Absorbance at 280 nm is a commonly used first approximation of protein concentration. Upon amino acid analysis of the protein, however, we found that it had a higher extinction coefficient than originally approximated, with 1 A_{280} unit of rsT4.2 being equivalent to 0.5 mg of the protein.

We then carried out a p24 replication assay as described above, except that the soluble T4 was added to the infected cultures during refeeding at days 3, 7 and 10, in order to maintain a constant
 5 rsT4 concentration throughout the infection period. The results of this assay are shown in the table below.

**INHIBITION OF HIV REPLICATION
 WITH CONSTANT CONCENTRATION OF rsT4**

10	rsT4.2 (μ g/ml)	p24 (ng/ml)
	0.008	770
	0.031	970
	0.125	85
15	0.5	0
	5.0	0
	0	1120
	uninfected	0

These results demonstrate that when soluble
 20 ble T4 protein according to this invention was maintained at a constant concentration throughout the infection period, as little as 0.125 μ g/ml of the protein substantially blocked replication of 250 TCID₅₀/ml of HIV-1.

25 Advantageously, soluble T4 protein according to this invention, at concentrations far exceeding those required to block viral replication, did not exert immunotoxic effects in vitro, as measured by three lymphocyte proliferation assays -- mixed
 30 lymphocyte response, phytohemagglutinin, and tetanus toxoid-stimulated response.

Syncytia Inhibition Assay

To further assess the effect of soluble T4 on HIV env-T4 binding, we evaluated the effect of two
 35 preparations of our soluble T4 protein on the syncytiagenic properties of HIV in the co-cultivation assay. We carried out a C8166 cell fusion assay

-79-

as described in B. D. Walker et al., Proc. Natl. Acad. Sci. USA, 84, pp. 8120-24 (1987).

We incubated 1×10^9 H9 cells chronically infected with HTLV-IIIB for 1 hour at 37°C in 5% CO₂ with various concentrations of one of two preparations of rST4.2 in 150 µl RPMI-1640 media supplemented with 20% fetal calf serum. We then added 3×10^4 C8166 cells in 50 µl media (a T4⁺ transformed human umbilical cord blood lymphocyte line [Sodroski et al., supra], to a final volume of 0.2 ml in each well. Final well concentrations of soluble T4 were 0.5 µg/ml* and 5.0 µg/ml* for preparation #1 and 1.25 µg/ml* and 12.5 µg/ml* for preparation #2. We then counted total number of syncytia per well at 2 hours and 4 hours after adding the C8166 cells at 37°C in 5% CO₂. Parallel co-cultivations used buffer alone (negative control) or OKT4A at 25 µg/ml (positive control) as controls. We considered a positive result as a 50% reduction in syncytia compared to controls, at a time when at least 100 syncytia per 10^4 infected H9 cells were present in the control cultivations. The results of this assay are shown below and in Figure 40 (2 hour data).

25

* These concentrations were initially believed to be, respectively, 1 µg/ml, 10 µg/ml, 2.5 µg/ml and 25 µg/ml, based upon our preliminary approximation that 1 unit of absorbance at 280 nm ("A₂₈₀"), was equivalent to 1 mg of rST4.2. Upon amino acid analysis of the protein, however, we found that it had a higher extinction coefficient than originally approximated, with 1 A₂₈₀ unit of rST4.2 being equivalent to 0.5 mg of the protein.

30

3729

89085519

-80-

INHIBITION IN C8166 FUSION ASSAY

	<u>Preparation</u>	<u>[rsT4.2] ($\mu\text{g/ml}$)</u>	<u>% Inhibition*</u>	
			<u>2 Hrs</u>	<u>4 Hrs</u>
	buffer	0	0	0
5	rsT4.2	0.5**	30	42
	rsT4.2	5.0**	54	47
	rsT4.2	1.25**	16	21
	rsT4.2	12.5**	77	55
	OKT4A (25 $\mu\text{g/ml}$)	0	100	100

10 As demonstrated in this table and in Figure 40, soluble T4 according to this invention at 5.0 $\mu\text{g/ml}$ and 12.5 $\mu\text{g/ml}$ inhibited syncytia formation at 2 hours, as compared to buffer alone. By 4 hours after the addition of C8166 cells, soluble T4 at 12.5 $\mu\text{g/ml}$ continued to inhibit greater than 50% syncytia formation, as compared to the negative control.

15 We also evaluated the effect of two preparations of our soluble T4 protein rsT4.7 on the syncytiagenic properties of HIV in a similar co-cultivation assay. The results of this assay are shown below.

25 * All assays were carried out in triplicate, and the number of syncytia counted per well was averaged to calculate % inhibition. The % inhibition represents the difference between the average number of syncytia in the negative control (without rsT4 or OKT4A) and the average number of syncytia counted when either rsT4 or OKT4A were present during the assay, divided by the average syncytia count for the negative control and multiplied by 100.

30 ** These concentrations were initially believed to be, respectively, 1 $\mu\text{g/ml}$, 10 $\mu\text{g/ml}$, 2.5 $\mu\text{g/ml}$ and 25 $\mu\text{g/ml}$, based upon our preliminary approximation that 1 unit of absorbance at 280 nm (" A_{280} "), was equivalent to 1 mg of rsT4.2. Upon amino acid analysis of the protein however, we found that it had a higher extinction coefficient than originally approximated, with 1 A_{280} unit of rsT4.2 being equivalent to 0.5 mg of the protein.

-81-

INHIBITION IN C8166 FUSION ASSAYAssay date: day 1

	<u>Preparation</u>	<u>rsT4.7 (μg/ml)</u>	<u>Average Syncytia/50μl aliquot</u>	<u>% Inhibition at 2 Hrs</u>
5	H9 cells (control)	0	0	N/A
	C8166 cells (control)	0	0	N/A
10	HIV-infected H9 cells added to C8166 cells (control)	0	118	0
15	OKT4A (control)	0	0	100
	Prep. 1 of rsT4.7	$\approx 5.0^*$	43	63.6

- 20 * This concentration was initially believed to be 10 μ g/ml, based upon our preliminary approximation that 1 unit of absorbance at 280 nm (" A_{280} "), was equivalent to 1 mg of rsT4.2. Upon amino acid analysis of the protein, however, we found that it
- 25 had a higher extinction coefficient than originally approximated, with 1 A_{280} unit of rsT4.2 being equivalent to 0.5 mg of the protein.

3731

89085519

-82-

Assay date: day 13

	<u>Preparation</u>	<u>rsT4.7 (μg/ml)</u>	<u>Average Syncytia/50μl aliquot</u>	<u>% Inhibition at 2 Hrs</u>
5	H9 cells (control)	0	0	N/A
	C8166 cells (control)	0	1	N/A
10	HIV-infected H9 cells added to C8166 cells (control)	0	141	0
	OKT4A (control)	0	0	100
15	Prep. 2 of rsT4.7	$\approx 5.0^*$	27	80.9

* This concentration was initially believed to be 10 μ g/ml, based upon our preliminary approximation that 1 unit of absorbance at 280 nm (" A_{280} "), was equivalent to 1 mg of rsT4.2. Upon amino acid analysis of the protein, however, we found that it had a higher extinction coefficient than originally approximated, with 1 A_{280} unit of rsT4.2 being equivalent to 0.5 mg of the protein.

3732

89085519

-83-

Assay date: day 14

	<u>Preparation</u>	<u>rsT4.7 (μg/ml)</u>	<u>Average Syncytia/50μl aliquot</u>	<u>% Inhibition at 2 Hrs</u>
5	H9 cells (control)	0	0	N/A
	C8166 cells (control)	0	0	N/A
10	HIV-infected H9 cells added C8166 cells (control)	0	128	0
	OKT4A (control)	0	0	100
15	Prep. 1 of rsT4.7	$\approx 5.0^*$	35	72.7
	Prep. 2 of rsT4.7	$\approx 5.0^*$	2	98.4

As demonstrated in these tables, soluble T4 protein rsT4.7 inhibited syncytia formation in HIV-infected H9 cells.

We also evaluated the effect of rsT4.113.1 and rsT4.111 on the syncytiogenic properties of HIV in a co-cultivation assay. We carried out a C8166 cell fusion assay as described in Walker et al., supra.

We incubated 1×10^4 H9 cells chronically infected with HTLV-III_B for 1 hour at 37°C in 5% CO₂, with from 5 to 50 μ g/ml rsT4.113.1 or rsT4.111 in 150 μ l RPMI-1640 media supplemented with 20% fetal calf serum in 96-well microtiter plates. We

30

*- This concentration was initially believed to be 10 μ g/ml, based upon our preliminary approximation that 1 unit of absorbance at 280 nm ("A₂₈₀"), was equivalent to 1 mg of rsT4.2. Upon amino acid analysis of the protein, however, we found that it had a higher extinction coefficient than originally approximated, with 1 A₂₈₀ unit of rsT4.2 being equivalent to 0.5 mg of the protein.

35

3733

89085519

-84-

then added 3×10^4 C8166 cells to the wells in 50 μ l aliquots. The plates were incubated for 2 hours at 37°C in 5% CO₂ and, following this incubation, the number of syncytia per well were counted.

5 Syncytia were defined as cells containing a ballooning cytoplasm greater than three cell diameters. All samples were counted twice. Parallel co-cultivation used OKT4A alone or rsT4.3 alone at a concentration of 25 μ g/ml (positive controls) or H9
10 cells alone or C8166 cells alone (negative controls). The results of this assay are shown below and in Figure 41.

INHIBITION IN C8166 FUSION ASSAY

	<u>Preparation</u>	<u>rsT4(μg/ml)</u>	<u>% Inhibition</u>
15	H9 cells (control)	0	0
	C8166 cells (control)	0	0
	rsT4.113.1	1.25	35
	rsT4.113.1	2.5	63
	rsT4.113.1	4.25	63
20	rsT4.113.1	6.25	82
	rsT4.113.1	12.5	96
	rsT4.3	12.5	100
	OKT4A (25 μ g/ml)	0	100

As demonstrated in this table and in
25 Figure 41, rsT4.113.1 exhibited a dose-dependent inhibition of HIV-induced syncytia formation. The molar specific inhibitory activity of rsT4.113.1 appeared to be reduced by an order of magnitude by comparison to anti-viral activity of longer forms of
30 recombinant soluble T4. Thus, whereas rsT4.113.1 is effective toward neutralization of HIV-dependent cell fusion in vitro, its molar specific inhibitory

-85-

activity is decreased by a factor of 10. It is undetermined whether this decreased potency is due to incomplete renaturation of the E.coli-derived protein, the presence of three additional amino acids at the N-terminus of rsT4.113.1 (Met-Gln-Gly) lacking in rsT4.2 or rsT4.3 produced in mammalian cells, or the absence of additional structure in rsT4.113.1 required for high-affinity binding to HIV.

We also carried out a C8166 cell fusion assay with rsT4.111, as described for rsT4.113.1. The results of this assay are shown below.

INHIBITION IN C8166 FUSION ASSAY

	<u>Preparation</u>	<u>rsT4(µg/ml)</u>	<u>% Inhibition</u>
15	H9 cell (control)	0	0
	C8166 cells (control)	0	0
-	rsT4.111	1.25	0
	rsT4.111	2.5	40
	rsT4.111	4.25	20
20	rsT4.111	6.25	67
	rsT4.111	12.5	100
	rsT4.111	25.0	100
	rsT4.3	12.5	100
	rsT4.3	25.0	100
25	OKT4A (25 µg/ml)	0	100

As demonstrated in this table, rsT4.111 exhibited a dose-dependent inhibition of HIV-induced syncytia formation. At a concentration of 12.5 µg/ml and 25.0 µg/ml, complete inhibition of cell fusion was achieved.

Kinetics Of Intramuscular Injection Of Soluble T4

We examined the kinetics of the appearance of a recombinant soluble T4 protein according to this invention (specifically, rsT4.3 from the pBG381-transfected cell line BG381) in serum after intramuscular injection as follows.

3735

89085519

-86-

We obtained two cynomolgus monkeys (Macaca fascicularis) who were free of infectious disease and in good health. Each monkey had been subjected to a 6 week quarantine period prior to administration of the soluble T4 protein. Throughout the administration period, each monkey was maintained on a conventional diet of monkey chow supplemented with fresh fruit. A catheter and a vascular access port were surgically placed in a femoral vein of each animal prior to treatment in order to facilitate blood collection.

Over a period of 28 days, each animal received recombinant soluble T4 protein twice daily by intramuscular injection to the large muscles of the thighs or buttocks. Injections were administered to each animal 8 hours apart and each injection contained a volume of 0.15 ml/kg (0.25 mg/kg) of rsT4.3 (from the pBG381-transformed cell line BG381), for a total dose of 0.5 mg/kg/day/monkey. Serum samples for clearance determination were collected on day 1 before the first treatment and at 1, 2, 4 and 8 hours after the first injection, as well as 1, 2, 4, 14 and 16 hours after the second injection on days 7, 14 and 28.

We found that intramuscularly injected soluble T4 reached the maximum level in serum between 1 and 2 hours after injection, with the level falling off slowly and reaching half-maximum value at approximately 6 hours post-injection. According to data obtained for intravenous administration (not shown), the level of rsT4.3 in serum should drop below that attained via intramuscular injection approximately 2 hours after intravenous injection. Thus, while the maximum rsT4.3 level in serum after intramuscular injection does not reach that attainable via intravenous injection, it is slowly released into the blood stream, remaining detectable in serum for a

- 3736

Q0000510

-87-

much longer time. This slow release mechanism associated with intramuscular routes of injection is advantageous because a higher level of soluble T4 protein is available over a longer period of time over a given concentration; thus remaining in a sustained level. Intramuscular administration of soluble T4 protein is particularly useful in treating early stage HIV-infected patients, to prevent the virus from disseminating, or in treating patients who have been exposed to the virus and who are not yet seropositive.

We determined serum levels of rsT4.3 using an ELISA assay. Throughout this assay, dilutions were made in blocking solution and, between each step, we washed the plates with PBS/0.05% Tween-20. More specifically, we coated wells of Immulon 2 plates with .01 OD (280 nm)/ml of OKT4 (IgG2b) in 0.05 M bicarbonate buffer to a volume of 50 µl/well and incubated the plates overnight at 4°C. We then blocked the plates with 5% bovine serum albumin in PBS, 200 µl/well, and incubated for 30 minutes at room temperature.

Subsequently, we added 50 µl of sample or standard to each well, incubating for 4 hours at room temperature. We then added 50 µl/well of OKT4A at 0.1 µg/ml and incubated overnight at 4°C. Using a Byclone Kit (Byclone) we then carried out the following steps. First, we added 1 drop of rabbit anti-mouse IgG2a to each well and incubated the plates for 1 hour at room temperature. We then added 100 µl of peroxidase-labeled anti-rabbit IgG, diluted 1:4000 with 5% BSA/PBS to each well, and incubated for 1 hour at room temperature.

We prepared a substrate reagent as follows. We diluted substrate reagent 1:10 in distilled water and added two O-phenyl-ethylene-diamine ("OPD") chromophore tablets per 10 ml of substrate. We let

3737

89085519

-88-

the mixture dissolve thoroughly by mixing with a vortex. Alternatively, a TMB peroxidase substrate system (Kirkegaard & Perry Catalogue #50-76-00) may be used. Subsequently, we added 100 μ l of the
5 chromophore solution to each well, incubated for 10-15 minutes at room temperature and then stopped the color development with 100 μ l of 1N H_2SO_4 . We then measured OD at 490 nm, using an ELISA plate reader.

10 The results of the assay are demonstrated in the tables below.

89085519

-89-

Monkey #7-91

		<u>rsT4 Level</u> <u>(ng/ml)</u>			
	<u>Time(hr)</u>	<u>Day 1</u>	<u>Day 7</u>	<u>Day 14</u>	<u>Day 28</u>
5	0	22.7*	96.5	158.0	19.8
	1	278.8	199.6	360.7	238.3
	2	281.8	366.8	306.4	441.1
	4	214.9	246.6	363.9	393.2
	5				290.4
10	8	72.3	105.0	199.4	
	9**	246.2			
	10	259.6			
	12	136.0			
	22	23.8			
15	24	13.4			

Monkey #7-92

		<u>rsT4 Level</u> <u>(ng/ml)</u>			
	<u>Time(hr)</u>	<u>Day 1</u>	<u>Day 7</u>	<u>Day 14</u>	<u>Day 28</u>
20	0	6.7*	56.0	106.3	60.9
	1	87.2	225.8	178.0	437.7
	2	254.2	377.9	253.2	770.6
	4	170.0	167.3	308.2	821.5
	5				898.3
25	8	118.9	101.2	176.5	
	9**	405.1			
	10	523.5			
	12	371.5			
	22	48.4			
30	24	39.4			

* - background

** - second injection administered after the collection of the 8 hour sample.

Polyvalent Forms Of Recombinant Soluble T4

35 Receptors may be characterized by their affinity for specific ligands, such that, at equilibrium, the intrinsic affinity (K_a) between monovalent receptor and monovalent ligand can be defined as

40 $[RL]/[R_f][L_f]$, where $[RL]$ is the concentration of receptor (R) bound to ligand (L) and $[R_f]$ and $[L_f]$ are the concentrations of free receptor and ligand,

-90-

respectively [P. A. Underwood, in Advances In Virus Research, ed. K. Maramorosch et al., 34, pp. 283-309 (1988)].

For a polyvalent receptor (with a valency of n) binding to a polyvalent ligand (with a valency of m), a functional affinity can be defined as $n[R_b]/n[R_f]m[L_f]$, where $[R_b]$ is the concentration of bound receptor sites, and $n[R_f]$ and $m[L_f]$ are, respectively, the concentrations of free receptor and ligand binding sites. The effect of increasing the valence (the number of binding sites) is to enhance the stability of ligand-receptor complexes. The affinity of a polyvalent receptor for a polyvalent ligand will depend on three factors: the intrinsic association constant of each binding site, the valency (number of binding sites) and the topological relationship between the receptor and ligand binding sites. Under some circumstances, polyvalent binding interactions will lead to higher functional affinity. The decreased dissociation rate of polyvalent ligands with polyvalent receptors results in an increased functional affinity [C. L. Hornick and F. Karush, Immunochemistry, 9, pp. 325-40 (1972); I. Otterness and F. Karush, "Principles Of Antibody Reactions", in Antibody As A Tool, ed. J. J. Marchalonis and G.W. Warr, pp. 97-137 (1982)].

The simplest case for receptor polyvalency increasing functional affinity is represented by a bivalent soluble receptor, such as an antibody molecule, which has two identical ligand binding sites, each capable of independently binding antigen with equal affinity. If the antigen is displayed polyvalently, for example, chemically coupled to a solid support such that the spacing between antigenic sites can be bridged by the antibody's two antigen binding arms, the functional affinity of the antibody for the antigen coupled to the solid support would be

-91-

greater than the intrinsic affinity of the antibody binding site for the monovalent antigen [D. Crothers and H. Metzger, Immunochemistry, 9, pp. 341-57 (1972)]. Because virus particles represent poly-

5 valent antigens, the greater functional affinity of antibodies for polyvalent antigens is an important factor for antibody-directed virus neutralization.

The association of recombinant soluble T4 and the HIV major envelope glycoprotein gp120 is an

10 example of monovalent receptor binding to monovalent ligand. The affinity of this interaction has been measured, and the association between T4 and gp120 has a dissociation constant $K_d = 4 \times 10^{-9}$ M [L. Lasky et al., Cell, 50, pp. 975-88 (1987)].

15 Using the antibody analogy, we believe that polyvalent rsT4 will demonstrate a greater affinity for HIV-infected cells displaying gp120 than monovalent rsT4 and the topological relationship between gp120 on the virus particle or the

20 infected cell surface, will determine the degree to which polyvalent rsT4 exhibits higher functional affinity than monovalent rsT4. One example of a polyvalent rsT4 is described below, with respect to the production of a recombinant bivalent rsT4 con-

25 sisting of two tandem repeats of amino acids 3-178, followed by the C-terminal 199 amino acids of rsT4.3. According to this invention, a "polyvalent" receptor possesses two or more binding sites for a given ligand. Furthermore, the intrinsic affinity of each

30 ligand binding site of a given polyvalent receptor need not be identical.

As shown in Figure 42, to construct bivalent rsT4, we digested pBG391 with NheI, which cleaves after the valine at position 178 in rsT4, and removed

35 the NheI 5' overhang with mung bean nuclease. Next, we cleaved with BglII to remove the C-terminal half of the rsT4 coding sequence in pBG391. Finally, we

-92-

ligated a DraI-BglII fragment containing the coding sequence for rsT4 amino acids 3 (lysine) through 377 (isoleucine) to the cleaved pBG391 to create pBiv.1, a plasmid coding for a fusion protein with a tandem
5 duplication of the N-terminal 176 amino acids of rsT4, followed by the C-terminal 199 amino acids of rsT4.3. The protein produced by this plasmid, therefore, contains two adjacent N-terminal gp120-binding or OKT4A-binding domains (defined by amino
10 acid residues 3 through 111 of rsT4.111), followed by one OKT4-binding C-terminal domain (Figure 43).

pBiv.1 was transfected by electroporation into COS 7 cells to test expression of the bivalent rsT4 protein. Three days later, we tested the con-
15 ditioned medium of the transfected cells for the presence of the rsT4 bivalent protein by immuno-precipitation, followed by Western blot analysis of the precipitated protein. Both OKT4A and OKT4 were used for immuno-precipitation to determine that the
20 OKT4 epitope and at least one of the OKT4A epitopes had folded correctly. Both antibodies precipitated a protein of the predicted apparent molecular weight (60,000d) from the conditioned medium of the cells.

Bivalent rsT4 may be purified by immuno-
25 affinity purification from an OKT4 column and the purified protein may then be used to perform quantitative competition assays with rsT4.3. We believe that the bivalent molecule would demonstrate equivalent competition against rsT4.3 for OKT4 binding,
30 but significantly greater competition against monovalent rsT4 for OKT4A binding. The ability of bivalent recombinant soluble T4 to block syncytium formation may also be demonstrated in the C8166 fusion assay. We also believe that bivalent
35 recombinant soluble T4 would block syncytium formation at significantly lower concentrations than monovalent rsT4; based upon the higher

-93-

functional affinity of bivalent recombinant soluble T4 for gp120.

According to alternate embodiments of this invention, other methods for producing polyvalent rsT4 may be employed. For example, polyvalent rsT4 may be produced by chemically coupling rsT4 to any clinically acceptable carrier molecule, a polymer selected from the group consisting of Ficoll, polyethylene glycol or dextran, using conventional coupling techniques. Alternatively, rsT4 may be chemically coupled to biotin, and the biotin-rsT4 conjugate then allowed to bind to avidin, resulting in tetravalent avidin/biotin/rsT4 molecules. And rsT4 may be covalently coupled to dinitrophenol (DNP) or trinitrophenol (TNP) and the resulting conjugate precipitated with anti-DNP or anti-TNP-Igm, to form decameric conjugates with a valency of 10 for rsT4 binding sites.

Alternatively, a recombinant chimeric antibody molecule with rsT4 sequences substituted for the variable domains of either or both of the immunoglobulin molecule heavy and light chains may be produced. Because recombinant soluble T4 possesses gp120 binding activity, the construction of a chimeric antibody having two soluble T4 domains and having unmodified constant region domains could serve as a mediator of targeted killing of HIV-infected cells that express gp120.

For example, chimeric rsT4/IgG₁ may be produced from two chimeric genes -- an rsT4/human kappa light chain chimera (rsT4/C_{kappa}) and an rsT4/human gamma 1 heavy chain chimera (rsT4/C_{gamma-1}). Both C_{kappa} and C_{gamma-1} regions have been isolated from human recombinant DNA libraries, and each has been subcloned into animal cell selection vectors containing either the bacterial neo resistance or bacterial gpt markers

3743

89085519

-94-

for selection in animal cell hosts against the antibiotic G418 or mycophenolic acid, respectively.

To construct rsT4/C_{gamma-1} and rsT4/C_{kappa} chimeric genes, an rsT4 gene segment, including at least the secretory signal sequence and the N-terminal 110 amino acid residues of the mature rsT4 coding sequence and including a splice donor or portion thereof, is placed upstream of the gamma-1 and kappa constant domain exons. A suitable restriction enzyme may be used to cut within the intron downstream of the desired rsT4 coding sequence, thus providing a donor splice site. Subsequently, a suitable restriction enzyme is used to cut within the introns upstream of the kappa and gamma-1 coding regions. The rsT4 sequence is then joined to the kappa or gamma-1 constant region sequence, such that the rsT4 intron sequence is contiguous with the gamma-1 and kappa introns. In this way, an acceptor splice site is provided by the kappa or gamma-1 constant region intron. Alternatively, rsT4 chimeric genes may be constructed without the use of introns, by fusing a suitable rsT4 cDNA gene segment directly to the gamma-1 or kappa coding regions.

The rsT4/C_{gamma-1} and rsT4/C_{kappa} vectors may then be cotransfected, for example, by electroporation into lymphoid or non-lymphoid host cells. Following transcription and translation of the two chimeric genes, the gene products may assemble into chimeric antibody molecules.

Expression of the chimeric gene products may be measured by an enzyme-linked immunoadsorbant assay (ELISA) that utilizes monoclonal anti-T4 antibody OKT4A, as described infra, or in gp120 competition assays and radioimmunoassays, as described infra. Activity of the rsT4/IgG₁ chimeras may be measured by incubating them with HIV-infected cells in the presence of human complement, followed by quantitating

subsequent complement-mediated lysis of these cells. Alternatively, activity may be measured in HIV replication and HIV syncytium assays as described infra.

5 In order to determine if bivalent rsT4 has a greater potency than monovalent rsT4, we mixed OKT4, at various concentrations, together with a constant concentration of rsT4, so that the molar ratio of OKT4:rsT4 varied between 0.2 and 4. After preincubating the mixture overnight at 4°C, we added
10 aliquots to the HIV syncytium assay described infra. OKT4 has no observable effect in this assay when used alone. In addition, the concentration of recombinant soluble T4 chosen did not cause inhibition in this assay. Accordingly, we looked for indications that
15 the OKT4/rsT4 mixture was more potent than rsT4 alone. We observed that at ratios of OKT4:rsT4 greater than 0.2, partial to complete inhibition of syncytium formation occurred. We believe that under conditions where two rsT4 molecules are bound to 1 OKT4 molecule,
20 the greatest inhibitory effect should be found.

Thus, polyvalent, as well as monovalent forms of recombinant soluble T4 are useful in the compositions and methods of this invention.

25 Microorganisms and recombinant DNA molecules prepared by the processes of this invention are exemplified by cultures deposited in the In Vitro International, Inc. culture collection, in Linthicum, Maryland, on September 2, 1987, and identified as:

30 BG378: E.coli MC1061/pBG378
199-7: E.coli MC1061/p199-7
170-2: E.coli JA221/p170-2
EC100: E.coli JM83/pEC100
BG377: E.coli MC1061/pBG377
BG380: E.coli MC1061/pBG380
35 BG381: E.coli MC1061/pBG381

These cultures were assigned accession
3745 numbers IVI 10143-10149, respectively.

89085519

-96-

In addition, microorganisms and recombinant DNA molecules according to this invention are exemplified by cultures deposited in the In Vitro International, Inc. culture collection, in Linthicum,

5 Maryland, on January 6, 1988, and identified as:

BG-391: E.coli MC1061/pBG391

BG-392: E.coli MC1061/pBG392

BG-393: E.coli MC1061/pBG393

BG-394: E.coli MC1061/pBG394

10 BG-396: E.coli MC1061/pBG396

203-5 : E.coli SG936/p203-5.

These cultures were assigned accession numbers IVI 10151-10156, respectively.

15 Microorganisms and recombinant DNA molecules according to this invention are also exemplified by cultures deposited in the In Vitro International, Inc. culture collection, in Linthicum, Maryland, on August 24, 1988 and identified as:

211-11: E.coli A89/pBG211-11

20 214-10: E.coli A89/pBG214-10

215-7 : E.coli A89/pBG215-7

These cultures were assigned accession numbers IVI 10183-10185 respectively.

25 While we have hereinbefore described a number of embodiments of this invention, it is apparent that our basic constructions can be altered to provide othe embodiments which utilize the processes and compositions of this invention. Therefore, it will be appreciated that the scope of this
30 invention is to be defined by the claims appended hereto rather than by the specific embodiments which have been presented hereinbefore by way of example.

89085519

CLAIMS

We claim:

1. A DNA sequence selected from the group consisting of:

5 (a) the DNA inserts of p199-7, pBG377, pBG380, pBG381, p203-5, pBG391, pBG392, pBG393, pBG394, pBG395, pBG396, pBG397, p211-11, p214-10 and p215-7;

(b) DNA sequences which hybridize to one or more of the foregoing DNA inserts and which
10 code on expression for a soluble T4-like polypeptide; and

(c) DNA sequences which code on expression for a soluble T4-like polypeptide coded for on expression by any of the foregoing DNA inserts
15 and sequences.

2. The DNA sequence according to claim 1, wherein said DNA sequence (b) codes on expression for a soluble T4-like polypeptide which inhibits adhesion between T4⁺ lymphocytes and infective agents
20 which target T4⁺ lymphocytes and which inhibits interaction between T4⁺ lymphocytes and antigen presenting cells and targets of T4⁺ lymphocyte mediated killing.

3. A recombinant DNA molecule comprising
25 a DNA sequence selected from the group consisting of the DNA sequences of claim 1 or 2, said DNA sequence being operatively linked to an expression control sequence in said recombinant DNA molecule.

4. The recombinant DNA molecule according
30 to claim 3, wherein said expression control sequence is selected from the group consisting of the early or late promoters of SV40 or adenovirus, the lac system, the trp system, the TAC system, the TRC

-98-

system, the major perator and plomoter regions of phage λ , the contr 1 regions of fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase, the polyhedron promoter of the baculovirus system and the promoters of the yeast α -mating factors.

5. A unicellular host transformed with a recombinant DNA molecule selected from the group consisting of the recombinant DNA molecules of claim 3 or 4.

6. The host according to claim 5, wherein said host is selected from the group consisting of strains of E.coli, Pseudomonas, Bacillus, Streptomyces, fungi, animal cells, plant cells, insect cells and human cells in tissue culture.

7. A polypeptide coded for on expression by a DNA sequence selected from the group consisting of the DNA sequences of claim 1 or 2, said polypeptide being essentially free of other proteins of human origin.

8. The polypeptide according to claim 7, wherein said polypeptide is selected from the group consisting of a polypeptide of the formula $AA_{-23}-AA_{362}$ of Figure 3, a polypeptide of the formula AA_1-362 of Figure 3, a polypeptide of the formula $Met-AA_1-362$ of Figure 3, a polypeptide of the formula AA_1-374 of Figure 3, a polypeptide of the formula $Met-AA_1-374$ of Figure 3, a polypeptide of the formula AA_1-377 of Figure 3, a polypeptide of the formula $Met-AA_1-377$ of Figure 3, a polypeptide of the formula $AA_{-23}-AA_{374}$ of Figure 3, a polypeptide of the formula $AA_{-23}-AA_{377}$ of Figure 3.

89085519

-99-

9. The polypeptide according to claim 7, wherein said polypeptide is selected from the group consisting of a polypeptide of the formula AA₋₂₃-AA₁₈₂ of Figure 16, a polypeptide of the formula AA₁-AA₁₈₂ of Figure 16, a polypeptide of the formula Met-AA₁₋₁₈₂ of Figure 16, a polypeptide of the formula AA₋₂₃-AA₁₈₂ of Figure 16, followed by the amino acids asparagine-leucine-glutamine-histidine-serine-leucine, a polypeptide of the formula AA₁-AA₁₈₂ of Figure 16, followed by the amino acids asparagine-leucine-glutamine-histidine-serine-leucine, a polypeptide of the formula Met-AA₁₋₁₈₂ of Figure 16, followed by the amino acids asparagine-leucine-glutamine-histidine-serine-leucine, a polypeptide of the formula AA₋₂₃-AA₁₁₃ of Figure 16, a polypeptide of the formula AA₁-AA₁₁₃ of Figure 16, a polypeptide of the formula Met-AA₁₋₁₁₃ of Figure 16, a polypeptide of the formula AA₋₂₃-AA₁₁₁ of Figure 16, a polypeptide of the formula AA₁-AA₁₁₁ of Figure 16, a polypeptide of the formula Met-AA₁₋₁₁₁ of Figure 16, a polypeptide of the formula AA₋₂₃-AA₁₃₁ of Figure 16, a polypeptide of the formula AA₁-AA₁₃₁ of Figure 16, a polypeptide of the formula Met-AA₁₋₁₃₁ of Figure 16, a polypeptide of the formula AA₋₂₃-AA₁₄₅ of Figure 16, a polypeptide of the formula AA₁-AA₁₄₅ of Figure 16, a polypeptide of the formula Met-AA₁₋₁₄₅ of Figure 16, a polypeptide of the formula AA₋₂₃-AA₁₆₆ of Figure 16, a polypeptide of the formula AA₁-AA₁₆₆ of Figure 16, a polypeptide of the formula Met-AA₁₋₁₆₆ of Figure 16, or portions thereof.

10. The polypeptide according to claim 7, wherein said polypeptide is selected from the group consisting of a polypeptide of the formula AA₋₂₃-AA₃₆₂ of mature T4 protein, a polypeptide of the formula AA₁-AA₃₆₂ of mature T4 protein, a polypeptide of the formula Met-AA₁₋₃₆₂ of mature T4 protein, a polypep-

89085519

-100-

tide of the formula AA_{1-374} of mature T4 protein, a
 polypeptide of the formula $Met-AA_{1-374}$ of mature T4
 protein, a polypeptide of the formula AA_{1-377} of
 mature T4 protein, a polypeptide of the formula
 5 $Met-AA_{1-377}$ of mature T4 protein, a polypeptide of
 the formula $AA_{-23-AA_{374}}$ of mature T4 protein, a poly-
 peptide of the formula $AA_{-23-AA_{377}}$ of mature T4 pro-
 tein, or portions thereof.

11. The polypeptide according to claim 7,
 10 wherein said polypeptide is selected from the group
 consisting of a polypeptide of the formula $AA_{-23-AA_{182}}$
 of mature T4 protein, a polypeptide of the formula
 AA_1-AA_{182} of mature T4 protein, a polypeptide of the
 formula $Met-AA_{1-182}$ of mature T4 protein, a polypep-
 15 tide of the formula $AA_{-23-AA_{182}}$ of mature T4 protein,
 followed by the amino acids asparagine-leucine-
 glutamine-histidine-serine-leucine, a polypeptide of
 the formula AA_1-AA_{182} of mature T4 protein, followed
 by the amino acids asparagine-leucine-glutamine-
 20 histidine-serine-leucine, a polypeptide of the formula
 $Met-AA_{1-182}$ of mature T4 protein, followed by the
 amino acids asparagine-leucine-glutamine-histidine-
 serine-leucine, a polypeptide of the formula
 $AA_{-23-AA_{113}}$ of mature T4 protein, a polypeptide of
 25 the formula AA_1-AA_{113} of mature T4 protein, a polypep-
 tide of the formula $Met-AA_{1-113}$ of mature T4 protein,
 a polypeptide of the formula $AA_{-23-AA_{111}}$ of mature
 T4 protein, a polypeptide of the formula AA_1-AA_{111}
 of mature T4 protein, a polypeptide of the formula
 30 $Met-AA_{1-111}$ of mature T4 protein, a polypeptide of
 the formula $AA_{-23-AA_{131}}$ of mature T4 protein, a poly-
 peptide of the formula AA_1-AA_{131} of mature T4 protein,
 a polypeptide of the formula $Met-AA_{1-131}$ of mature
 T4 protein, a polypeptide of the formula $AA_{-23-AA_{145}}$
 35 of mature T4 protein, a polypeptide of the formula
 AA_1-AA_{145} of mature T4 protein, a polypeptide of the

3750

89085519

-101-

5 formula Met-AA₁₋₁₄₅ of mature T4 protein, a polypeptide of the formula AA₂₃₋₁₆₆ of mature T4 protein, a polypeptide of the formula AA₁₋₁₆₆ of mature T4 protein, a polypeptide of the formula Met-AA₁₋₁₆₆ of mature T4 protein, or portions thereof.

10 12. A method for producing a polypeptide selected from the group consisting of the polypeptides of any one of claims 7 to 11 comprising the step of culturing a unicellular host transformed with a recombinant DNA molecule selected from the group consisting of the recombinant DNA molecules of claim 3 or 4.

15 13. A pharmaceutical composition comprising an immunotherapeutic or immunosuppressive effective amount of a polypeptide selected from the group consisting of the polypeptides of any one of claims 7 to 11 and a pharmaceutically acceptable carrier.

20 14. A method for treating patients comprising the step of treating them in a pharmaceutically acceptable manner with a composition selected from the group consisting of the composition of claim 13.

15. The method according to claim 14, wherein the patient is treated by intramuscular injection of the composition.

25 16. A diagnostic composition for detecting or for monitoring the course of HIV infection comprising a diagnostic effective amount of a polypeptide selected from the group consisting of the polypeptides of any one of claims 7 to 11.

30 17. A method for detecting or for monitoring the course of HIV infection comprising the

step of employing as a diagnostic a composition selected from the group consisting of the compositions of claim 16.

18. A means for detecting or for monitoring the course of HIV infection comprising a composition selected from the group consisting of the compositions of claim 16.

19. A pharmaceutical composition comprising an immunotherapeutic or immunosuppressive effective amount of antibody to a polypeptide selected from the group consisting of the polypeptides of any one of claims 7 to 11 and a pharmaceutically acceptable carrier.

20. A method for treating patients comprising the step of treating them in a pharmaceutically acceptable manner with a composition according to claim 19.

21. The use of a polypeptide selected from the group consisting of the polypeptides of any one of claims 7 to 11 to purify HIV virus.

22. The use according to claim 20, wherein the HIV virus is purified from a biological sample.

23. A method for purifying HIV virus from a sample comprising the step of exposing the sample to a polypeptide selected from the group consisting of the polypeptides of any one of claims 7 to 11.

24. The method according to claim 22, wherein the sample is a biological sample.

25. A DNA sequence comprising the DNA insert of p170-2, said sequence coding on expression for a T4-like polypeptide.

5 26. A recombinant DNA molecule comprising a DNA sequence selected from the group consisting of the DNA sequence of claim 25, said DNA sequence being operatively linked to an expression control sequence in said recombinant DNA molecule.

10 27. A unicellular host transformed with a recombinant DNA molecule according to claim 26.

28. A polypeptide coded for on expression by a DNA sequence of claim 25, said polypeptide being essentially free of other proteins of human origin.

15 29. A pharmaceutical composition comprising an immunotherapeutic or immunosuppressive amount of a soluble protein receptor and a pharmaceutically acceptable carrier.

20 30. A method for treating patients comprising the step of treating them in a pharmaceutically acceptable manner with a pharmaceutical composition of claim 29.

25 31. A diagnostic composition for detecting or for monitoring the course of viral infection comprising a diagnostic effective amount of a soluble protein receptor.

32. A method for detecting or for monitoring the course of a viral infection comprising the step of employing as a diagnostic a diagnostic effective amount of a soluble protein receptor.

33. A means for detecting or for monitoring the course of a viral infection comprising a soluble protein receptor.

34. A DNA sequence selected from the group consisting of:

- (a) the DNA insert of pBiv.1;
- (b) DNA sequences which hybridize to the DNA insert of pBiv.1 and which code on expression for a polyvalent soluble T4-like polypeptide; and
- (c) DNA sequences which code on expression for a polyvalent soluble T4-like polypeptide coded for by the DNA insert of pBiv.1.

35. A recombinant DNA molecule comprising a DNA sequence selected from the group consisting of the DNA sequences of claim 34, said DNA sequence being operatively linked to an expression control sequence in said recombinant DNA molecule.

36. A unicellular host transformed with a recombinant DNA molecule according to claim 35.

37. A polypeptide coded for on expression by a DNA sequence selected from the group consisting of the DNA sequences according to claim 34, said polypeptide being essentially free of other proteins of human origin.

38. The polypeptide according to claim 7, wherein said polypeptide is polyvalent.

39. A method for producing a polyvalent polypeptide comprising the steps of:

- (a) culturing a unicellular host transformed with a recombinant DNA molecule according to claim 3 or 4 to produce a polypeptide; and

-105-

(b) coupling said polypeptid to a carrier to form a polyvalent polypeptide.

40. A DNA sequence comprising:

(a) a first portion comprising a DNA sequence coding for the constant region of an immunoglobulin light chain; and

(b) a second portion comprising a DNA sequence according to claim 1 or 2, or portions thereof, said second portion being joined upstream of said first portion.

41. A DNA sequence comprising:

(a) a first portion comprising a DNA sequence coding for the constant region of an immunoglobulin heavy chain; and

(b) a second portion comprising a DNA sequence according to claim 1 or 2, or portions thereof, said second portion being joined upstream of said first portion.

42. An expression vector comprising the DNA sequence according to claim 40.

43. An expression vector comprising the DNA sequence according to claim 41.

44. An expression vector comprising the DNA sequence according to claim 40 and the DNA sequence according to claim 41.

45. A method for producing a chimeric rST4/IgG₁ comprising the step of co-transfecting a host cell with the expression vector according to claim 42 and the expression vector according to claim 43.

3755

89085519

-106-

46. A method for producing a chimeric rsT4/IgG₁ comprising the step of transfecting a host cell with the expression vector according to claim 44.

47. A chimeric rsT4/IgG₁ produced by the method according to claim 45 or 46.

48. A pharmaceutical composition comprising an immunotherapeutic or immunosuppressive effective amount of a polypeptide according to claim 37 or 38.

49. A method for treating patients comprising the step of treating them in a pharmaceutically acceptable manner with a composition according to claim 48.

50. A diagnostic composition for detecting or for monitoring the course of HIV infection comprising a diagnostic effective amount of a polypeptide according to claim 37 or 38.


51. A pharmaceutical composition comprising an immunotherapeutic or immunosuppressive effective amount of a chimeric rsT4/IgG₁ according to claim 47.

52. A method for treating patients comprising the step of treating them in a pharmaceutically acceptable manner with a composition according to claim 51.

89085519

3756

1/93




3757

FIG. 1

89085519

2/93


FIG. 2



3758

89085519

FIG. 3



4/93

FIG 3(cont'd)

	H		B		S
	i		NBSN	F F	AaS
	n		laps		vui
	f		anlp	k k	a9n
	1		4222	1 1	261
			//		//

361 AAAACTCCAACCAGATAAAGATTCTGGGAAATCAGGGCTCCTTCTTAACATAAGGTCCAT

 TTTTGAGGTTGGTCTATTTCTAAGACCCTTTAGTCCCGAGGAAGAATTGATTTCAGGTA

 N S N Q I K I L G N Q G S F L T K G P S -

	S FH	H		S	REX
A	MNDa niHT	i	M	MANaSS	Splice
1	bdpu unhh	n	b	bvluit	Acceptor
u	oen3 DPaa	f	o	oaz9ny	
1	121A 2111	1	2	224611	
	//			///	

421 CCAAGCTGAATGATCGCGCTGACTCAAGAAGAAGCCTTCGGGACCAAGGAACTTTCCCC

 GGTTCGACTTACTAGCGCGACTGAGTCTTCTTCGGAAGCCCTGGTTCCTTTGAAAGGGG

 K L N D R A D S R R S L R D Q G N F P L -

S	H		H		S
BMDNa	i	A	i D M M	MM	MANMaM
cbpdu	n	f	nd b b	nn	nvnun
lone3	f	1	f e o o	1 1	lall91
1112A	1	2	1 1 2 2	1 1	121161
	//				/

481 TGATCATCAAGAATCTTAAGATAGAAGACTCAGATACTTACATCTGTGAAGTGGAGGACC

 ACTAGTAGTTCCTAGAATTCATCTTCTGAGTCTATGAATGTAGACACTTCACCTCCTGG

 I I K N L K I E D S D T Y I C E V E D Q -

S		M		E
i		a		c
n		e		o
1		1		b
/				

541 AGAAGGAGGAGGTGCAATTGCTAGTGTTCGGATTGACTGCCAACTCTGACACCCACCTGC

 TCTTCCTCCTCCACGTTAACGATCACAAGCCTAACTGACGGTTGAGACTGTGGGTGGACG

 K E E V Q L L V F G L T A N S D T H L L -

B B		BBE S	
s s		BsscNc	D
P P		aptosr	d
M M		niNRpF	e
1 1		221221	1
		///	

601 TTCAGGGGCAGAGCCTGACCCTGACCTGGAGAGCCCCCTGGTAGTAGCCCTCAGTGC

 AAGTCCCCGTCTCGGACTGGGACTGGAACCTCTCGGGGGGACCATCATCGGGGAGTCACG

 Q G Q S L T L T L E S P P G S S P S V Q -

M M	H	S		M	M	M DM	As
nn	i	t		b	b	n dn	lp
1 1	f	y		o	o	1 el	uB
1 1	1	1		2	2	1 11	12
						/	/

661 AATGTAGGAGTCCAAGGGGTAAAAACATACAGGGGGGGAAGACCCTCTCCGTGTCTCAGC

 TTACATCCTCAGGTTCCCATTTTGTATGTCCCCCTTCTGGGAGAGGCACAGAGTCG

 C R S P R G K N I Q G G K T L S V S Q L -

89085519

3760

720

FIG. 3(cont'd)

M	MM	A	HS	M	MM	M
b	ba	l	at	n	nn	n
o	ee	u	eu	l	ll	l
2	ll	l	3l	l	ll	l

A
l
u
l

AGAGGGGGAACAGGTGGAGTTCCTTCCACTCGCCTTTACAGTTGAAAAGCTGACGG
-----+-----+-----+-----+-----+ 900
TCTCCCCCTTGTCACCTCAAGAGGAAGGGTGAGCGGAATGTCAACTTTTCGACTGCC

E G E Q V E P S F P L A F T V E K L T G -

M	M	M	M	B	BES	P	S
m	m	b	b	sm	scc	ADNpPDa	s A
l	l	o	o	ta	tor	vrlusdui	l
1	1	2	2	Ee	NRF	aaaMse9n	u
				23	121	224111161	1

[illegible]

FIG. 3(cont'd)

	BES	S		S	S	BES
H	scc	MHMa		f	f	scc
p	tor	nanu		a	a	tor
h	NRF	le19		N	N	NRF
1	121	1316		1	1	121

1081 CTGGAAACCTCACCCTGGCCCTTGAAGCGAAACAGGAAAGTTGCATCAGGAAGTGAACC 1140

GACCTTTGGAGTGGGACCGGGAACCTTCGCTTTTGTCTTTCAACGTAGTCCTTCACTTGG

G N L T L A L E A K T G K L H Q E V N L -

	H	H D	A		M M		P S
	p	p d	l		n n		ADNNpPa
	h	h e	u		l l		vrllusu
	1	1 1	1		1 1		aaaaMs9
							2244116

1141 TGGTGGTGTATGAGAGCCACTCAGCTCCAGAAAATTGACCTGTGAGGTGTGGGGACCCA 1200

ACCACCACTACTCTCGGTGAGTCGAGGTCTTTTAAACTGGACACTCCACACCCCTGGGT

V V M R A T Q L Q K N L T C E V W G P T -

	S	S					
s	f	Df	MAM	DE	A	M M	T
i	a	da	nln	ds	l	n n	a
n	N	eN	lul	ep	u	l l	q
1	1	11	111	11	1	1 1	1

1201 CCTCCCCCTAAGCTGATGCTGAGCTTGAAACTGGAGAACAAAGGAGGCAAAGGTCTCGAAGC 1260

GGAGGGGATTGACTACGACTCGAAGCTTTGACCTCTTGTTCCTCCGTTTCAGAGCTTCG

S P K L M L S L K L E N K E A K V S K R -

		M M		DM		F	FD	M	H
		n n		ds		o	od	a	i
		l l		et		k	ke	e	f
		1 1		12		1	11	3	1

1261 GGGAGAAGCGCGGTGTGGGTGCTGAACCTTGAGGCGGGGATGTGGCAGTGTCTGCTGAGTGT 1320

CCCTCTTCCGCCACACCCACGACTTGGGACTCCGCCCTACACCGTCACAGACGACTCAC

E K A V W V L N P E A G M W Q C L L S D -

		P S		H		S		S
A		ADpPas		i		ANaS		HMNC
v		vrusui		n		vlui		pscr
a		aaMs9n		f		aa9n		apiP
1		221161		1		2361		2111

1321 ACTCGGGACAGGTCTGCTGGAATCCAACATCAAGGTTCTGCCCACATGGTCCACCCCGG 1380

TGAGCCCTGTCCAGGACGACCTTAGGTTGTAGTTCCAAGACGGGTGTACCAGGTGGGGCC

S G Q V L L E S N I K V L P T W S T P V -

	F		S			H			
	n		aH B	B	H	H	Ag	HMNH	M M M
	u		ua b	b	g	g	hi	psaa	n n b
	4		9e v	v	a	a	aD	spee	l l o
	H		63 1	1	1	1	21	2113	1 1 2

3762 TGCAGCCAATGGCCCTGATTGTGCTGGGGGCGTGGCCGGCCTCCTGCTTTTCATTGGGC 1440

ACGTCGGTTACCGGGACTAACACGACCCCCCGCAGCGCGCGGAGGACGAAAAGTAACCCG

Q P M A L I V L G G V A G L L L P I G L -

FIG. 3(cont'd)

M S f f B N HMBNN H H BAGHBHINN D
 b a a a l psaal ahihbanal d
 N N n a apnea naDaePra e
 2 1 1 1 4 21114 121112114 1
 TAGGCATCTTCTTCTGTGTCAGGTGCCGGCCACCGAAGGCGCCAAGCAGACGGATGTCTC
 ATCCGTTAGAAGAAGACACAGTCCACGGCCGTGGCTTCCGCGGTTCGTCTCGCCTACAGAG
 G I F F C V R C R H R R R Q A E R M S Q -
 S H B BB
 MNDaP F 1 D M M MH MMH Ms ssn HM M M
 bdpuo o n d n n bp bbb bp pps ps n n
 oen3k k f e l l oh ooh om lmp ap l l
 121A1 1 1 1 1 21 221 21 212 21 1 1
 AGATCAAGAGACTCCTCAGTGAGAAGAAGACCTGCCAGTGCCCTCACCGGTTTCAGAAGA
 TCTAGTTCTCTGAGGAGTCACTCTTCTTCTGACGGTCAACGGGAGTGGCCAAAGTCTTCT
 I K R L L S E K K T C Q C P H R P Q K T -
 N BE S S F BES
 MNDaP M M sChc MNDaX n ecc BM BM
 bnlbnp n n toar bdpuh u tor bn bn
 olaloH l l NRaP oen3o 4 NRP vl vl
 213121 1 1 1231 121A2 H 121 11 11
 Stop REX Stop
 CATGTAGCCCCATTTCAGGACGAGGCCAGGCAGATCCCACTGTAGCCTCCCCAGGTGT
 GTACATCGGGGTAAACTCCGTGCTCCGGTCCGTCTAGGGTGAAGTCCGAGGGGTCCACA
 C S P I G T R P G R S H L Q P P Q V S -
 F S SBES B
 nT AaS MNDaScCX HS s H M
 uh vui bdputorb at t e n
 Da a9n oen3NRfo eu X e l
 21 261 121A1212 31 1 3 1
 CTGCCCCCGGTTTCTGCCTGCGGACCAGATGAATGTAGCAGATCCCAAGGCTCTGGCCT
 GACGGGGCGCAAAGGACGGACGCCTGGTCTACTTACATCGTCTAGGGTCCGGAGACCGGA
 A P R F L P A D Q M N V A D P R P L A S -
 M M M BES SS
 n n n scc DHNPHDNaC
 l l l tor ralspscur
 1 1 1 NRP aeasapi9F
 1 1 1 121 234121161
 CCTGPTCGCCTCCTCTACAATTGCCATTGTTTCTCCTGGGTAGGCCCCGGCTTCACTG
 GGACAAGCGGAGGAGATGTTAAACGGTAACAAAGAGGACCCAATCCGGGGCCGAAGTGAC
 C S P P L Q P A I V S P G L G P G F T G -
 M M M M M
 nan nan M M
 lel lel l l
 111 111
 GTTGAGTGTGCTCTCTAGTTTCCAGAGGCTTAATCACACCGTCTCCACGCCATTTCCT
 CAACTCACAACGAGAGATCAAAGGTCTCCGAATTAGTGTGGCAGGAGGTGCGGTAAAGGA
 V L L S S P Q R L N H T V L H A I S F -

FIG. 3(cont'd)

M M M M
 a a a a
 e e e e
 1 1 1 1
 1801 TTTCCTTCAAGCCTAGCCCTTCTCTCATTATTTCTCTCTGACCCTCTCCCCACTGCTCAT 1860
 AAAGGAAGTTCCGATCGGGAAGAGAGTAATAAGAGAGACTGGGAGAGGGGTGACGAGTA
 S P K P S P S L I I S L * P S P H C S F -
 B BE SS S BE S
 aMDNscMacX a H scMcM H H
 mbpdtolurh u a tonrn n n
 HoneNRa3Po 9 e NRlPl 1 1
 1112124A12 6 3 12111 1 1
 / / / / / /
 1861 TTGGATCCCAGGGGAGTGTTCAGGGCCAGCCCTGGCTGGCTGGAGGGGTGAGGCTGGGTGT 1920
 AACCTAGGGTCCCCCTCACAAGTCCCCGGTCCGGACCGACCGACCTCCCACTCCGACCCACA
 G S Q G S V Q G Q P W L A G G * G W V S -
 N N BE P SSS
 1 1 ADNscMPhANPaacSS
 a a vrltonunvlsuurii
 3 3 aaNRlMlaas99Fnn
 2241211124166111
 / / / / / /
 1921 CTGGAAGCATGGAGCATGGGACTGTTCTTTTACAAGACAGGACCCTGGGACCACAGAGGG 1980
 GACCTTCGTACCTCGTACCCTGACAAGAAAATGTTCTGTCTGGGACCCTGGTGTCTCCC
 G S M E H G T V L L Q D R T L G P Q R A -
 S S S
 f f M M MNDPaxFF F
 a a n n bdpouhoo o
 N N 1 1 oenk3okk k
 1 1 1 1 1211A211 1
 / /
 1981 CAGGAACTTGCACAAAATCACACAGCCAAGCCAGTCAAGGATGGATGCAGATCCAGAGGT 2040
 GTCCTTGAACGTGTTTTAGTGTGTCTGGTTCGGTCAAGTTCCTACCTACGTCTAGGTCTCCA
 G T C T K S H S Q A S Q G W M Q I Q R P -
 F F
 n R B BB B M M HnN H AFNF
 u s b bb b n n pul p volo
 4 a v vv v 1 1 h4a h akak
 H 1 1 11 1 1 1 1H3 1 2141
 / /
 2041 TTCTGGCAGCCAGTACCTCCTGCCCCATGCTGCCCGCTTCTCACCCTATGTGGGTGGGAC 2100
 AAGACCGTCTGGTTCATGGAGGACGGGGTACGACGGGGCAAGAGTGGGATACACCCACCCTG
 L A A S T S C P M L P A S H P M W V G P -
 S H
 aS i 89085519 M M NR
 ui n n n ls
 9n f 1 1 1 aa
 61 1 1 1 31
 / /
 2101 CACAGACTCACATCCTGACCTTGCACAAACAGCCCTCTGGACACAGCCCCATGTACAG 2160
 GTGTCTGAGTGTAGGACTGGAACGTGTTGTCTGGGGAGACCTGTGTCTGGGGTACATGTGC
 Q T H I L T L H K Q P L W T Q P H V H G -

9/93

FIG. 3(cont'd)

R	F	F	M	M	F	F	M	M	D
a			n	n			n	n	d
e	k	k	l	l	k	k	l	l	e
3	1	1	1	1	1	1	1	1	1

2161 GCCTCAAGGGATGTCTCACATCCTCTGTCTATTTGAGACTTAGAAAAATCCTACAAGGCT 2220

CGGAGTTCCTACAGAGTGTAGGAGACAGATAAACTCTGAATCTTTTAGGATGTTCCGA

L K G C L T S S V Y L R L R K I L Q G W -

		S		BH
A	D	BMDNa	M	M ADBaGNS
c	d	cbpdu	n	n ldapias
c	e	lone3	l	l uenlApct
1	1	1112A	1	1 11221211

GGCAGTAGACAGAACTAAGATGATCATCTCCAGTTTATAGACCAGAACCAGAGCTCAGAG

2221 CCGTCATCTGTCTTGATTCTACTAGTAGAGGTCAAATATCTGGTCTTGGTCTCGAGTCTC 2280

CCGTCATCTGTCTTGATTCTACTAGTAGAGGTCAAATATCTGGTCTTGGTCTCGAGTCTC

Q * T E L R * S S P V Y R P E P E L R E -

M		HM	M	H	BES
a		ps	a	l	acc
e		ap	e	n	tor
1		21	1	f	NRP

AGCCTAGATGATTGATTACCAAGTGCCGGACTAGCAAGTGCTGGAGTCGGGACTAACC

2281 TCCGATCTACTAATAATGGTTCACGGCCTGATCGTTCACGACCTCAGCCCTGATTGGGT 2340

TCCGATCTACTAATAATGGTTCACGGCCTGATCGTTCACGACCTCAGCCCTGATTGGGT

A R * L I T K C R T S K C W S R D * P R -

P	S		F		H	H
ADNPas	B	B	n	B	M	MB
vrlusuf	b	b	u	s	n	ns
aaaMs9n	v	v	4	m	l	lm
2241161	1	1	H	1	1	11

GGTCCCTTGTGCCAAGTTCACCTGCTGCTCTTGAATGCAGGGACAAATGCCACACGGCT

2341 CCAGGGAACAGGGTTCAGGTGACGACGGGAGAACTTACGTCCCTGTTTACGGTGTGCCGA 2400

CCAGGGAACAGGGTTCAGGTGACGACGGGAGAACTTACGTCCCTGTTTACGGTGTGCCGA

S L V P S S T A A S * M Q G Q M P H G S -

	M		R		R
	a		s		s
	e		a		a
	1		1		1

CTCACCAGTGGCTAGTGGTGGGTACTCAATGTGTACTTTTGGGTTCACAGAAGCACACGA

2401 GAGTGGTCAACGATCACCACCCATGAGTTACACATGAAAACCCAAGTGTCTTCGTGTCTG 2460

GAGTGGTCAACGATCACCACCCATGAGTTACACATGAAAACCCAAGTGTCTTCGTGTCTG

H Q W L V V G T Q C V L L G S Q K H S T -

		S		HS	P	P	MFH
SN	N	ANas	D	at	o	o	non
tc	1	vlu1	d	eu	k	k	1kl
yo	a	aa9n	e	31	1	1	111
11	3	2461	1				

CCCATGGGAAGGGTCCATCTCAGAGAATTACGAGCAGGGATGAAGGCCTCCCTGTCTAA

2461 GGGTACCCTTCCCAGGTAGAGTCTCTTAAATGCTCGTCCCTACTTCCGGAGGGACAGATT 2520

GGGTACCCTTCCCAGGTAGAGTCTCTTAAATGCTCGTCCCTACTTCCGGAGGGACAGATT

3765 H G K G P S Q R I Y E Q G * R P P C L K -

89085519


10/93

FIG. 3(cont'd)

F M M N H H M M M M M M
 o n n s i M M M b b
 k n n p n n n a o o
 l 1 1 B f l l e 2 2
 1 1 1 2 1 1 13 2 2
 2521 AATCCCTCCTTCATCCCCCGCTGGTGGCAGAATCTGTTACCAGAGGACAAAGCCTTTGGC 2580
 TTAGGGAGGAAGTAGGGGGCGACCACCGTCTTAGACAATGGTCTCCTGTTTCGGAACCG
 S L L H P P L V A E S V T R G Q S L W L -
 H BH
 i H A sqN P B B M
 n h l p l s s s a
 P a u l a p t m m e
 1 1 1 212 1 1 13
 //
 2581 TCTTCTAATCAGAGCGCAAGCTGGGAGCACAGGCACTGCAGGAGAGAATGCCCAGTGACC 2640
 AGAAGATTAGTCTCGCGTTTCGACCCTCGTGTCCGTGACGTCCTCTCTTACGGGTCAGTGG
 F * S E R K L G A Q A L Q E R M P S D Q -
 BES
 M scc M M A M M H A M M N
 a cor n n l n n n l a n h
 e NRP 1 1 u l l l u e l e
 3 121 1 1 11 1 1111
 //
 2641 AGTCACTGACCCCTGTGCAGAACCTCCTGGAAGCGAGCTTTGCTGGGAGAGGGGGTAGCTA 2700
 TCAGTGACTGGGACACGTCCTTGGAGGACCTTCGCTCGAAACGACCCTCTCCCCCATCGAT
 S L T L C R T S W K R A L L G E G V A S -
 P S BES H
 D ADMONpPaS M M HscCH i H
 d 1 d v r n l u s u i n n p t o r p n h
 e a e a a l l a m s 9 n 1 1 h N R F h P a
 1 4 1 221141161 1 1 11211 1 1
 //
 2701 GCCTGAGAGGGGAACCCCTCTAAGGGACCTCAAAGGTGATTGTGCCAGGCTCTGCGCCTGCC 2760
 CGGACTCTCCCTTGGGAGATTCCCTGGAGTTTCCACTAACACGGTCCGAGACGCGGACGG
 L R G N P L R D L K G D C A R L C A C P -
 M M M M M M M M
 n n n n n n n n
 1 1 1 11 1 1 1 1
 1 1 1 11 1 1 1 1
 2 32
 2761 CCACACCCTCCCTTACCCCTCCTCAGACCATTTCAGGACACAGGGAATCAGGGTTACAAA 2820
 GGTGTGGGAGGGAATGGGAGGAGGTCTGGTAAGTCCTGTGTCCCTTTAGTCCCAATGTTT
 T P S L T L L Q T I Q D T G K S G L Q I -
 S B S
 MNDa D M aMDMNaX H M H M M M D
 b d p u d b m b p b d l u h p n p n n n d
 o e n 3 e o H o n o e a 3 o h 1 h 1 1 1 e
 121A 1 2 111224A2 1 1 11 1 1 1
 //
 2821 TCTTCTTGATCCACTTCTCTCAGGATCCCTCTCTTCTTACCCTTCCTCACCACCTTCCTT 2880
 AGAAGAAGTCTAGGTGAAGAGAGTCTTAGGGGAGAGAAGGATGGGAAGGAGTGGTGAAGGGA
 F L I H F S Q D P L S S Y P S S P L P S -


12/93

FIG. 4



13/93

FIG. 5



* numbers in parentheses refer to PBL T4 cDNA coordinates

14/93

FIG. 6


AMINO ACID SEQUENCE COMPARISON AT POSITIONS 36 AND 231 OF T4

Position No.	Maddon et al.	PBL Clone	Rex Clone	Genomic	Mouse	Sheep
3	N AAC	K AAG	-	-	K AAG	K -
64	W TGG	R CGG	W TGG	W TGG	W TGG	-
231	F TTT	S TCT	F TTT	-	F TTC	-

3770

89085519

15/93



KVLPTWSTPVQPMALIVLGGVAGLLFLFIGLGFVRCRHR

C. Mutant




FIG. 7A








FIG.7B



18/93

FIG. 8B*p178A (full length T4)**arg → trp*↓ *Ava I, EcoRI, Isolate 1.2 kbp fragment*↙ *Linkers 46/47, 48/49*↘ *pBG 368-EcoRI/Bgl II*↓ *Ligase*

(EcoRI)


↓ *EcoRI/SalI, Isolate 1.3 kbp fragment*↙ *Linkers 50/51*↘ *pBG 368-EcoRI/Bgl II*

3774

89085519

19/93

"Met-perfect" cassette for *E. coli* expression of *rsT4*




3775

FIG. 8C


20/93

FIG.8D



21/93

FIG. 9A



22/93




FIG. 9B

23/93

FIG. 9C

198-2

	180					<u>StuI</u>					
CTA	GCT	TTC	CAG	AAG	GCC	TCC	AGC	ATA	GTC	TAT	

deletion

203-5

	180		182							
CTA	GCT	TTC	CAG	AAC	CTC	CAG	CAT	AGT	CTA	TAA
leu	ala	phe	gln	asn	leu	gln	his	ser	leu	stop

← T4 frameshift →

89085519

24/93

*FIG.10*48

5' CTG CCC ACA TGG TCG ACC CCG GTG CAG CCA ATG TGA 3'

49

5' GAT CTC ACA TTG GGT GCA CCG GCG TCG ACC ATG T 3'

50

5' TCG ACC CCG GTG CAG CCA ATG GCC CTG ATT TGA 3'

3' GG GGC CAC GTC GGT TAC CCG GAC TAA ACT CTA G 5'

5141

5' GAA GAA GGT TGT GCG ACC AAG 3'

34

5' TCG GGA CAG GTC CTG CTG GAA TCC AAC ATC AAG TGA A 3'

3' CTG TCC ACG ACG ACC TTA GGT TGT AGT TCA CTT CTA G 5'

3555

5' CAG CCA CCC AAG GAA ACA AAG TCG 3'

46

5' TCG GGA CAG GTC CTG CTG GAA TCC AAC ATC AAG GTT 3'

47

5' GCG CAG AAC CTT GAT GTT GGA TTC CAG CAG GAC CTG TC 3'

89085519

25/93

*FIG. 10(cont'd)*54

5' AGC TTC GAC TCG AGG ATG CAG GGA AAC AAA GTG GTG 3'

← 57 58 →
5' AATTCTTACACTTAACTTAAATTCGCTAACTTTATAGATTACAAACTT
GAATGTGAATCAATTTAAGGATTGAAATATCTAATCTTTTGAA
← 59 →

ACGAAATCGATTTCATGG

TCCTTTAGCTTAAAGGTTACCTAG 5'
← 60 →

89085519

3781

27/93

FIG. 11(cont'd)

361 GTGTTGGATTGACTGCCAACTCTGACACCCACCTGCTTCAGGGCCAGACCTGACCCCTG 420
 CACAAGCCTAACTGACCGTTGAGACTGTGGGTGGACCAAGTCCCCGTCTCGGACTGGAC
 ValPheGlyLeuThrAlaAsnSerAspThrHisLeuLeuGlnGlyGlnSerLeuThrLeu -

BBZ S
 S BaccNc D M H S
 c aptosr d n i c
 y nLRpF e l z y
 l 221221 1 1 1 1
 / / / /
 421 ACCTTGGAGAGCCCCCTGGTAGTACCCCTCAGTCCAATGTAGGAGTCCAAGCGGTAAA 480
 TGGAACTCTCTGGGGGGGACCATCATCGGGGAGTCAGGTACATCCTCAGGTTCCTCCAT--
 ThrLeuGluSerProProGlySerSerProSerValGlnCysArgSerProArgGlyLys -

N BBZH SB BES
 M M D AsP A BaccGNScsS B Nacc
 b n d lpv l aptoisarts a ltor
 o l e uBu u nLRAPcFXc a aNRP
 2 1 1 122 1 2212121111 1 4:21
 // / / / / / / / /
 481 AACATACAGGGGGGGAAGACCTCTCCGTGTCTCAGCTGGAGCTCCAGGATAGTGGCACC 540
 TTGTATGTCCCCCTCTCTGGGAGAGGCACAGAGTCCAGCTCCAGGTCTTATCACCCTGG
 AsnIleGlnGlyGlyLysThrLeuSerValSerGlnLeuGluLeuGlnAspSerGlyThr -

N
 Ns
 lp
 aH
 31
 /
 541 TGGACATGCACTGTCTTCCAGAACAGAGAAGCTGGAGTTCAAAATAGACATCGTGGTG 600
 ACCTGTACGTGACAGAACGTCTTGGTCTTCTTCCACCTCAAGTTTATCTGTAGCACCAC
 TrpThrCysThrValLeuGlnAsnGlnLysLysValGluPheLysIleAspIleValVal -

M A HS M M
 a l ac n n
 e u eu l l
 l l 31 1 1
 /
 601 CTAGCTTTCAGAAAGGCTTCAGCATAGTCTATAAGAAAGAGGGGGAACAGGTGGAGTTC 660
 GATCGAAAGGTCTTCCGGAGGTGCTATCAGATATCTCTTCTCCCCCTTGTCCACCTCAAG
 LeuAlaPheGlnLysAlaSerSerIleValTyrLysLysGluGlyGluGlnValGluPhe -

A A M
 l l n
 u u l
 l l l
 661 TCCTTCCCACTCGCCTTACAGTTGAAAAGCTACGGGAGTGGCGAGCTGTGGTGGCAG 720
 AGCAAGCGGTGACCCGAAATGTGAACTTTCGACTGCCCGTACCGCTCGACACCACCGTC
 SerPheProLeuAlaPheThrValGluLysLeuThrGlySerGlyGluLeuTrpTrpGln -

3783

89085510

28/93

FIG. 11(cont'd)

	P	S	
MH	Mf	MNDa	M
np	nl	bdpu	b
lh	lm	en3	o
ll	ll	121A	2

721 CCGGAGAGGGCTTCTCTCTCCAAGTCTTGGATCACTTTGACCTGAAGAACAAGGAAGTC
 ----- 750
 CGCCTCTCCCGAAGGAGGAGGTTCAGAACCTAGTGCAAACTGGACTTCTTGTCTCTTCAC
 AlaGluArgAlaSerSerSerLysSerTrpIleThrPheAspLeuLysAsnLysGluVal -

	BES	P	S		
SH	scc	ADNPPDaS	A		
ta	cor	vrllusui	l	A	H
Ze	HRP	aaaMse9n	u	u	p
23	121	22411161	1	1	h

781 TCTGTAAACCGGTTACCCAGGACCCTAAGCTCCAGATGGGCAAGAAGCTCCCGCTCCAC
 ----- 840
 AGACATTTTCCCAATCGGTCCTGGGATTCGAGGTCTACCCGTTCTTCGAGGGCGAGGTC
 SerValLysArgValThrGlnAspProLysLeuGlnMetGlyLysLysLeuProLeuHis -

	BES					BES	S
M	scc	HS	D	M	H	scc	MHa
n	cor	ac	d	n	p	cor	baa
1	HRP	eu	e	1	h	HRP	le9
1	121	31	1	1	1	121	136

841 CTCACCCCTGCCCCAGGCTTGGCTCAGTATGCTGGCTCTGGAAACCTCACCCCTGGCCCTT
 ----- 900
 CAGTGGGACGGGCTCCGGAACGAGTCATACGACCAGACCTTTGCACTGGGACCGGGAA
 LeuThrLeuProGlnAlaLeuProGlnTyrAlaGlySerGlyAsnLeuThrLeuAlaLeu -

		S	BES				
		f	scc			H	D
		a	cor			p	d
		M	HRP			h	e
		1	121			1	1

901 GAACCGAAAACAGGAAGTTGCATCAGGAAGTGAACCTGGTGGTATGAGAGCCACTCAG
 ----- 960
 CTTCGCTTTTGTCTTTCAACGTAGTCTCTCACTTGGACCACCACTACTCTCGGTGAGTC
 GluAlaLysThrGlyLysLeuHisGlnGluValAsnLeuValValMetArgAlaThrGln -

		P	S	S			
M		ADNPPa3	f	D	MA	D	
n		vrllusui	a	d	nl	d	
1		aaaMse9n	M	e	lu	e	
1		22441161	1	1	11	1	

961 CTCAGAAAAATTTGACCTGTGAGGTGTGGGGACCCACCTCCCCCTAAGCTGATGCTGAGT
 ----- 1020
 GAGGTCTTTTAACTGGACACTCCACACCCCTGGGTGGAGGGGATTGCACTACCACTCA
 LeuGlnLysAsnLeuThrCysGluValTrpGlyProThrSerProLysLeuMetLeuSer -

89085519

	M			M
	n			n
	1			1
	1			1

1021 TTGAAACTGGAGAACAAGGAGGCCAAAGGTCTCGAAGCGGGAGAAGCGGTGTGGGTGCTG
 ----- 1080
 AACTTTGACCTCTTGTCTCTCCGTTTCCAGAGCTTCGCCCTCTTCCGCCACACCCACGAC
 LeuLysLeuGluAsnLysGluAlaLysValSerLysArgGluLysAlaValTrpValLeu -

3784

29/93

FIG. 11(cont'd)

DM	P	D	M	I	A	P	S	H
ds		d	a	n	v	ADPpAs		1
et	k	e	e	f	a	vrusui		n
12	1	1	3	1	1	aaMs9n		f
/						221161		1
						// //		

1081 AACCTGAGCGCGCGGATGTGCCAGTGTCTGCTGAGTGACTCGGGACAGGTCTCTGCTGGAA
 TTGGGACTCCGCCCCCTACACCGTCACAGACGACTCACTGAGCCCTGTCCAGGACGACCTT 1140

AsnProGluAlaGlyMetTrpGlnCysLeuLeuSerAspSerGlyGlnValLeuLeuGlu -

	H	S	P	S	P
	NSAIT	MDMc	n	BMDNaX	TP
	lacna	pser	u	bbspduh	us
	alccq	apiP	4	vonajo	4c
	31121	2111	H	1112A2	H1
	/	///		///	

1141 TCCAACATCAAGGTTCTGCCACATGGTGACCCCGGTCCAGCCAAATGTGAGATCTCTGCA
 AGGTTGTAGTTTCAAGACGGGTGTACCAGCTGGGGCCACGTCCGTACACTTAGGACCT 1200

SerAsnIleLysValLeuProThrTrpSerThrProValGlnProMetEndAspProAla -
 STOP

	P	S	P	S	S
A	MB	ADNNpHPAs	ADNpPas	MDNaX	
1	nb	vrlluasui	vrllusui	bcpuh	
u	lv	aaaaMes9n	aaMs9n	oenjo	
1	11	224411161	2241161	121A2	
	///	///	///	///	

1201 GCCCAGCTTGGGGACCTTAGAGGTCCCTTTAATTTGAATGGGAGATCCCAATCTT
 CCGGTGGAACCCCTGGGATCTCCAGGGGAAAAATAAACTTAACCTCTAGGGTAAAGA 1260

AlaGlnLeuGlyAspProArgGlyProLeuPheTyrPheGluLeuGlyAspProAsnSer -

Clal

N	A	CT	AMN	A
1	1	14	1ah	1
a	u	aa	uee	u
3	1	11	111	1

1261 CATGTTTACAGCTTATCATCGATAAGCTAGCTTTAATCCGGTAGTTTATCACAGTTAA
 GTACAAACTGTGGAATAGTAGCTATTCGATCGAAATTACGCCATCAAACTAGTGTCAATT 1320

HisValEndGlnLeuIleIleAspLysLeuAlaLeuMetArgEndPheIleThrValLys -

89085519

B	H	P	I	H	HB
a	1	o	n	b	pa
n	a	k	p	a	nn
1	4	1	1	1	11

1321 TTGCTAACGCAGTCAGGCACCGTGTATGAAATCTAACAAATCCGCTCATCGTCACTCTCGG
 AACGATTCCGTCAGTCCGTGGCACATACTTTAGATTGTTACCGCAGTAGCAGTAGGAGCC 1380

LeuLeuThrGlnSerGlyThrValTyrGluIleEndGlnCysAlaHisArgHisProArg -

S	BES	P	HM	R	S	S
N	MMf	scc			MDNaHC	
1	ana	tor	o	s	pecuar	
a	eln	NRF	k	a	api9eP	
4	311	121	1	21	211631	
	//	/		/	///	

1381 CACCGTCACCCCTGGATGCTGTAGGCATAGGCTTGGTTATGCCGGTACTGCCGGGCGCTCTT
 GTGGCAGTGGGACCTACGACATCCGTATCCGAACCAATACGGCCATGACGGCCCCGGAGAA 1440

3785 HisArgHisProGlyCysCysArgHisArgLeuGlyTyrAlaGlyThrAlaGlyProL u -

30/93

FIG. 11(cont'd)

M	E		S		P	H	S
n	C		MBf		n NM	i HHf	
1	R		aba		u ha	n ha	
1	V		evM		4 ee	P aeM	
			311		H 11	1 121	

1441 GCGGGATATCGTCCATTCCGACAGCATCGCCAGTCACTATGCGGTGCTGCTAGCGCTATA

CGCCCTATAGCAGGTAAGGCTGTCTAGCGGTCACTGATACCGCACGACGATCGCGATAT 1500

AlaGlyTyrArgProPheArgGlnHisArgGlnSerLeuTTPArgAlaAlaSerAlaIle -

H		BH		P
1PHM		sgM		C Hn
nshs		pis		f au
Ppat		lap		r ea
1111		212		1 3H

1501 TCGGTTGATGCAATTTCTATGCCACCCGTTCTCGGAGCACGTGTCGACCCGCTTTGGCCG

ACGCAACTACGTTAAAGATACCGGTGGGCAAGACCCCTCGTGACAGGCTGGCGAAACCGGC 1560

CysValAspAlaIleSerMetArgThrArgSerArgSerThrValArgProLeuTTPPro -

P			P	S
n		M	nMDat	N
u		1	ubdpuh	1
4		a	Doenja	a
H		4	2121A1	3

1561 CCGCCCAGTCTCTGCTCGCTTCGCTACTTCGAGCCACTATCGACTACCGGATCATGGCCAC

GGCGGGTCAGGACGAGCGAAGCGATGAACCTCGGTGATAGCTGATCGCCTAGTACCCCTG 1620

ProProSerProAlaArgPheAlaThrTTPSerHisTyrArgLeuArgAspHisGlyAsp -

H		S		H	H
i		CHHHHMMf		HMBAGHHHIN	
n	HM	fgappsa		psahihbana	
2	ps	rcaahpeN		apnaDaeePr	
1	ap	11321111		2112111211	
	21	/	/	/	/

1621 CACACCCGTCCTGTGGATTCTCTACGCCGGACCCATCGTGGCCGGCATCACCGGCCCCAC

GTGTGGCCAGGACACCTAAGAGATCGCCCTCGGTAGCACGGCCCGTAGTGGCCCGGGTG 1680

HisThrArgProValAspSerLeuArgArgThrHisArgGlyArgHisHisArgArgHis -

S		H	H		S	B
Nf		BAGHHHINM	H		MNDa	BaMN
1a		ahihbanal	p		bdpu	apbs
aN		naDaeePra	h		oen3	olop
41		121112114	1		121A	2222
//		/	/	/	/	/

1681 AGGTGCGGTTGCTGCGGCTTATATCGCCGACATCACCGATGGGGAAGATCGGGCTCGCCA

TCACGCCAAGCACCGCGGATATAGCGGCTGTAGTGGCTACCCCTTCTAGCCCCAGCGGT 1740

ArgCysGlyCysTTPArgLeuTyrArgArgHisHisArgTTPGlyArgSerGlySerPro -

B		H		S		S
BeN	N	i	HH		DHNPa	C HHNC
aps	1	n	ha		zalsu	f apct
nlp	a	P	ae		aeas9	r eapif
222	3	1	12		23416	1 32111
//		/	/	/	/	/

1741 CTTGCGGGCTCATGAGCGCTTGTTCGGCGTGGGTATCGTGGCAGGCCCCGTGGCCGGGGG

GAAGCCCCGAGTACTCGCGAACAAGCCGACCCATACCACCGTTCGGGGCACCGGCCCCC 1800

LeuArgAlaHisGluArgLeuPheArgArgGlyTyrGlyGlyArgProArgGlyArgGly -

89085519

31/93

FIG. 11(cont'd)

	H	H		P		BH	
	BAqHBHINN			n	n	sqN	H
	shihbanal			u	u	pls	a
	naDa ePra			4	4	lap	e
	121112114			H	H	212	3
	/	////				//	

1801 ACTGTTGGGCGCCATCTCCTTGCACGCACCATTCCTTGGCGCGCGGGTCTCAACGGCGCT 1860

TCACAACCCGCGGTAGAGGAACGTGCGTGCTAAGGAACGCCCGCCCGCCAGAGTTGCCGGA

ThrValGlyArgHisLeuLeuAlaArgThrIleProCysGlyGlyGlyAlaGlnArgPro -

	B	M		P		H	S
	b	n		n		i	2
	v	l		u		g	a
	1	1		4		4	H
	1	1		H		1	1

1861 CAACCTACTACTGGGCTGCTTCTTAATGCAGGAGTCCCATTAAGGCAGACCGTCTCCGAT 1920

GTTGCGATGATGACCCGACGAAGGATTACGTCTCTACCGTATTCCTCTCTCCAGCAGGCTA

GlnProThrThrGlyLeuLeuProAsnAlaGlyValAlaEndGlyArgAlaSerSerAsp -

				A	HM	H P	
				1	ps	i nHT	N
				u	ap	n uhh	1
				1	21	P Daa	a
						1 211	3
				/		//	

1921 GCGCTTGAGAGCCTTCAACCCAGTCAGCTCTTCCGGTGGCGCGCGGGCATGACTATCGT 1980

CGGGAACCTCTCGGAAGTTGGGTCAGTCGACGAAGGCCACCCCGCGCCCGTACTGATAGCA

AlaLeuGluSerLeuGlnProSerGlnLeuLeuProValGlyAlaGlyHisAspTyrArg -

	P						P H
	n	M		N			B N HDN n i HH
	u	b		1			a l psa u n ha
	4	o		a			n a spe 4 P ae
	H	2		3			1 4 211 H 1 12
							//

1981 CCGCGCACTTATGACTGTCTTCTTTATCATGCAACTCGTAGGACAGGTGCCCGCCAGCGCT 2040

CGCGCGTGAATACTGACAGAAGAAATAGTACGTTGAGCATCTGTCTCCAGCGCCGTCCGCA

ArgArgThrTyrAspCysLeuLeuTyrHisAlaThrArgArgThrGlyAlaGlySerAla -

	BH		S		H P		S
	bn		Aas		i nHT		MNDa H
	vl		vul		n uhh		bdpu a
	11		asn		P Daa		oen3 e
			261		1 211		121A 3
			//		//		//

2041 CTGGGTCAATTTTCGGCGAGGACCGCTTTCGCTGGAGCGCGACGATGATCGGCTGTTCGCT 2100

GACCCAGTAAAAGCCGCTCTCGCGAAGCGACCTCGCGCTGCTACTAGCCGGACAGCGA

LeuGlyHisPheArgArgGlyProLeuSerLeuGluArgAspAspAspArgProValAla -

	H					S
	i			M	M	AaNS
	n			n	a	vuli
	f			1	e	asn
	1			1	3	2641
						//

37872101 TCGGTATTTCGGAATCTTCACGCCCTCGCTCAAGCCTTCGTCAGTGGTCCCGCCACCAA 2160

ACGCCATAAGCCTTAGAACGTGCGGGAGCGAGTTCCGAAGCAGTGACCAGCGCGCTGCTT

CysGlyIleArgAsnLeuAlaArgProArgSerSerLeuArgHisTrpSerArgHisGln -

89085519

FIC. 11(cont'd)

M		H		HGN	B	N	CENK	FM	niHT	H	M
a		a		psa	g	1	fauam	unhh	g	a	
e				ape	1	a	rg4ea	DPaa	a	e	
2		3		211	1	3	11H33	2111	1	2	

2161 ACGTTTCGGCGAGAACCGAGCCATTATCGCCGGCATGGCCGGCCGACGGCTGGGCTACGT

 TGCAAAGCCGCTCTTCGTCGGTAATAGCGCCGCTACCGCCGGCTGCGCGACCCGATGCA 2220
 ThrPheArgArgGluAlaGlyHisTyrArgArgHisGlyGlyArgArgAlaGlyLeuArg -

	M	P	P	H	H	FM	H		H
	n	urh	uh	g	a	ob	n		p
	1	Dua	Da	a	e	ko	2		a
	1	211	21	1	3	12	1		2

2221 CTTGCTGGCGCTTCGGACCGGAGGCTGGATGGCCCTTCCCAATTATGATTCTTCTCGCTTC

 GAACGACCGCAAGCGCTCGGCTCCGACCTACCGGAAGCGGTAATACTAAGAAGACCGAAG 2280
 LeuAlaGlyValArgAspAlaArgLeuAspGlyLeuProHisTyrAspSerSerArgPhe -

SP	S	P		B	BES
M in	f	nt	P	sn	scC
s au	a	uh	o	pl	tor
p M4	M	Da	k	Ma	NRP
1 1H	1	21	1	13	121

2281 CCGCGGCATCGGGATCGCCCGCTTCAGGCCATCGCTGCCAGGCAGGTAGATGACGACCA

 GCGCGCGTAGCCCTACGGCGCGCAACGTCCGGTACGACAGGTCCGTCCATCTACTGCTCGT 2340
 ArgArgHisArgAspAlaArgValAlaGlyHisAlaValGlnAlaGlyArgEndArgPro -

	A	S	FP		S	S
	1	MDa	nnT		TDa	AsS
	u	bcpa	uuh		abcpa	vui
	1	oen3	D4a		qoen3	a9n
		121A	2H1		1121A	261

2341 TCAGGGACAGCTTCAAGGATCGCTCGCGGCTCTTACCAGCCTAACTTCGATCACTGGACC

 AGTCCCTGTGGAAGTTCTTAGCGAGCGCCGAGAATCGTCCGATTGAAGCTAGTGACCTGG 2400
 SerGlyThrAlaSerArgIleAlaArgGlySerTyrGlnProAsnPheAspHisTrpThr -

N	S		P		BH		M
s MNDaM			n	B	M	sqN	M
p bcpa			u	g	n	pis	1
B oen3e			4	1	1	lap	a
2 121A3			H	1	1	212	3

2401 GCTGATCGTCACCGCGATTATCGCCGCTCGCGGAGCACATGGAACGGGTGGCATGGAT

 CGACTAGCAGTCCCGCTAAATACGGCGGAGCCGCTCGTGTACCTTGCCCAACCGTACCTA 2460
 AlaAspArgHisGlyAspLeuCysArgLeuGlyGluHisMetGluArgValGlyMetAsp -

	H	P	H		P	P		N	N	HGN
	BAGHBnHINN			H	ntM	nt		1	1	psc
	ahihbusnal			g	uhn	uh		a	a	api
	naDae4ePfa			a	Dal	Da		3	4	211
	12111H2114			1	211	21				

2461 TGTAGGCGCCGCCCTATACCTTGTCTCCCTCCCCCGGTTGCGTCCGGTGCATGGAGCCG

 ACATCCGCGCGCGGATATGGAACAGACCGAGCGCGCCAACGCAGCGCCACGTACCTCGGC 2520
 CysArgArgArgProIleProCysGluArgProArgValAlaSerArgCysMetGluPro -

FIG. 11(cont'd)

S S P H
 aHc T M HON nB N H M I
 uar a n psa ua l p n n
 9eP q l ape 4n a b l f
 631 l l 211 H1 4 l l l
 // //

2521 GGGCACCTCGACCTGAATCGAAGCCGGCGGCACCTCGCTAACGGATTACCACTCCAAGA
 ----- 2580
 CCGGTGGAGCTGGACTTACCTTCGGCCGGCGTGGACCGATTGCCTAAGTGGTGAGGTTCT

GlyHisLeuAspLeuAsnGlySerArgArgHisLeuAlaAsnGlyPheThrThrProArg -

P H
 2 1
 1 1
 M a
 1 4
 H 1P3HM
 1 1 nsehs
 1 1 Ppmat
 1 1 11111
 1 1 11

2581 ATTGGAGCCAATCAATTCTTGGCGAGAACTGTGAATCGCCAAACCAACCTTCGGCAGAAC
 ----- 2640
 TAACCTCGGTAGTTAAGAAGCCCTCTTGACACTTACCGGTTTGGTTGGGAACCGTCTTC

IleGlyAlaAsnGlnPheLeuArgArgThrValAsnAlaGlnThrAsnProTrpGlnAsn -

P P P P S S
 nT n n nnBITH A B f BMDNa A
 uh u u unbnhh v b a cbpdu l
 Da a 4 D4vPaa a v M lone3 u
 21 H H 2H1111 1 1 1 1112A 1
 // //

2641 ATATCCATCGCGTCCGCCATCTCCAGCAGCCCGACCGCCCATCTTCGGCGGATCATCAG
 ----- 2700
 TATACGTAGCCGAGCGCGTAGAGGTCTGTCGGCGTCCCGCCGTAGACCCCTTACTAGTC

IleSerIleAlaSerAlaIleSerSerSerArgThrArgArgIleSerGlyAspAspGln -

FN FHP H H M FN S
 nsPF nlnHMT H M MnaA HONc B
 upvo unubhnh P p n lupl pect b
 4Buk DPDaaia h h 1 a4Hu apif v
 H221 2121111 1 1 1 3H11 2111 1
 // // //

2701 CTGCCTCGCGCGTTTCGGTCAATGACGGTGAACCTCTGACACATCCAGGTCCCGGAGAC
 ----- 2760
 GACCGAGCGCGCAAGCCACTACTGCCACTTTTGGAGACTGTGTACGTGAGGGCCCTCTG

LeuProArgAlaPheArgZndZndArgZndLysProLeuThrHisAlaAlaProGlyAsp -

M A S S H P N
 a 1 a HONc P H 1 nHT a
 e u N pect o g n ubh p
 3 1 1 2111 1 1 1 211 2
 // //

2761 GGTACAGCTTGTCTGTAAGCGGATGCCGGCAGCAGACAAGCCCGTCAGGGCCGCTCAGC
 ----- 2820
 CCAGTGTGGAACAGACATTGCCCTACGGCCCTCTGCTGTTGGG-CAGTCCCGCCAGTCC

GlyHisSerLeuSerValSerGlyCysArgGluGlnThrSerProSerGlyArgValSer -

H P
 1 H2 N TBM M A
 n zu 1 tba a C
 P a4 a hve e C
 1 1H 3 113 2 1

2821 GGGTGTGGCGGGTGTGCGGGCGCAGCCATGACCCAGTCACGTAGCCATAGCCGAGTGTA
 ----- 2880
 CCCACAACCGCCACAGCCCGCGTCTGGTACTGGGTGAGTGCATCGCTATCGCCTCACAT

GlyCysTrpArgValSerGlyArgSerHisAspProValThrZndArgZndArgSerVal -

34/93

FIG. 11 (cont'd)

	P	S	R D	A	BH
	n	2		p	sgNN
	u	a	s d	a	plsd
	4	N	a e	L	lApe
	H	1	1 1	1	2121

TACTGGCTTAACTATGCGGCATCAGAGCAGATTGTACTGAGAGTGCACCATATGCGGTGT
 2881 ----- 2940
 ATGACCGAATTGATACGCCGTAGTCTCGTCTAACATGACTCTCAGCTGGTATACGCCACA
 TTTTPlouAsnTyrAlaAlaSerGluGlnIleValLeuArgValHisHisMetArgCys -

S					
2					
a					
N					
1					

GAAATACCCACAGATCGGTAAGGAGAAAATACCGCATCAGCGCGTCTTCCGCTTCCTCG
 2941 ----- 3000
 CTTTATGCGGTGTCTACCGATTCTCTTTTATGCGGTAGTCCCGCAGAAGCCGAAGGAGC
 GluIleProHisArgCysValArgArgLysTyrArgIleArgArgSerSerAlaSerSer -

CTCAGTCACTCGCTCGGCTCGGTCTTCCGCTCGCGCGAGCGGTATCAGCTCACTCAAG
 3001 ----- 3060
 CAGTCACTCAAGCGACCGAGCCAGCAAGCCGACCGCGCTCGCCATAGTTCAGTCACTTC
 LeuThrAspSerLeuArgSerValValArgLeuArgArgAlaValSerAlaHisSerLys -

CCGGTAATACGGTTATCCACAGAATCAGCGGATAACCGCAGAAAGAACATGTGAGCAAAA
 3061 ----- 3120
 CGCCATTATGCCAATAGGTGTCTTAGTCCCTATTGCGTCTCTCTGTACACTCGTCTT
 AlaValIleArgLeuSerThrGluSerGlyAspAsnAlaGlyLysAsnMetEndAlaLys -

GCGCAGCAAAAGCCAGGAACCGTAAAAAGGCGCGCTTCTGCGGTCTTCCATAGCGTC
 3121 ----- 3180
 CCGGTCTGCTTCCGCTCTCTGCACTTCTCGGCGCAAGCAGCCGCAAAAGGTATCCGAG
 GlyGlnGlnLysAlaArgAsnArgLysLysAlaAlaLeuLeuAlaPhePheHisArgLeu -

CCCCCCGCTGACGAGCATCACAATAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACA
 3181 ----- 3240
 CCGGGGGGAGCTGCTCGTAGTCTTTTACGCTCGAGTTCAGTCTCCACCGCTTTGCGGTGT
 ArgProProAspGluHisHisLysArgArgArgSerSerGlnArgTTPArgAsnProThr -

35/93

FIG. 11'(cont'd)

BES	BES	A	H
SCC	SCC	1	1 HM
cor	cor	1	n hn
NRP	NRP	u	P al
121	121	1	1 11

3241 GGACTATAAAGATACAGCGGTTTCCCCCTCGGAAGCTCCCTCGTCCGCTCTCTGTTCCG 3300
 CCTCATATTTCTATGCTCCGCAAGGGGGACCTTCGAGCGAGCACCGGAGAGGACAAGGC
 GlyLeuEndArgTyrGlnAlaPheProProGlySerSerLeuValArgSerProValPro -

F	HM	H
n	ps	i HH
u	ap	n hn
4	21	P ae
H		1 12

3301 ACCCTGCGCGCTTACCGGATACCTGTCCCGCTTCTCCCTTCGGGAAGCGTGGCGCTTCT 3360
 TCGGACGGCGAATGGCTATGACACGGCGGAAGAGCGGAAGCCCTTCGACCGCGAAAGA
 ThrLeuProLeuThrGlyTyrLeuSerAlaPheLeuProSerGlySerValAlaLeuSer -

D	A
d	1
e	u
1	1

3361 CAATGCTCAGCGTGTAGGTATCTCAGTTCGGTGTAGGTCTGCTTCCCTCCAAGCTGGGCTGT 3420
 GTTACGAGTGGGACATCCATAGAGTCAAGCCACATCCAGCAAGCGAGGTTTCGACCCGACA
 GlnCysSerArgCysArgTyrLeuSerSerValEndValValArgSerLysLeuGlyCys -

A	BH	B	NF H	HM	H
P	sgN	b	sn i H	psa	i
a	pis	v	pu n h	spe	n
L	lap	1	B4 P a	213	e
1	212		2H 1 1		1

3421 GTGCACGAACCCCCCGTTCAGCCCCGACCGCTCGCCCTTATCCGGTAACTATCGTCTTGAG 3480
 CACGTGCTTGGGGGGCAAGTCCGGCTCGCGACCGCGAATAGGCCATTGATAGCAGAACTC
 ValHisGluProProValGlnProAspArgCysAlaLeuSerGlyAsnTyrArgLeuGlu -

S	F	F	BM B	H
HMNC	n	n	ba b	n
pscr	u	u	ve v	1
apiP	4	4	13 1	1
2111	H	H		

3481 TCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAAACAGGATTAGC 3540
 AGGTTGGGCCATTCTGTGCTGAATAGCGGTGACCGTCCGTCCGTGACCAATTGCTCTAATG
 SerAsnProValArgHisAspLeuSerProLeuAlaAlaAlaThrGlyAsnArgIleSer -

H
a
e
3

3541 AGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCCTAACTACGGCTAC 3600
 TCTCGCTCCATACATCCGCCACGATGTCTCAAGAACTTCACCACCGGATTGATGCCGATG
 ArgAlaArgTyrValGlyGlyAlaThrGluPheLeuLysTyrTyrProAsnTyrGlyTyr -

3791

89085519

36/93

FIG. 11(cont'd)

M
a
●
1
 H
1 H
a h
P a
1 1
 M
a
●
3
 ACTAGAAGGACAGTATTTGGTATCTCCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGA
 1601 ----- 1660
 TGATCTTCCTGTCATAAACCATAGACGGCAGACGACTTCGGTCAATGGAAGCCTTTCT
 ThrArgArgThrValPheGlyIleCysAlaLeuLeuLysProValThrPheGlyLysArg -

S
A MNDaX
1 bdpups
u oen3a
1 121A21
 H
a
p
B
2
 GTTCGTAGCTCTTGATCCGGCAAAACAACCCACCGCTGGTAGCGGTGGT-----GTTTGC
 1661 ----- 1720
 CAACCATCGAGAACTAGCCCGTTTCTTTTGGTGGCGACCATCGCCACCAAAAAACAACG
 ValGlySerSerEndSerGlyLysGlnThrThrAlaGlySerGlyGlyPhePheValCys -

P F H S S
a BnHIT MNDaX MNDaX MNDaX
u buhnh bdpuh bdpuh bdpba
4 vDaPa oen3o oen3o oen3o
H 12111 121A2 121A2 1212A
 1721 ----- 1780
 AACCCAGCAGATTACCCCGAGAAAAAAGGATCTCAAGAGATCCTTTGATCTTTCTACG
 TTCGTCTCTAATCCCGCTCTTTTCTCTAGAGTTCTTCTAGGAACTAGAAAAGATGC
 LysGlnGlnIleThrArgArgLysLysGlySerGlnGluAspProLeuIlePheSerThr -

D H M N H
d g a 1 b
● a ● a o
1 1 2 3 2
 1781 ----- 1840
 GGGTCTGACGCTCAGTCCGAACGAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCA
 CCCAGACTCGGAGTCACCTTGCTTTTGAGTGCAATTCCCTAAAACCACTACTCTAATAGT
 GlySerAspAlaGlnTrpAsnGluAsnSerArgEndGlyIleLeuValMetArgLeuSer -

S S S S A
H MNDaX H MNDaX BMDNaX MNDaX
p bdpuh a bdpuh gbpduh bdpu 1
h oen3o e oen3o lone3o oen3 u
1 121A2 1 121A2 2112A2 121A 1
 1841 ----- 1900
 AAAAGGATCTTCACCTAGATCCTTTTCAGATCTCCCGATCTTTAGCTGTCTTGGTTTCCG
 TTTTCTAGAAAGTGGATCTAGGAAAAGTCTAGAGGGCTAGAAATCGACAGAACCAACGG
 LysArgIlePheThrEndIleLeuPheArgSerProAspLeuEndLeuSerTrpPheAla -

H
1 H
a h
P a
1 1
 D H R
d ns
● la
1 11
 1901 ----- 1960
 CAAAGCGCATTCATATAATCTTTACGGGTTATGCGTTGTTCCATACAACCTCCTTAGTACA
 GTTTCGGGTAACGTATTAGAAAAGTCCCAATACCGCAACAAAGGTATGTTGGAGGAATCATGT
 3792 GlnSerAlaLeuHisAsnLeuSerGlyLeuCysValValProTyrAsnLeuLeuSerThr -

89085519

37/93

FIG. 11(cont'd)

N			H
NsH	M		1
lpp	n		n
aHh	1		c
311	1		2

3961 TCCAACCATTATCACCGCCAGAGGTAAAATAGTCAACACGCACGGTGTAGATAATTTATC 4020

 ACGTTGGTAATAGTGGCGGTCTCCATTTTATCAGTTGTCCGTGCCACAATCTATAAATAG
 CysAsnHisTyrHisArgGlnArgEndAsnSerGlnHisAlaArgCysEndIlePheIle -

	EM	BH		P
	pa	sqn		n A
	he	pis		u l
	12	lap		1 4 u
		212		1 H 1

4021 CCTTCCGGTGTATAGATTTAACGTATGAGCACAAAAAGAACCATTAACACAGACCAGC 4080

 GGAACGCCACTATCTAAATTGCATCTCGTCTTTTCTTTGGTAAATGTGTTCTCGTCC
 ProCysGlyAspArgPheAsnValEndAlaGlnLysArgAsnHisEndHisLysSerSer -

	B	M	H
	b	a	g
	v	e	a
	1	2	1

4081 TTGAGGACCCACGTCGCCCTTAAGCAATTTATCAAAAAAGAAAAATCAACTTGGCTTAT 4140

 AACTCTCGGTGCAGCGGAATTCCTAAATACCTTTCTTTTACTTGAACCGAATA
 LeuArgThrHisValAlaLeuLysGlnPheMetLysLysArgLysMetAsnLeuAlaTyr -

BES	H	
scc	1	P
cor	n	o
NRP	2	k
121	1	1

4141 CCCAGGAATCTGTCCAGACAAGATGGCGATGGCGCAGTCAGCCGTGGTGTCTTTATTTA 4200

 GCGTCTTTAGACACCGCTCTGTCTACCCCTACCCCGTCAGTCCCAACCAAGAAATAAT
 ProArgAsnLeuSerGlnThrArgTrpGlyTrpGlySerGlnAlaLeuValLeuTyrLeu -

	S	P
	2N	n
	as	u
	N1	4
	11	H

4201 ATGGCATCAATGCATTAAATGCTTATAACGGCGCATTCCTTACAAAAATCTCAAAGTTA 4260

 TACCGTAGTTACGTAAATTACGAATATTCGGCGCTAACGAATGTTTTTAAGAGTTTCAAT
 MetAlaSerMetHisEndMetLeuIleThrProHisCysLeuGlnLysPheSerLysLeu -

	M
	b
	o
	2

4261 CGGTTGAAGAATTTAGCCCTTCAATGCCAGAGAAATCTACGAGATGTATGAAGCGGTTA 4320

 CGCAAGTTCTTAATCGGGAAGTTAGCGGTCTCTTTAGATGCTCTACATAGTTCCGCAAT
 AlaLeuLysAsnLeuAlaLeuGlnSerProGluLysSerThrArgCysMetLysArgLeu -

3793

89085519

38/93

FIG. 11(cont'd)

P M D B R N
 n a d b s l
 u e e v a a
 4 3 1 1 1 3
 H 3 1 1 1 3
 4321 GTATGCAGCCGTCACCTTAGAAGTGAGTATGAGTACCCTGTTTTTCTCATGTTTCAGGCAG 4380

 CATACGTCCGCAGTGAATCTTCACTCATACTCATGGGACAAAAAGAGTACAAGTCCGTC
 ValCysSerArgHisLeuGluValSerMetSerThrLeuPhePheLeuMetPheArgGln -

H H S H
 DP1 AD f a P H
 don ld a N 1
 ekd ue 1 1 1
 113 11 1 1 1
 4381 GGATGTTCTCACCTAAGCTTAGAACCTTACCAAGGTGATCCGGACAGATCGGTAAGCA 4440

 CCTACAAGAGTGGATTGGAATCTTGGAAATGGTTTCCACTACCGCTCTCTACCCATTCTGT
 GlyCysSerHisLeuSerLeuGluProLeuProLysValMetArgArgAspGlyEndAla -

H BM N
 i en l
 n ml a
 f 11 3
 1 11 3
 4441 CAACCAAAAAAGCCAGTCAATTCCTGCTTACGTTCAAGGTAAATTCATGACCG 4500

 GTTCGTTTTTTCGGTCACTAAGAGCTAAGACCG ACTCCAATTCCATTAAAGGTACTGGC
 GlnProLysLysProValIleLeuHisSerGlyLeuArgLeuLysValIleProEndPro -

H H D
 i A i d
 n l n e
 d u c e
 3 1 2 1
 4501 CACCACAGGCTCCAAGCCAAGCTTTCCTGACGGAAATGTTAATTCCTCGTTGACCCCTGAGC 4560

 GTGGTTCTCCGAGGTTCCGTTCCGAAGGACTGCCCTTACAATTAAGAGCAACTGGGACTCG
 HisGlnGlnAlaProSerGlnAlaPheLeuThrGluCysEndPheSerLeuThrLeuSer -

Scc H H
 cor p p
 NRP h h
 121 1 1
 4561 AGGCTGTTGAGCCAGGTGATTTCTGCATAGCCAGACTTGGGGGTGATGAGTTTACCTTCA 4620

 TCCGACAACTCGGTCCACTAAAGACGTATCGGTCTGAACCCCCACTACTCAAATGGAAAT
 ArgLeuLeuSerGlnValIleSerAlaEndProAspLeuGlyValMetSerLeuProSer -

R
 s
 a
 1
 4621 AGAAACTAATTAGGGATAGCGGTCAAGTGTTTTTACAACCACTAAACCCACAGTACCCAA 4680

 TCTTTGATTAAATCCCTATCGCCAGTCCACAAAAATGTTGGTGATTTCGGGTGTCATGGGT
 ArgAsnEndLeuGlyIleAlaValArgCysPheTyrAsnHisEndThrHisSerThrGln -

89085519

39/93

FIG. 11(cont'd)

S
 MNDA 1
 bdpu 1
 oen3 1
 121A 1
 / /
 4681 TGATCCCATGCAATGAGAGTTGTTCCGTTGTGGGGAAGTTATCGCTAGTCAGTGGCCTG 4740
 ACTAGCGGTACGTTACTCTCAACAAGGCAACACCCCTTCAATAGCGATCAGTCACCGGAC
 EndSerHisAlaMetArgValValProLeuTyrGlyLysLeuSerLeuValSerGlyLeu -
 S
 M M MNDA
 a b bdpu
 e o oen3
 2 2 121A
 / /
 4741 AAGAGACGTTTCGCTGATCCGCCAAGGTTTCTGGTGGGCGCATAGCTGATAACAAATTGAG 4800
 TTCTCTGCAACCGGACTAGCCGTTCCACAAGACCAGCCCGGTATCGACTATTGTTAACTC
 LysArgArgLeuAlaAspArgGlnGlyValLeuValGlyAlaEndLeuIleThrIleGlu -
 S
 H PS P MNDA
 1B af bHP
 nb ua ugs
 fv 4M 4at
 11 H1 H11
 / /
 4801 CAAGAATCTTCATCGGGGCTGCAGCCCAACCATCGCTCCGGCGGTAGAGGATCTCTACCTA 4860
 GTTCTTAGAAGTAGCCCCGACGTCGGGTCTACCCAGCCCGCATCTCTAGAGAGTGGAT
 GlnGluSerSerSerGlyLeuGlnProThrMetArgProAlaEndArgIleSerHisLeu -
 S
 MNDA
 pbdpu
 hoen3
 1121A2
 / /
 4861 CCAACAAATGCCCCCTGCAAAAAATAAATTCATATAAAAAACATACAGATAACCATCTG 4920
 GGTTCGTTACGGGGGGACGTTTATTATTAAGTATATTTTGTATGCTATTCGTAGAC
 ProAsnAsnAlaProLeuGlnLysIleAsnSerTyrLysLysHisThrAspAsnHisLeu -
 S
 H H BH
 p h Hsg
 b c ppl
 1 2 1 121
 /
 4921 CGGTGATAAATTATCTCTGGCGGTGTTGACATAAATACCACTGGCGGTGATACTGAGCAC 4980
 GCCACTATTTAATAGAGACCGCCACAACCTGTATTTATGGTGACCGCCACTATGACTCGTG
 ArgEndEndIleIleSerGlyGlyValAspIleAsnThrThrGlyGlyAspThrGluHis -
 S
 N H N M HB
 s g 1 a GP
 p a 1 a
 2 1 3 3 11
 /
 3795 4981 ATCAGCAGGACCGCACTGACCACCATGAAGGTGACGCTCTTAAAAATTAAGCCCTGAAGAAG 5040
 TAGTCGTCTCTCGGTGACTGGTGGTACTTCCACTCGGAGAAATTTTAAATCGGGACTTCTTC

89085519

FIG. 11(cont'd)

7
 n BM B
 u sb b
 4 mo v
 H 12 l
 5041 GGCACCATTCAAAGCAGAAGGCTTTGGGGTGTGTGATACGAAACGAACCAATT 5092
CCGTCGTAACTTTCGTCTTCCGAAACCCACACACTATGCTTTGCTTCGTAA

GlySerIleGlnSerArgArgLeuTrpGlyValIlendTyrGluThrLysHis???

Enzymes that do cut:


AccI	Aha2	Afl2	AluI	ApaI	AvaI	Ava2	BanI	Ban2	BbeI	BbvI
BclI	BglI	Bgl2	BemI	SepI	BspMI	BstE2	BstNI	BstXI	CfrI	ClaI
DdeI	DpnI	Dra2	EagI	EcoB	EcoK	EcoRI	EcoR2	EcoRV	FnuD2	Fnu4H
PokI	PspI	Bae2	Bae3	HgaI	HgiA1	HgiD1	HhaI	Hinc2	Hind3	HinfI
HinPI	Hpa2	HphI	MaeI	Mae2	Mae3	MboI	Mbo2	MnlI	HspI	HstI
Hst2	MaeI	NarI	NciI	NdeI	Nde2	NheI	Nla3	Nla4	NruI	NotI
Hsp2	HspB2	HspH1	PflMI	PpuMI	PseI	PstI	Pvu2	RsaI	SacI	Sall
Sau3A	Sau96	ScrPI	SfaNI	SinI	SetI	StuI	StyI	TaqI	ThaI	TclI
Xho2	Xma3									

Enzymes that do not cut:

Aat2	Aha3	Apal	Asp70	Asp71	Asu2	Avr2	BalI	BamHI	BspM2	BssH2
DraI	Dra3	EspI	HpaI	KpnI	MluI	NcoI	NotI	PaeR7	PvuI	Rsr2
Sac2	Scal	Sfil	SmaI	SnaBI	SpeI	SphI	SepI	Set2	XbaI	XhoI
XmaI	XmnI	Xor2								

4/1/93

FIG. 12



lane 1 = Pre-induced

lane 4 = uninduced

lane 5 = 3 hr. post-induction

lane 11 = overnight post-induction (~16 hr)


MW = Molecular wt. markers

89085519

3797


SUBSTITUTE SHEET

FIG. 13



43/93

FIG. 14A




3799

89085519

44/93

FIG. 14B




89085519

3800

SUBSTITUTE SHEET

45/93

FIG. 14C



46'93

pBG391
 :BG368 backbone
 :soluble T4#3
 :AA #3 = LYS

FIG. 15

bg381.seq Length: 6151

```

1  GAATTAATTC CAGCTTGCTG TGAATGTGT GTCAGTTAGG GTGTGGAAAG
51  TCCCCAGGCT CCCCAGCAGG CAGAAGTATG CAAAGCATGC ATCTCAATTA
101 GTCAGCAACC AGGTGTGGAA AGTCCCCAGG CTCCCCAGCA GGCAGAAGTA
151 TGCAAAGCAT GCATCTCAAT TAGTCAGCAA CCATAGTCCC GCCCCTAACT
201 CCGCCCATCC CGCCCTAAC TCCGCCAGT TCCGCCATT CTCCGCCCA
251 TGGCTGACTA ATTTTTTTTA TTTATGCAGA GGCCGAGGCC GCCTCGGCCT
301 CTGAGCTATT CCAGAAGTAG TGAGGAGGCT TTTTGGAGG GGTCTCTCTC
351 GTATAGAAAC TCGGACCACT CTGAGACGAA GGCTCGCGTC CAGGCCAGCA
401 CGAAGGAGGC TAAGTGGGAG GGGTAGCGGT CGTTGTCCAC TAGGGGGTCC
451 ACTCGCTCCA GGGTGTGAAG ACACATGTCG CCCTCTTCGG CATCAAGGAA
501 GGTGATTGGT TTATAGGTGT AGGCCACGTG ACCGGGTGTT CCTGAAGGGG
551 GGCTATAAAA GGGGGTGGGG GCGCGTTCGT CCTCACTCTC TTCCGCATCG
601 CTGTCTGCGA GGGCCAGCTG TTGGGCTCGC GGTGAGGAC AAACCTCTCG
651 CGGTCTTTCC AGTACTCTTG GATCGGAAAC CCGTCGGCCT CCGAACGGTA
701 CTCCGCCACC GAGGGACCTG AGCGAGTCCG CATCGACCGG ATCGGAAAAC
751 CTCTCGAGAA AGGCGTCTAA CCAGTCACAG TCGCAAGGTA GGCTGAGCAC
801 CGTGGCGGGC GGCAGCGGGT GCGGTTCGGG GTTGTCTCTG GCGGAGGTGC
851 TGCTGATGAT GTAATTAAAG TAGGCGGTCT TGAGACGGCG GATGGTCGAG
901 GTGAGGTGTG GCAGGCTTGA GATCGATCTG GCCATACACT TGAGTGACAA
951 TGACATECAC TTTGCCTTTC TCTCCACAGG TGTCCACTCC CAGGTCCAAC
1001 TGGATCCAAG CTTCGACTCG AGGAATTCCC CGAAGGAACA AAGCACCCCTC
1051 CCCACTGGGC TCCTGGTTGC AGAGCTCCAA GTCCTCACAC AGATACGCCT
1101 GTTTGAGAAG CAGCGGGCAA GAAAGACGCA AGCCCAGAGG CCCTGCCATT
1151 TCTGTGGGCT CAGGTCCCTA CTGGCTCAGG CCCCTGCCTC CCTCGGCAAG
1201 GCCACAATGA ACCGGGGAGT CCCTTTTAGG CACTTGCTTC TGGTGCTGCA
1251 ACTGGCGCTC CTCCCAGCAG CCACTCAGGG AAAGAAAGTG GTGCTGGGCA
1301 AAAAAGGGGA TACAGTGGAA CTGACCTGTA CAGCTTCCCA GAAGAAGAGC
1351 ATACAATTCC ACTGGAAAAA CTCCAACCAG ATAAAGATTC TGGGAAATCA
1401 GGGCTCCTTC TTAACCTAAG GTCCATCCAA GCTGAATGAT CGCGCTGACT

```

3802

891085519

47/93

FIG. 15 (con'd)

1451 CAAGAAGAAG CTTGTGGGAC CAAGGAAACT TTCCCTGAT CATCAAGAAT
 1501 CTTAAGATAG AAGACTCAGA TACTTACATC TGTGAAGTGG AGGACCAGAA
 1551 GGAGGAGGTG CAATTGCTAG TGTTCGGATT GACTGCCAAC TCTGACACCC
 1601 ACCTGCTTCA GGGGCAGAGC CTGACCCTGA CCTTGGAGAG CCCCCCTGGT
 1651 AGTAGCCCCCT CAGTGCAATG TAGGAGTCCA AGGGGTAAAA ACATACAGGG
 1701 GGGGAAGACC CTCTCCGTGT CTCAGCTGGA GCTCCAGGAT AGTGGCACCT
 1751 GGACATGCAC TGTCTTGACAG AACCAGAAGA AGGTGGAGTT CAAAATAGAC
 1801 ATCGTGGTGC TAGCTTTCCA GAAGGCCTCC AGCATAGTCT ATAAGAAAGA
 1851 GGGGGAACAG GTGGAGTTCT CCTTCCCACT CGCCTTTACA GTTGAAAAGC
 1901 TGACGGGCAG TGGCGAGCTG TGGTGGCAGG CGGAGAGGGC TTCCTCCTCC
 1951 AAGTCTTGGA TCACCTTTGA CCTGAAGAAC AAGGAAGTGT CTGTAAAACG
 2001 GGTTACCCAG GACCCTAAGC TCCAGATGGG CAAGAAGCTC CCGCTCCACC
 2051 TCACCCTGCC CCAGGCCTTG CCTCAGTATG CTGGCTCTGG AAACCTCACC
 2101 CTGGCCCTTG AAGCGAAAAC AGGAAAGTTG CATCAGGAAG TGAACCTGGT
 2151 GGTGATGAGA GCCACTCAGC TCCAGAAAAA TTTGACCTGT GAGGTGTGGG
 2201 GACCCACCTC CCCTAAGCTG ATGCTGAGTT TGAAACTGGA GAACAAGGAG
 2251 GCAAAGGTCT CGAAGCGGGA GAAGGCGGTG TGGGTGCTGA ACCCTGAGGC
 2301 GGGGATGTGG CAGTGTCTGC TGAGTGA CTC GGGACAGGTC CTGCTGGAAT
 2351 CCAACATCAA GGTTCTGCCC ACATGGT CGA CCCCAGTGCA GCCAATGGCC
 2401 CTGATTTGAG ATCTTTGTGA AGGAACCTTA CTTCTGTGGT GTGACATAAT
 2451 TGGACAAACT ACCTACAGAG ATTTAAAGCT CTAAGGTAAA TATAAAATTT
 2501 TTAAGTGTAT AATGTGTAA ACTACTGATT CTAATTGTTT GTGTATTTTA
 2551 GATTCCAACC TATGGAAGT ATGAATGGGA GCAGTGGTGG AATGCCTTTA
 2601 ATGAGGAAAA CCTGTTTTGC TCAGAAGAAA TGCCATCTAG TGATGATGAG
 2651 GCTACTGCTG ACTCTCAACA TTCTACTCCT CCAAAAAAGA AGAGAAAGGT
 2701 AGAAGACCCC AAGGACTTTC CTTCAGAATT GCTAAGTTTT TTGAGTCATG
 2751 CTGTGTTTAG TAATAGA ACT CTTGCTTGCT TTGCTATTTA CACCACAAAG
 2801 GAAAAAGCTG CACTGCTATA CAAGAAAATT ATGGAAAAAT ATTCTGTAAC
 2851 CTTTATAAGT AGGCATAACA GTTATAATCA TAACATACTG TTTTTTCTTA
 2901 CTCCACACAG GCATAGAGTG TCTGCTATTA ATA ACTATGC TCAAAAATTG
 2951 TGTACCTTTA GCTTTTTAAT TTGTAAAGGG GTTAATAAGG AATATTTGAT
 3001 GTATAGTGCC TTGACTAGAG ATCATAATCA GCCATACCAC ATTTGTAGAG
 3051 GTTTTACTTG CTTTAACTGACAC CTCCCTCTGA ACCTGAAACA

48/93

F1G.15(cont'd)

3101 TAAATGAAT GCAATTGTTG TTGTTAALIT GTTTATTGCA GCTTATAATG
 3151 GTTACAAATA AAGCAATAGC ATCACAAATT TCACAAATAA AGCATTTTTT
 3201 TCACTGCATT CTAGTTGTGG TTTGTCCAAA CTCATCAATG TATCTTATCA
 3251 TGTCTGGATC CTCTACGCCG GACGCATCGT GGCCGGGCATC ACCGGCGCCA
 3301 CAGGTGCGGT TGCTGGCGCC TATATCGCCG ACATCACCGA TGGGGAAGAT
 3351 CGGGCTCGCC ACTTCGGGCT CATGAGCGCT TGTTTCGGCG TGGGTATGGT
 3401 GGCAGGCCCC TGGCCGGGGG ACTGTTGGGC GCCATCTCCT TGCATGCACC
 3451 ATTCTTGCG GCGGCGGTGC TCAACGGCCT CAACCTACTA CTGGGCTGCT
 3501 TCCTAATGCA GGAGTCGCAT AAGGGAGAGC GTCGACCGAT GCCCTTGAGA
 3551 GCCTTCAACC CAGTCAGCTC CTTCCGGTGG GCGCGGGGCA TGA CTATCGT
 3601 CGCCGCACTT ATGACTGTCT TCTTTATCAT GCAACTCGTA GGACAGGTGC
 3651 CGGCAGCGCT CTGGGTCATT TTCGGCGAGG ACCGCTTTCG CTGGAGCGCG
 3701 ACGATGATCG GCCTGTCGCT TGCGGTATTC GGAATCTTGC ACGCCCTCGC
 3751 TCAAGCCTTC GTCACTGGTC CCGCCACCAA ACGTTTCGGC GAGAAGCAGG
 3801 CCATTATCGC CGGCATGGCG GCCGACGCGC TGGGCTACGT CTTGCTGGCG
 3851 TTCGCGACGC GAGGCTGGAT GGCCTTCCCC ATTATGATTC TTCTCGCTTC
 3901 CGGCGGCATC GGGATGCCCG CGTTGCAGGC CATGCTGTCC AGGCAGGTAG
 3951 ATGACGACCA TCAGGGACAG CTTCAAGGAT CGCTCGCGGC TCTTACCAGC
 4001 CTA ACTTCGA TCACTGGACC GCTGATCGTC ACGGCGATTT ATGCCGCCTC
 4051 GGCGAGCACA TGGAACGGGT TGGCATGGAT TG TAGGCGCC GCCCTATACC
 4101 TTGTCTGCCT CCCC GCGTTG CGTCGCGGTG CATGGAGCCG GGCCACCTCG
 4151 ACCTGAATGG AAGCCGGCGG CACCTCGCTA ACGGATTCAC CACTCCAAGA
 4201 ATTGGAGCCA ATCAATTCTT GCGGAGAACT GTGAATGCGC AAACCAACCC
 4251 TTGGCAGAAC ATATCCATCG CGTCCGCCAT CTCCAGCAGC CGCAGCGGGC
 4301 GCA-TCTCGG CCGCGTTGCT GCGGTTTTTC CATAGGCTCC GCCCCCTGA
 4351 CGAGCATCAC AAAAATCGAC GCTCAAGTCA GAGGTGGCGA AACCCGACAG
 4401 GACTATAAAG ATACCAGGCG TTTCCCCCTG GAAGCTCCCT CGTGCGCTCT
 4451 CCTGTTCCGA CCCTGCCGCT TACCGGATAC CTGTCCGCCT TTCTCCCTTC
 4501 GGGAAAGCGTG GCGCTTTCTC AATGCTCACG CTGTAGGTAT CTCAGTTCGG
 4551 TG TAGGTCGT TCGCTCCAAG CTGGGCTGTG TGCACGAACC CCCC GTTCAG
 4601 CCCGACCGCT GCGCCTTATC CGGTA ACTAT CGTCTTGAGT CCAACCCGGT
 4651 AAGACACGAC TTATCGCCAC TGGCAGCAGC CACTGGTAAC AGGATTAGCA
 470: GAGCGAGGTA TG TAGGCGAT CCTACAGAGT TCTTGAAGTG GTGGCCTAAC

89085574

49/93

FIG. 15(cont'd)

4751 TACGGCTACA CTAGAAGGAC AGTATTTGGT ATCTGCGCTC TGCTGAAGCC
4801 AGTTACCTTC GGAAAAAGAG TTGGTAGCTC TTGATCCGGC AAACAAACCA
4851 CCGCTGGTAG CCGTGGTTTT TTTGTTTGCA AGCAGCAGAT TACGCGCAGA
4901 AAAAAAGGAT CTCAAGAAGA TCCTTTGATC TTTTCTACGG GGTCTGACGC
4951 TCAGTGGAAC GAAAACTCAC GTTAAGGGAT TTTGGTCATG AGATTATCAA
5001 AAAGGATCTT CACCTAGATC CTTTTAAATT AAAAATGAAG TTTTAAATCA
5051 ATCTAAAGTA TATATGAGTA AACTTGGTCT GACAGTTACC AATGCTTAAT
5101 CAGTGAGGCA CCTATCTCAG CGATCTGTCT ATTTCGTTCA TCCATAGTTG
5151 CCTGACTCCC CGTCGTGTAG ATAACTACGA TACGGGAGGG CTTACCATCT
5201 GGCCCCAGTG CTGCAATGAT ACCGCGAGAC CCACGCTCAC CGGCTCCAGA
5251 TTTATCAGCA ATAAACCAGC CAGCCGGAAG GGCCGAGCGC AGAAGTGGTC
5301 CTGCAACTTT ATCCGCCTCC ATCCAGTCTA TTAATTGTTG CCGGGAAGCT
5351 AGAGTAAGTA GTTCGCCAGT TAATAGTTTG CGCAACGTTG TTGCCATTGC
5401 TGCAGGCATC GTGGTGTAC GCTCGTCGTT TGGTATGGCT TCATTCAGCT
5451 CCGGTTCCCA ACGATCAAGG CGAGTTACAT GATCCCCCAT GTTGTGCAAA
5501 AAAGCGGTTA GCTCCTTCGG TCCTCCGATC GTTGTGAGAA GTAAGTTGGC
5551 CGCAGTGTTA TCACTCATGG TTATGGCAGC ACTGCATAAT TCTCTTACTG
5601 TCATGCCATC CGTAAGATGC TTTTCTGTGA CTGGTGAGTA CTCAACCAAG
5651 TCATTCTGAG AATAGTGTAT GCGGCGACCG AGTTGCTCTT GCCCGGCGTC
5701 AACACGGGAT AATACCGCGC CACATAGCAG AACTTTAAAA GTGCTCATCA
5751 TTGGAAAACG TTCTTCGGGG CGAAAACTCT CAAGGATCTT ACCGCTGTTG
5801 AGATCCAGTT CGATGTAACC CACTCGTGCA CCCAACTGAT CTTCAGCATC
5851 TTTTACTTTC ACCAGCGTTT CTGGGTGAGC AAAAACAGGA AGGCAAAATG
5901 CCGCAAAAAA GGAATAAGG GCGACACGGA AATGTTGAAT ACTCATACTC
5951 TTCCTTTTTT AATATTATTG AAGCATTTAT CAGGGTTATT GTCTCATGAG
6001 CGGATACATA TTTGAATGTA TTTAGAAAAA TAAACAAATA GGGGTTCCGC
6051 GCACATTTCC CCGAAAAGTG CCACCTGACG TCTAAGAAAC CATTATTATC
6101 ATGACATTAA CCTATAAAAA TAGGCGTATC ACGAGGCCCT TTCGTCTTCA

6151 A

89085519

3805

SUBSTITUTE SHEET

PB6392
 :BG368 backb n
 :soluble T4#7
 :AA #3 = LVS
 :182AA+6AA
 :from 203-5

FIG. 16

bg392.seq Length: 6149

1 GAATTAATTC CAGCTTGCTG TGAATGTGT GTCAGTTAGG GTGTGGAAG
 51 TCCCCAGGCT CCCCAGCAGG CAGAAGTATG CAAAGCATGC ATCTCAATTA
 101 GTCAGCAACC AGGTGTGGAA AGTCCCCAGG CTCCCCAGCA GGCAGAAGTA
 151 TGCAAAGCAT GCATCTCAAT TAGTCAGCAA CCATAGTCCC GCCCCTAACT
 201 CCGCCCATCC CGCCCCCTAAC TCCGCCCAGT TCCGCCCATT CTCCGCCCCA
 251 TGGCTGACTA ATTTTTTTTA TTTATGCAGA GGCCGAGGCC GCCTCGGCCT
 301 CTGAGCTATT CCAGAAGTAG TGAGGAGGCT TTTTGGAGG GGTCTCTCTC
 351 GTATAGAAAC TCGGACCACT CTGAGACGAA GGCTCGCGTC CAGGCCAGCA
 401 CGAAGGAGGC TAAGTGGGAG GGGTAGCGGT CGTTGTCCAC TAGGGGGTCC
 451 ACTCGCTCCA GGGTGTGAAG ACACATGTCT CCCTCTTCGG CATCAAGGAA
 501 GGTGATTGGT TTATAGGTGT AGGCCACGTG ACCGGGTGTT CCTGAAGGGG
 551 GGCTATAAAA GGGGGTGGGG GCGCGTTCGT CCTCACTCTC TTCCGCATCG
 601 CTGTCTGCGA GGGCCAGCTG TTGGGCTCGC GGTTGAGGAC AAACCTCTTCG
 651 CGGTCTTTCC AGTACTCTTG GATCGGAAAC CCGTCGGCCT CCGAACGGTA
 701 CTCCGCCACC GAGGGACCTG AGCGAGTCCG CATCGACCGG ATCGGAAAAC
 751 CTCTCGAGAA AGGCGTCTAA CCAGTCACAG TCGCAAGGTA GGCTGAGCAC
 801 CGTGGCGGGC GGCAGCGGGT GCGGGTCGGG GTTGTTTCTG GCGGAGGTGC
 851 TGCTGATGAT GTAATTAAAG TAGGCGGTCT TGAGACGGCG GATGGTCGAG
 901 GTGAGGTGTG GCAGGCTTGA GATCGATCTG GCCATACACT TGAGTGACAA
 951 TGACATCCAC TTTGCCTTTC TCTCCACAGG TGTCCACTCC CAGGTCCAAC
 1001 TGGATCCAAG CTTCGACTCG AGGAATTCCC CGAAGGAACA AAGCACCTC
 1051 CCCACTGGGC TCCTGGTTGC AGAGCTCCAA GTCCTCACAC AGATACGCCT
 1101 GTTTGAGAAG CAGCGGGCAA GAAAGACGCA AGCCCAGAGG CCCTGCCATT
 1151 TCTGTGGGCT CAGGTCCCTA CTGGCTCAGG CCCCTGCCTC CCTCGGCAAG
 1201 GCCACA^{MET}ATGA ACCGGGGAGT CCCTTTTAGG CACTTGCTTC TGGTGCTGCA
 1251 ACTGGCGCTC CTCCCAGCAG CCACT^{AA-23}CAGGG AAAGAAAGTG GTGCTGGGCA
 1301 AAAAAGGGGA TACAGTGGAA^{AA1}CTGACCTTA CAGCTTCCCA GAAGAAGAGC
 1351 ATACAATTCC ACTGGAAAAA CTCCAACCAG ATAAAGATTC TGGGAAATCA

51/93

FIG. 16(cont'd)

1401 GGGCTCCTTC TTAACATAAG GTCCATCCAA GCTGAATGAT CGCGCTGACT
 1451 CAAGAAGAAG CTTGTGGGAC CAAGGAAACT TTCCCCTGAT CATCAAGAAT
 1501 CTTAAGATAG AAGACTCAGA TACTTACATC TGTGAAGTGG AGGACCAGAA
 1551 GGAGGAGGTG CAATTGCTAG TGTTCGGATT GACTGCCAAC TCTGACACCC
 1601 ACCTGCTTCA GGGGCAGAGC CTGACCCTGA CCTTGGAGAG CCCCCCTGGT
 1651 AGTAGCCCCCT CAGTGCAATG TAGGAGTCCA AGGGGTAAAA ACATACAGGG
 1701 GGGGAAGACC CTCTCCGTGT CTCAGCTGGA GCTCCAGGAT AGTGGCACCT
 1751 GGACATGCAC TGTCTTGCAG AACCAGAAGA AGGTGGAGTT CAAAATAGAC
 1801 ATCGTGGTGC TAGCTTTCCA GAACCTCCAG CATAGTCTAT ^{STOP} AAGAAAGAGG
 1851 GGGAACAGGT GGAGTTCTCC TTCCCCTCG CCTTTACAGT TGAAAAGCTG
 1901 ACGGGCAGTG GCGAGCTGTG GTGGCAGGCG GAGAGGGCTT CCTCCTCCAA
 1951 GTCTTGGATC ACCTTTGACC TGAAGAACAA GGAAGTGTCT GTAAAACGGG
 2001 TTACCCAGGA CCCTAAGCTC CAGATGGGCA AGAAGCTCCC GCTCCACCTC
 2051 ACCCTGCCCC AGGCCTTGCC TCAGTATGCT GGCTCTGGAA ACCTCACCTC
 2101 GGCCCTTGAA GCGAAAACAG GAAAGTTGCA TCAGGAAGTG AACCTGGTGG
 2151 TGATGAGAGC CACTCAGCTC CAGAAAAATT TGACCTGTGA GGTGTGGGGA
 2201 CCCACCTCCC CTAAGCTGAT GCTGAGTTTG AACTGGAGA ACAAGGAGGC
 2251 AAAGGTCTCG AAGCGGGAGA AGGCGGTGTG GGTGCTGAAC CCTGAGGCGG
 2301 GGATGTGGCA GTGTCTGCTG AGTGA CTGG GACAGGTCTT GCTGGAATCC
 2351 AACATCAAGG TTCTGCCAC ATGGTCGACC CCGGTGCAGC CAATGGCCCT
 2401 GATTTGAGAT CTTTGTGAAG GAACCTTACT TCTGTGGTGT GACATAATTG
 2451 GACAACTAC CTACAGAGAT TTAAAGCTCT AAGGTAAATA TAAAATTTTT
 2501 AAGTGATAA TGTGTTAAAC TACTGATTCT AATTGTTTGT GTATTTTAGA
 2551 TTCCAACCTA TGGAAGTAT GAATGGGAGC AGTGGTGGAA TGCCTTTAAT
 2601 GAGGAAAACC TGTTTTGCTC AGAAGAAATG CCATCTAGTG ATGATGAGGC
 2651 TACTGCTGAC TCTCAACATT CTACTCCTCC AAAAAAGAAG AGAAAGGTAG
 2701 AAGACCCCAA GGACTTTCCT TCAGAATTGC TAAGTTTTTT GAGTCATGCT
 2751 GTGTTTAGTA ATAGA ACTCT TGCTTGCTTT GCTATTTACA CCACAAAGGA
 2801 AAAAGCTGCA CTGCTATACA AGAAAATTAT GGAAAAATAT TCTGTAACCT
 2851 TTATAAGTAG GCATAACAGT TATAATCATA ACATACTGTT TTTTCTTACT
 2901 CCACACAGGC ATAGAGTGTC TGCTATTAAT AACTATGCTC AAAAATTGTG
 2951 TACCTTTAGC TTTTAAATTT GTAAAGGGT TAATAAGGAA TATTTGATGT
 3001 ATAGTGCCTT GACTAGAGAT CATAATCAGC CATACCACAT TTGTAGAGGT

89085519

52/93

FIG. 16(cont'd)

3051 TTTACTTGCT TTAACAAAAC TCCCACACCT CCCCCTGAAC CTGAAACATA
3101 AAATGAATGC AATTGTTGTT GTTAACTTGT TTATTGCAGC TTATAATGGT
3151 TACAAATAAA GCAATAGCAT CACAAATTTC ACAAATAAAG CATTTTTTTC
3201 ACTGCATTCT AGTTGTGGTT TGTCCAAACT CATCAATGTA TCTTATCATG
3251 TCTGGATCCT CTACGCCGGA CGCATCGTGG CCGGCATCAC CGGCGCCACA
3301 GGTGCGGTTG CTGGCGCCTA TATCGCCGAC ATCACCGATG GGAAGATCG
3351 GGCTCGCCAC TTCGGGCTCA TGAGCGCTTG TTTCGGCGTG GGTATGGTGG
3401 CAGGCCCGTG GCCGGGGGAC TGTGCGGCGC CATCTCCTTG CATGCACCAT
3451 TCCTTGCGGC GGCGGTGCTC AACGGCCTCA ACCTACTACT GGGCTGCTTC
3501 CTAATGCAGG AGTCGCATAA GGGAGAGCGT CGACCGATGC CCTTGAGAGC
3551 CTTCAACCCA GTCAGCTCCT TCCGGTGGGC GCGGGGCATG ACTATCGTCG
3601 CCGCACTTAT GACTGTCTTC TTTATCATGC AACTCGTAGG ACAGGTGCCG
3651 GCAGCGCTCT GGGTCATTTT CGGCGAGGAC CGCTTTCGCT GGAGCGCGAC
3701 GATGATCGGC CTGTCGCTTG CGGTATTCGG AATCTTGCAC GCCCTCGCTC
3751 AAGCCTTCGT CACTGGTCCC GCCACCAAAC GTTTCGGCGA GAAGCAGGCC
3801 ATTATCGCCG GCATGGCGGC CGACGCGCTG GGCTACGTCT TGCTGGCGTT
3851 CGCGACCGGA GGCTGGATGG CCTTCCCAT TATGATTCTT CTCGCTTCCG
3901 GCGGCATCGG GATGCCCGCG TTGCAGGCCA TGCTGTCCAG GCAGGTAGAT
3951 GACGACCATC AGGGACAGCT TCAAGGATCG CTCGCGGCTC TTACCAGCCT
4001 AACTTCGATC ACTGGACCGC TGATCGTCAC GGCGATTTAT GCCGCCTCGG
4051 CGAGCACATG GAACGGGTTG GCATGGATTG TAGGCGCCGC CCTATACCTT
4101 GTCTGCCTCC CCGCGTTGCG TCGCGGTGCA TGGAGCCGGG CCACCTCGAC
4151 CTGAATGGAA GCCGGCGGCA CCTCGCTAAC GGATTCACCA CTCCAAGAAT
4201 TGGAGCCAAT CAATTCTTGC GGAGAACTGT GAATGCGCAA ACCAACCCTT
4251 GGCAGAACAT ATCCATCGCG TCCGCCATCT CCAGCAGCCG CACGCGGCGC
4301 ATCTCGGGCC GCGTTGCTGG CGTTTTTCCA TAGGCTCCGC CCCCCTGACG
4351 AGCATCACAA AAATCGACGC TCAAGTCAGA GGTGGCGAAA CCCGACAGGA
4401 CTATAAAGAT ACCAGGCGTT TCCCCCTGGA AGCTCCCTCG TGCGCTCTCC
4451 TGTTCCGACC CTGCCGCTTA CCGGATACCT GTCCGCCTTT CTCCCTTCGG
4501 GAAGCGTGGC GCTTTCTCAA TGCTCACGCT GTAGGTATCT CAGTTCGGTG
4551 TAGGTCGTTT GCTCCAAGCT GGGCTGTGTG CACGAACCCC CCGTTCAGCC
4601 CGACCGCTGC GCCTTATCCG CTAAGTATCG TCTTGAGTCC AACC CGGTAA
4651 GACACGACTT ATCGCCACTG GCAGCAGCCA CTGGTAACAG GATTAGCAGA

3808

89085519


53/93

FIG. 16(cont'd)

4701 GCGAGGTATG TAGGCGGTGC TACAGAGTTC TTGAAGTGGT GGCCTAACTA
4751 CGGCTACACT AGAAGGACAG TATTTGGTAT CTGCGCTCTG CTGAAGCCAG
4801 TTACCTTCGG AAAAAGAGTT GCTAGCTCTT GATCCGGCAA ACAAACCACC
4851 GCTGGTAGCG GTGGTTTTTT TGTTCGCAAG CAGCAGATTA CGCGCAGAAA
4901 AAAAGGATCT CAAGAAGATC CTTTGATCTT TTCTACGGGG TCTGACGCTC
4951 AGTGGAAACGA AAATCAGCT TAAGGGATTT TGGTCATGAG ATTATCAAAA
5001 AGGATCTTCA CCTAGATCCT TTAAATTA AAATGAAGTT TTAAATCAAT
5051 CTAAAGTATA TATGAGTAAA CTTGGTCTGA CAGTTACCAA TGCTTAATCA
5101 GTGAGGCACC TATCTCAGCG ATCTGTCTAT TTCGTTTCATC CATAGTTGCC
5151 TGAATCCCCG TCGTGTAGAT AACTACGATA CGGGAGGGCT TACCATCTGG
5201 CCCCAGTGCT GCAATGATAC CGCGAGACCC ACGCTCACCG GCTCCAGATT
5251 TATCAGCAAT AAACCAGCCA GCCGGAAGGG CCGAGCGCAG AAGTGGTCTT
5301 GCAACTTTAT CCGCCTCCAT CCAGTCTATT AATTGTTGCC GGAAGCTAG
5351 AGTAAGTAGT TCGCCAGTTA ATAGTTTGCG CAACGTTGTT GCCATTGCTG
5401 CAGGCATCGT GGTGTCACGC TCGTCGTTTG GTATGGCTTC ATTCAGCTCC
5451 GGTTCCCAAC GATCAAGGCG AGTTACATGA TCCCCCATGT TGTGCAAAAA
5501 AGCGGTTAGC TCCTTCGGTC CTCCGATCGT TGTCAGAAGT AAGTTGGCCG
5551 CAGTGTTATC ACTCATGGTT ATGGCAGCAC TGCATAATTC TCTTACTGTC
5601 ATGCCATCCG TAAGATGCTT TTCTGTGACT GGTGAGTACT CAACCAAGTC
5651 ATTCTGAGAA TAGTGTATGC GGCGACCGAG TTGCTCTTGC CCGGCGTCAA
5701 CACGGGATAA TACCGCGCCA CATAGCAGAA CTTTAAAAGT GCTCATCATT
5751 GGAAAACGTT CTTCGGGGCG AAAACTCTCA AGGATCTTAC CGCTGTTGAG
5801 ATCCAGTTCC ATGTAACCCA CTCGTGCACC CAACTGATCT TCAGCATCTT
5851 TTACTTTTAC CAGCGTTTCT GGGTGAGCAA AAACAGGAAG GCAAAATGCC
5901 GCAAAAAAGG GAATAAGGGC GACACGGAAA TGTTGAATAC TCATACTCTT
5951 CCTTTTTCAA TATTATTGAA GCATTTATCA GGGTTATTGT CTCATGAGCG
6001 GATACATATT TGAATGTATT TAGAAAAATA AACAAATAGG GGTTCCGCGC
6051 ACATTTCCCC GAAAAGTGCC ACCTGACGTC TAAGAAACCA TTATTATCAT
6101 GACATTAACC TATAAAAATA GGCGTATCAC GAGGCCCTTT CGTCTTCAA

54/93

FIG. 17



3810

89085519

55/93

FIG. 18

61
5' CTA CCT TTT CCA GTG A 3'

62
5' GAT CTC ACT GGA AAG 3'

63
5' CCG CTG ATA GTA A 3'

64
5' GAT CTT ACT ATC A 3'

67
5' CCG GCA GAG CCT GAC CCT GAC CTT GGA GAG CCC C 3'

68
5' CCG GGG GGC TCT CCA AGG TCA CCG TCA GGC TCT C 3'

69
5' CCG GGT AGT AGC CCC TCA GTG CAA TCA 3'

70
5' GAT CTC ATT GCA CTG AGG GGC TAC TAC 3'

71
5' CCG GGT AGT AGC CCC TCA GTG CAA TGT AGG AGT C 3'

72
5' TAG GAC TCC TAC ATT GCA CTG AGG GGC TAC TAC 3'

73
5' CTA CCG GTA AAA ACA TAC AGG CCG GGA AGA CCT CA 3'

74
5' GAT CTC AGG TCT TTC CCC CCC TGT ATG TTT TTA CCC 3'

75
5' CCA GGA TAG TGG CAC CTG GAC ATG CAC TGT CTT GCA
GAA CTG A 3'

76
5' GAT CTC AGT TGT CCA AGA CAC TCC ATG TCC AGG TCC
CAC TAT CCT GCA CCT 3'

56/93

pg394
:BG368 backo file
:soluble T4#9
:AA #3 = LVS
:first 113 AA of T4
:basically up to V1J1

FIG. 19

pg394.seq Length: 5365

```

1  GAATTAATTC CAGCTTGCTG TGGAATGTGT GTCAGTTAGG GTGTGGAAAG
51  TCCCCAGGCT CCCCAGCAGG CAGAAGTATG CAAAGCATGC ATCTCAATTA
101 GTCAGCAACC AGGTGTGGAA AGTCCCCAGG CTCCCCAGCA GGCAGAAGTA
151 TGCAAAGCAT GCATCTCAAT TAGTCAGCAA CCATAGTCCC GCCCCTAACT
201 CCGCCCATCC CGCCCTAAC TCCGCCCAGT TCCGCCCATC CTCCGCCCCA
251 TGGCTGACTA ATTTTTTTTA TTTATGCAGA GGCCGAGGCC GCCTCGGCCT
301 CTGAGCTATT CCAGAAGTAG TGAGGAGGCT TTTTGGAGG GGTCTCTCTC
351 GTATAGAAAC TCGGACCACT CTGAGACGAA GGCTCGCGTC CAGGCCAGCA
401 CGAAGGAGGC TAAGTGGGAG GGGTAGCGGT CGTTGTCCAC TAGGGGGTCC
451 ACTCGCTCCA GGGTGTGAAG ACACATGTCG CCCTCTTCGG CATCAAGGAA
501 GGTGATTGGT TTAYAGGTGT AGGCCACGTG ACCGGGTGTT CCTGAAGGGG
551 GGCTATAAAA GGGGGTGGGG GCGCGTTCGT CCTCACTCTC TTCCGCATCG
601 CTGTCTGCGA GGGCCAGCTG TTGGGCTCGC GGTGAGGAC AAACCTCTCG
651 CGGTCTTTCC AGTACTCTTG GATCGGAAAC CCGTCGGCCT CCGAACGGTA
701 CTCCGCCACC GAGGGACCTG AGCGAGTCCG CATCGACCGG ATCGGAAAAC
751 CTCTCGAGAA AGGCGTCTAA CCAGTCACAG TCGCAAGGTA GGCTGAGCAC
801 CGTGGCGGGC GGCAGCGGGT GCGGTCGGG GTTGTTTCTG GCGGAGGTGC
851 TGCTGATGAT GTAATTAAAG TAGGCGGTCT TGAGACGGCG GATGGTCGAG
901 GTGAGGTGTG GCAGGCTTGA GATCGATCTG GCCATACACT TGAGTGACAA
951 TGACATCCAC TTTGCCTTTC TCTCCACAGG TGTCCACTCC CAGGTCCAAC
1001 TGGATCCAAG CTTCGACTCG AGGAATTCCC CGAAGGAACA AAGCACCCCTC
1051 CCCACTGGGC TCCTGGTTGC AGAGCTCCAA GTCCTCACAC AGATACGCCT
1101 GTTTGAGAAG CACCGGGCAA GAAAGACGCA AGCCAGAGG CCCTGCCATT
1151 TCTGTGGGCT CAGGTCCCTA CTGGCTCAGG CCCCTGCCTC CCTCGGCAAG
1201 GCCACAATGA ACCGGGGAGT CCCTTTTAGG CACTTGCTTC TGGTGCTGCA
1251 ACTGGCGCTC CTCCCAGCAG CCACTCAGGG AAAGAAAGTG GTGCTGGGCA
1301 AAAAAGGGGA TACAGTGGAA CTGACCTGTA CAGCTTCCCA GAAGAAGAGC
1351 ATACAATTCC ACTGAAATTC ACCAG ATAAAGATTC TGGGAAATCA

```

3812

89085519

F! G. 19 (cont'd)

1401 GGGCTCCTTC TTAACATAAG GTCCATCCAA GCTGAATGAT CGCGCTGACT
 1451 CAAGAAGAAG CTTGTGGGAC CAAGGAAACT TTCCCCTGAT CATCAAGAAT
 1501 CTTAAGATAG AAGACTCAGA TACTTACATC TGTGAAGTGG AGGACCAGAA
 1551 GGAGGAGGTG CAATTGCTAG TGTTCGGATT GACTGCCAAC TCTGACACCC
 1601 ACCTGCTTCA GGGGTGATAG TAAGATCTTT GTGAAGGAAC CTTACTTCTG
 1651 TGGTGTGACA TAATTGGACA AACTACCTAC AGAGATTAA AGCTCTAAGG
 1701 TAAATATAAA ATTTTAAAGT GTATAATGTG TTAAACTACT GATTCTAATT
 1751 GTTGTGTAT TTTAGATTCC AACCTATGGA ACTGATGAAT GGGAGCAGTG
 1801 GTGGAATGCC TTTAATGAGG AAAACCTGTT TTGCTCAGAA GAAATGCCAT
 1851 CTAGTGATGA TGAGGCTACT GCTGACTCTC AACATTCTAC TCCTCCAAAA
 1901 AAGAAGAGAA AGGTAGAAGA CCCAAGGAC TTTCCTTCAG AATTGCTAAG
 1951 TTTTTTGAGT CATGCTGTGT TTAGTAATAG AACTCTTGCT TGCTTTGCTA
 2001 TTTACACCAC AAAGGAAAAA GCTGCACTGC TATACAAGAA AATTATGGAA
 2051 AAATATTCTG TAACCTTTAT AAGTAGGCAT AACAGTTATA ATCATAACAT
 2101 ACTGTTTTTT CTTACTCCAC ACAGGCATAG AGTGTCTGCT ATTAATAACT
 2151 ATGCTCAAAA ATTGTGTACC TTTAGCTTTT TAATTTGTAA AGGGGTAAAT
 2201 AAGGAATATT TGATGTATAG TGCCTTGACT AGAGATCATA ATCAGCCATA
 2251 CCACATTTGT AGAGGTTTTA CTTGCTTTAA AAAACCTCCC ACACCTCCCC
 2301 CTGAACCTGA AACATAAAAT GAATGCAATT GTTGTGTGTA ACTTGTTTAT
 2351 TGCAGCTTAT AATGGTTACA AATAAAGCAA TAGCATCACA AATTTACAAA
 2401 ATAAAGCATT TTTTTCCTG CATTCTAGTT GTGGTTTGTC CAAACTCATC
 2451 AATGTATCTT ATCATGTCTG GATCCTCTAC GCCGGACGCA TCGTGGCCGG
 2501 CATCACCGGC GCCACAGGTG CGGTTGCTGG CGCCTATATC GCCGACATCA
 2551 CCGATGGGGA AGATCGGGCT CGCCACTTCG GGCTCATGAG CGCTTGTTTC
 2601 GGGGTGGGTA TGGTGGCAGG CCCGTGGCCG GGGGACTGTT GGGCGCCATC
 2651 TCCTTGCTAG CACCATTCCT TCGGGCGGCG GTGCTCAACG GCCTCAACCT
 2701 ACTACTGGGC TGCTTCCTAA TGCAGGAGTC GCATAAGGGA GAGCGTCGAC
 2751 CGATGCCCTT GAGAGCCTTC AACCCAGTCA GCTCCTTCGG GTGGGCGCGG
 2801 GGCATGACTA TCGTCGCCGC ACTTATGACT GTCTTCTTTA TCATGCAACT
 2851 CGTAGGACAG GTGCCGGCAG CGCTCTGGGT CATTTTCGGC GAGGACCGCT
 2901 TTCGCTGGAG CGCGACGATG ATCGGCCTGT CGCTTGCGGT ATTCGGAATC
 2951 TTGCACGCCC TCGCTCAAGC CTTCGTCACT GGTCCCGCCA CCAAACGTTT
 3001 CGGCGAGAAG CAGGCCATTA TCGCCGGCAT GCGGCGCGAC GCGCTGGGCT

58/93

FIG. 19 (cont'd)

305: ACGTCTTGCT GCGGTTGGLG ACGCGAGGCT GGATGGCCTT CCCCATTATG
 310: ATTCTTCTCG CTTCGGGCGG CATCGGGATG CCGCGTTGC AGGCCATGCT
 315: GTCCAGGCAG GTAGATGACG ACCATCAGGG ACAGCTTCAA GGATCGCTCG
 320: CGGCTCTTAC CAGCCTAACT TCGATCACTG GACCGCTGAT CGTCACGGCG
 325: ATTTATGCCG CCTCGGCGAG CACATGGAAC GGGTTGGCAT GGATTGTAGG
 330: CGCCGCCCTA TACCTTGCTT GCCTCCCCGC GTTGCGTCGC GGTGCATGGA
 335: GCCGGGCCAC CTCGACCTGA ATGGAAGCCG GCGGCACCTC GCTAACGGAT
 340: TCACCACTCC AAGAATTGGA GCCAATCAAT TCTTGCGGAG AACTGTGAAT
 345: GCGCAAACCA ACCCTTGGA GAACATATCC ATCGCGTCCG CCATCTCCAG
 350: CAGCCGCACG CGGCGCATCT CGGGCCGCGT TGCTGGCGTT TTTCCATAGG
 355: CTCCGCCCCC CTGACGAGCA TCACAAAAAT CGACGCTCAA GTCAGAGGTG
 360: GCGAAACCCG ACAGGACTAT AAAGATACCA GCGGTTTCCC CCTGGAAGCT
 365: CCCTCGTGCG CTCTCCTGTT CCGACCTGC CGCTTACCGG ATACCTGTCC
 370: GCCTTTCTCC CTTCGGGAAG CGTGGCGCTT TCTCAATGCT CACGCTGTAG
 375: GTATCTCAGT TCGGTGTAGG TCGTTCGCTC CAAGCTGGGC TGTGTGCACG
 380: AACCCCCCGT TCAGCCCGAC CGCTGCGCCT TATCCGGTAA CTATCGTCTT
 385: GAGTCCAACC CGGTAAGACA CGACTTATCG CCACTGGCAG CAGCCACTGS
 390: TAACAGGATT AGCAGAGCGA GGTATGTAGG CGGTGCTACA GAGTTCTTGA
 395: AGTGGTGGCC TAACTACGGC TACACTAGAA GGACAGTATT TGGTATCTGC
 400: GCTCTGCTGA AGCCAGTTAC CTTCGGAAAA AGAGTTGGTA GCTCTTGATC
 405: CCGCAAACAA ACCACCGCTG GTAGCGGTGG TTTTTTTGTT TGCAAGCAGC
 410: AGATTACGCG CAGAAAAAAA GGATCTCAAG AAGATCCTTT GATCTTTTCT
 415: ACGGGGTCTG ACGCTCAGTG GAACGAAAAAC TCACGTAAAG GGATTTTGGT
 420: CATGAGATTA TCAAAAAGGA TCTTCACCTA GATCCTTTTA AATTAAAAAT
 425: GAAGTTTTAA ATCAATCTAA AGTATATATG AGTAAACTTG GTCTGACAGT
 430: TACCAATGCT TAATCAGTCA GGCACCTATC TCAGCGATCT GTCTATTTCTG
 435: TTCATCCATA GTTGCTGAC TCCCCGTCGT GTAGATAACT ACGATACGGG
 440: AGGGCTTACC ATCTGGCCCC AGTGCTGCAA TGATACCGCG AGACCCACGC
 445: TCACCGGCTC CAGATTTATC AGCAATAAAC CAGCCAGCCG GAAGGGCCGA
 450: GCGCAGAAGT GGTCCTGCAA CTTTATCCGC CTCCATCCAG TCTATTAATT
 455: GTTGCCGGGA AGCTAGAGTA AGTAGTTCGC CAGTTAATAG TTTGCGCAAC
 460: GTTGTGCCA TTGCTGCGG CATCGTGGTG TCACGCTCGT CGTTTGGTAT
 465: GCTTCATTC AGCTCCGCTT CCAACGATC AAGGCGAGTT ACATGATCCC

59/93

FIG. 19 (cont'd)

4701 CCATGTTGTG CAAAAAAGCG GTTAGCTCCT TCGGTCCTCC GATCGTTGTC
4751 AGAAGTAAGT TGGCCGCAGT GTTATCACTC ATGGTTATGG CAGCACTGCA
4801 TAATTCTCTT ACTGTCATGC CATCCGTAAG ATGCTTTTCT GTGACTGGTG
4851 AGTACTCAAC CAAGTCATTC TGAGAATAGT GTATGCGGCG ACCGAGTTGC
4901 TCTTGCCCGG CGTCAACACG GGATAATACC GCGCCACATA GCAGAACTTT
4951 AAAAGTGCTC ATCATTGGAA AACGTTCTTC GGGGCGAAAA CTCTCAAGGA
5001 TCTTACCGCT GTTGAGATCC AGTTCGATGT AACCCACTCG TGCACCCAAC
5051 TGATCTTCAG CATCTTTTAC TTTCAACCAGC GTTTCTGGGT GAGCAAAAAC
5101 AGGAAGGCAA AATGCCGCAA AAAAGGGAAT AAGGGCGACA CGGAAATGTT
5151 GAATACTCAT ACTCTTCCTT TTTCAATATT ATTGAAGCAT TTATCAGGGT
5201 TATTGTCTCA TGAGCGGATA CATATTTGAA TGTATTTAGA AAAATAAACA
5251 AATAGGGGTT CCGCGCACAT TTCCCCGAAA AGTGCCACCT GACGTCTAAG
5301 AAACCATTAT TATCATGACA TTAACCTATA AAAATAGGCG TATCAGGAGG
5351 CCCTTTCGTC TTCAA

89085519

60/93

FIG. 20

pBG396
 :BG368 backbone
 :solubl T4#12
 :AA #3 = LYS

bg396.seq Length: 5518

```

1 GAATTAATTC CAGCTTGCTG TGAATGTGT GTCAGTTAGG GTGTGGAAAG
51 TCCCCAGGCT CCCCAGCAGG CAGAAGTATG CAAAGCATGC ATCTCAATTA
101 GTCAGCAACC AGGTGTGGAA AGTCCCCAGG CTCCCCAGCA GCCAGAAGTA
151 TGCAAAGCAT GCATCTCAAT TAGTCAGCAA CCATAGTCCC GCCCCTAACT
201 CCGCCCATCC CGCCCCTAAC TCCGCCCAGT TCCGCCCATT CTCCGCCCCA
251 TGGCTGACTA ATTTTTTTTA TTTATGCAGA GGCCGAGGCC GCCTCGGCCT
301 CTGAGCTATT CCAGAAGTAG TGAGGAGGCT TTTTGGAGG GGTCTCTCTC
351 GTATAGAAAC TCGGACCACT CTGAGACGAA GGCTCGCGTC CAGGCCAGCA
401 CGAAGGAGGC TAAGTGGGAG GGGTAGCGGT CGTTGTCCAC TAGGGGGTCC
451 ACTCGCTCCA GGGTGTGAAG ACACATGTCT CCCTCTTCGG CATCAAGGAA
501 GGTGATTGGT TTATAGGTGT AGGCCACGTG ACCGGGTGTT CCTGAAGGGG
551 GGCTATAAAA GGGGGTGGGG GCGCGTTCGT CCTCACTCTC TTCCGCATCG
601 CTGTCTGCGA GGGCCAGCTG TTGGGCTCGC GGTGAGGAC AACTCTTCG
651 CGGTCTTTCC AGTACTCTTG GATCGGAAAC CCGTCGGCCT CCGAACGGTA
701 CTCCGCCACC GAGGGACCTG AGCGAGTCCG CATCGACCGG ATCGGAAAAC
751 CTCTCGAGAA AGGCGTCTAA CCAGTCACAG TCGCAAGGTA GGCTGAGCAC
801 CGTGGCGGGC GGCAGCGGGT GCGGTCGGG GTTGTCTCTG GCGGAGGTGC
851 TGCTGATGAT GTAATTAAAG TAGGCGGTCT TGAGACGGCG GATGGTCGAG
901 GTGAGGTGTG GCAGGCTTGA GATCGATCTG GCCATACACT TGAGTGACAA
951 TGACATCCAC TTTGCCTTTC TCTCCACAGG TGTCCACTCC CAGGTCCAAC
1001 TGGATCCAAG CTTCGACTCG AGGAATTCCC CGAAGGAACA AAGCACCCCTC
1051 CCCACTGGGC TCCTGGTTGC AGAGCTCCAA GTCCTCACAC AGATACGCCT
1101 GTTTGAGAAG CAGCGGGCAA GAAAGACGCA AGCCAGAGG CCCTGCCATT
1151 TCTGTGGGCT CAGGTCCCTA CTGGCTCAGG CCCCTGCCTC CCTCGGCAAG
1201 GCCACAATGA ACCGGGGAGT CCCTTTTAGG CACTTGCTTC TGGTGCTGCA
1251 ACTGGCGCTC CTCCCAGCAG CCACTCAGGG AAAGAAAGTG GTGCTGGGCA
1301 AAAAAGGGGA TACAGTGGAA CTGACCTGTA CAGCTTCCCA GAAGAAGAGC
1351 ATACAATTCC ACTGGAAGAA GTGCAACAG ATAAAGATTC TGGGAAATCA
1401 GGGCTCTTTC TTAACATAAG GTCCATCCAA GCTGAATGAT CGCGCTGACT
  
```

3816

89085519
SUBSTITUTE SHEET

61/93

FIG. 20 (cont'd)

1451 CAAGAAGAAG CTTGTGGGAC CAAGGAAACT TTCCCCTGAT CATCAAGAAT
 1501 CTTAAGATAG AAGACTCAGA TACTTACATC TGTGAAGTGG AGGACCAGAA
 1551 GGAGGAGGTG CAATTGCTAG TGTTCCGATT GACTGCCAAC TCTGACACCC
 1601 ACCTGCTTCA GGGGCAGAGC CTGACCCTGA CCTTGGAGAG CCCCCCTGGT
 1651 AGTAGCCCCCT CAGTGCAATG TAGGAGTCCA AGGGGTAAAA ACATACAGGG
 1701 GGGGAAGACC CTCTCCCTGT CTCAGCTGGA GCTCCAGGAT AGTGGCACCT
 1751 GGACATGCAC TGTCTTGACAG AACTGAGATC TTTGTGAAGG AACCTTACTT
 1801 CTGTGGTGTG ACATAATTGG ACAAACCTACC TACAGAGATT TAAAGCTCTA
 1851 AGGTAAATAT AAAATTTTAA AGTGTATAAT GTGTTAAACT ACTGATTCTA
 1901 ATTGTTTGTG TATTTTAGAT TCCAACCTAT GGAACCTGATG AATGGGAGCA
 1951 GTGGTGGAAAT GCCTTTAATG AGGAAAACCT GTTTTGCTCA GAAGAAATGC
 2001 CATCTAGTGA TGATGAGGCT ACTGCTGACT CTCAACATTC TACTCCTCCA
 2051 AAAAAGAAGA GAAAGGTAGA AGACCCCAAG GACTTTCCTT CAGAATTGCT
 2101 AAGTTTTTTG AGTCATGCTG TGTTTAGTAA TAGAACTCTT GCTTGCTTTG
 2151 CTATTTACAC CACAAAGGAA AAAGCTGCAC TGCTATACAA GAAAATTATG
 2201 GAAAAATATT CTGTAACCTT TATAAGTAGG CATAACAGTT ATAATCATAA
 2251 CATACTGTTT TTTCTTACTC CACACAGGCA TAGAGTGTCT GCTATTAATA
 2301 ACTATGCTCA AAAATTGTGT ACCTTTAGCT TTTTAATTTG TAAAGGGGTT
 2351 AATAAGGAAT ATTTGATGTA TAGTGCCTTG ACTAGAGATC ATAATCAGCC
 2401 ATACCACATT TGTAAGAGTT TTAAGTGTCT TAAAAAACCT CCCACACCTC
 2451 CCCCTGAACC TGAACATAA AATGAATGCA ATTGTTGTTG TTAAGTTGTT
 2501 TATTGCAGCT TATAATGGTT ACAAATAAAG CAATAGCATC ACAAATTTCA
 2551 CAAATAAAGC ATTTTTTTCA CTGCATTCTA GTTGTGGTTT GTCCAAACTC
 2601 ATCAATGTAT CTTATCATGT CTGGATCCTC TACGCCGGAC GCATCGTGGC
 2651 CGGCATCACC GGCGCCACAG GTGCGGTTGC TGGCGCCTAT ATCGCCGACA
 2701 TCACCGATGG GGAAGATCGG GCTCGCCACT TCGGGCTCAT GAGCGCTTGT
 2751 TTCGGCGTGG GTATGGTGGC AGGCCCGTGG CCGGGGGACT GTTGGGCGCC
 2801 ATCTCTTGC ATGCACCATT CCTTGCGGCG GCGGTGCTCA ACGGCCTCAA
 2851 CCTACTACTG GGCTGCTTCC TAATGCAGGA GTCGCATAAG GGAGAGCGTC
 2901 GACCGATGCC CTTGAGAGCC TTCAACCCAG TCAGCTCCTT CCGGTGGGCG
 2951 CGGGGCATGA CTATCGTCGC CGCACTTATG ACTGTCTTCT TTATCATGCA
 3001 ACTCGTAGGA CAGGTGCGGAGGCTCTG GGTCATTTTC GGCGAGGACC
 3051 GCTTTGGTGG GAGCGCGACG ATGATCGGCC TGTGCTTGC GGTATTCGGA

3817

89085319

SEQUENCE LIST

62/93

FIG. 20 (cont'd)

3101 ATCTTGCAAG CCCTCGCTCA AGCCTTCGTC ACTGGTCCCG CCACCAAACG
3151 TTTCGGCGAG AAGCAGGCCA TTATCGCCGG CATGGCGGCC GACGCGCTGG
3201 GCTACGTCTT GCTGGCGTTC GCGACGCGAG GCTGGATGGC CTTCCCCATT
3251 ATGATTCTTC TCGCTTCCGG CGGCATCGGG ATGCCCGCGT TGCAGGCCAT
3301 GCTGTCCAGG CAGGTAGATG ACGACCATCA GGGACAGCTT CAAGGATCGC
3351 TCGCGGCTCT TACCAGCCTA ACTTCGATCA CTGGACCGCT GATCGTCACG
3401 GCGATTTATG CCGCCTCGGC GAGCACATGG AACGGGTTGG CATGGATTGT
3451 AGGCGCCGCC CTATACCTTG TCTGCCTCC CGCGTTGCGT CGCGGTGCAT
3501 GGAGCCGGGC CACCTCGACC TGAATGGAAG CCGGCGGCAC CTCGCTAACG
3551 GATTACCCAC TCCAAGAATT GGAGCCAA^TC AATTCTTGCG GAGAAGTGTG
3601 AATGCGCAAA CCAACCTTG GCAGAACATA TCCATCGCGT CCGCCATCTC
3651 CAGCAGCCGC ACGCGGCGCA TCTCGGGCCG CGTTGCTGGC GTTTTTCCAT
3701 AGGCTCCGCC CCCCTGACGA GCATCACAAA AATCGACGCT CAAGTCAGAG
3751 GTGGCGAAAC CCGACAGGAC TATAAAGATA CCAGGCGTTT CCCCCTGGAA
3801 GCTCCCTCGT GCGCTCTCCT GTTCCGACCC TGCCGCTTAC CGGATACCTG
3851 TCCGCCTTTC TCCCTTCGGG AAGCGTGGCG CTTTCTCAAT GCTCAGCTG
3901 TAGGTATCTC AGTTCGGTGT AGGTCGTTCT CTCCAAGCTG GGCTGTGTGC
3951 ACGAACCCCC CGTTCAGCCC GACCGCTGCG CCTTATCCGG TAACTATCGT
4001 CTTGAGTCCA ACCCGGTAAG ACACGACTTA TCGCCACTGG CAGCAGCCAC
4051 TGGTAACAGG ATTAGCAGAG CGAGGTATGT AGGCGGTGCT ACAGAGTTCT
4101 TGAAGTGGTG GCCTAACTAC GGCTACACTA GAAGGACAGT ATTTGGTATC
4151 TGCGCTCTGC TGAAGCCAGT TACCTTCGGA AAAAGAGTTG GTAGCTCTTG
4201 ATCCGGCAAA CAAACCACCG CTGGTAGCGG TGGTTTTTTT GTTTGCAAGC
4251 AGCAGATTAC GCGCAGAAAA AAAGGATCTC AAGAAGATCC TTTGATCTTT
4301 TCTACGGGGT CTGACGCTCA GTGGAACGAA AACTCACGTT AAGGGATTTT
4351 GGTCAATGAGA TTATCAAAAA GGATCTTCAC CTAGATCCTT TTAAATTA
4401 AATGAAGTTT TAAATCAATC TAAAGTATAT ATGAGTAAAC TTGGTCTGAC
4451 AGTTACCAAT GCTTAATCAG TGAGGCACCT ATCTCAGCGA TCTGTCTATT
4501 TCGTTCATCC ATAGTTGCCT GACTCCCCGT CGTGTAGATA ACTACGATAC
4551 GGGAGGGCTT ACCATCTGGC CCCAGTGCTG CAATGATACC GCGAGACCCA
4601 CGCTCACCGG CTCCAGATTT ATCAGCAATA AACCAGCCAG CCGGAAGGGC
4651 CGAGCGCAGA AGTGGTCTG CAAC^TTTATC CGCCTCCATC CAGTCTATTA
4701 ATTGTTGCCG GGAAGCTAGA GTAAGTAGTT CGCCAGTTAA TAGTTTGCGC

3818

89085519

63/93

FIG. 20 (cont'd)

4751 AACGTTGTTG CCATTGCTGC AGGCATCGTG GTGTCACGCT CGTCGTTTGG
4801 TATGGCTTCA TTCAGCTCCG GTTCCCAACG ATCAAGGCGA GTTACATGAT
4851 CCCCCATGTT GTGCAAAAAA GCGGTTAGCT CCTTCGGTCC TCCGATCGTT
4901 GTCAGAAGTA AGTTGGCCGC AGTGTTATCA CTCATGGTTA TGGCAGCACT
4951 GCATAATTCT CTTACTGTCA TGCCATCCGT AAGATGCTTT TCTGTGACTG
5001 GTGAGTACTC AACCAAGTCA TTCTGAGAAT AGTGTATGCG GCGACCGAGT
5051 TGCTCTTGCC CGGCGTCAAC ACGGGATAAT ACCGCGCCAC ATAGCAGAAC
5101 TTTAAAAGTG CTCATCATTG GAAAACGTTT TTCGGGGCGA AAACCTCTCA
5151 GGATCTTACC GCTGTTGAGA TCCAGTTCGA TGTAACCCAC TCGTGACCCC
5201 AACTGATCTT CAGCATCTTT TACTTTTACC AGCGTTTCTG GGTGAGCAAA
5251 AACAGGAAGG CAAAATGCCG CAAAAAAGGG AATAAGGGCG ACACGGAAAT
5301 GTTGAATACT CATACTCTTC CTTTTTCAAT ATTATTGAAG CATTTATCAG
5351 GGTATTGTG TCATGAGCGG ATACATATTT GAATGTATTT AGAAAAATAA
5401 ACAAATAGGG GTTCCGCGCA CATTTCCCCG AAAAGTGCCA CCTGACGTCT
5451 AAGAAACCAT TATTATCATG ACATTAACCT ATAAAAATAG GCGTATCAGG
5501 AGGCCCTTTC GTCTTCAA

89085519

3819

SUBSTITUTE SHEET

64/93

PBQ393
 :BG368 backbone
 :solubl T4#8
 :AA #3 = LYS
 : "perfect" Stu/first 182 AA of T4
 : basically up to V2J2

FIG. 21

bg393.seq Length: 5566

```

1  GAATTAATTC CAGCTTGCTG TGGAAATGTGT GTCAGTTAGG GTGTGGAAAG
51  TCCCCAGGCT CCCCAGCAGG CAGAAGTATG CAAAGCATGC ATCTCAATTA
101 GTCAGCAACC AGGTGTGGAA AGTCCCCAGG CTCCCCAGCA GGCAGAAGTA
151 TGCAAAGCAT GCATCTCAAT TAGTCAGCAA CCATAGTCCC GCCCCTAACT
201 CCGCCCATCC CGCCCCTAAC TCCGCCCAGT TCCGCCCAT TCTCCGCCCA
251 TGGCTGACTA ATTTTTTTTA TTTATGCAGA GGCCGAGGCC GCCTCGGCCT
301 CTGAGCTATT CCAGAAGTAG TGAGGAGGCT TTTTGGAGG GGTCCTCCTC
351 GTATAGAAAC TCGGACCACT CTGAGACGAA GGCTCGCGTC CAGGCCAGCA
401 CGAAGGAGGC TAAGTGGGAG GGGTAGCGGT CGTTGTCCAC TAGGGGGTCC
451 ACTCGCTCCA GGGTGTGAAG ACACATGTCG CCCTCTTCGG CATCAAGGAA
501 GGTGATTGGT TTATAGGTGT AGGCCACGTG ACCGGGTGTT CCTGAAGGGG
551 GGCTATAAAA GGGGGTGGGG GCGCGTTCGT CCTCACTCTC TTCCGCATCG
601 CTGTCTGCGA GGGCCAGCTG TTGGGCTCGC GGTTGAGGAC AAACCTCTTCG
651 CGGTCTTTCC AGTACTCTTG GATCGGAAAC CCGTCGGCCT CCGAACGGTA
701 CTCCGCCACC GAGGGACCTG AGCGAGTCCG CATCGACCGG ATCGGAAAAC
751 CTCTCGAGAA AGGCGTCTAA CCAGTCACAG TCGCAAGGTA GGCTGAGCAC
801 CGTGGCGGGC GGCAGCGGGT GGCGGTCGGG GTTGTCTCTG GCGGAGGTGC
851 TGCTGATGAT GTAATTAAAG TAGGCGGTCT TGAGACGGCG GATGGTCGAG
901 GTGAGGTGTG GCAGGCTTGA GATCGATCTG GCCATACACT TGAGTGACAA
951 TGACATCCAC TTTGCCTTTC TCTCCACAGG TGTCCACTCC CAGGTCCAAC
1001 TGGATCCAAG CTTCGACTCG AGGAATTCCC CGAAGGAACA AAGCACCTC
1051 CCCACTGGGC TCCTGGTTGC AGAGCTCCAA GTCCTCACAC AGATACGCCT
1101 GTTTGAGAAG CAGCGGGCAA GAAAGACGCA AGCCAGAGG CCCTGCCATT
1151 TCTGTGGGCT CAGGTCCCTA CTGGCTCAGG CCCCTGCCTC CCTCGGCAAG
1201 GCCACAATGA ACCGGGGAGT CCCTTTTAGG CACTTGCTTC TGGTGCTGCA
1251 ACTGGCGCTC CTCCCAGCAG CCACTCAGGG AAAGAAAGTG GTGCTGGGCA
1301 AAAAAGGGGA TACAGTGGAA CTCACCTGTA CAGCTTCCCA GAAGAAGAGC
1351 ATACAATTCC ACTGGAAAAA CTCCAACCAG ATAAAGATTG TGGGAAATCA

```

89085519

65/93

FIG. 21 (cont'd)

1401 GGGCTCCTTC TTAAGTAAAG GTCCATCCAA GCTGAATGAT CGCGCTGACT
1451 CAAGAAGAAG CTTGTGGGAC CAAGGAAACT TTCCCCTGAT CATCAAGAAT
1501 CTTAAGATAG AAGACTCAGA TACTTACATC TGTGAAGTGG AGGACCAGAA
1551 GGAGGAGGTG CAATTGCTAG TGTTCGGATT GACTGCCAAC TCTGACACCC
1601 ACCTGCTTCA GGGGCAGAGC CTGACCCTGA CCTTGGAGAG CCCCCCTGGT
1651 AGTAGCCCTT CAGTGCAATG TAGGAGTCCA AGGGGTAAAA ACATACAGGG
1701 GGGGAAGACC CTCTCCGTGT CTCAGCTGGA GCTCCAGGAT AGTGGCACCT
1751 GGACATGCAC TGTCTTGACG AACCAGAAGA AGGTGGAGTT CAAAATAGAC
1801 ATCGTGGTGC TAGCTTTCCA GTGAGATCTT TGTGAAGGAA CCTTACTTCT
1851 GTGGTGTGAC ATAATTGGAC AAACCTACCTA CAGAGATTTA AAGCTCTAAG
1901 GTAAATATAA AATTTTAAAG TGTATAATGT GTTAAACTAC TGATTCTAAT
1951 TGTTTGTGTA TTTTAGATTG CAACCTATGG AACTGATGAA TGGGAGCAGT
2001 GGTGGAATGC CTTTAATGAG GAAACCTGT TTTGCTCAGA AGAAATGCCA
2051 TCTAGTGATG ATGAGGCTAC TGCTGACTCT CAACATTCTA CTCCTCCAAA
2101 AAAGAAGAGA AAGGTAGAAG ACCCCAAGGA CTTTCCTTCA GAATTGCTAA
2151 GTTTTTTGAG TCATGCTGTG TTTAGTAATA GAACCTTGC TTGCTTTGCT
2201 ATTTACACCA CAAAGGAAAA AGCTGCACTG CTATACAAGA AAATTATGGA
2251 AAAATATTCT GTAACCTTTA TAAGTAGGCA TAACAGTTAT AATCATAACA
2301 TACTGTTTTT TCTTACTCCA CACAGGCATA GAGTGTCTGC TATTAATAAC
2351 TATGCTCAAA AATTGTGTAC CTTTAGCTTT TTAATTTGTA AAGGGGTAA
2401 TAAGGAATAT TTGATGTATA GTGCCTTGAC TAGAGATCAT AATCAGCCAT
2451 ACCACATTTG TAGAGGTTTT ACTTGCTTTA AAAACCTCC CACACCTCCC
2501 CCTGAACCTG AAACATAAAA TGAATGCAAT TGTTGTTGTT AACTTGTTTA
2551 TTGCAGCTTA TAATGGTTAC AAATAAAGCA ATAGCATCAC AAATTCACA
2601 AATAAAGCAT TTTTTCCTT GCAATCTAGT TGTGGTTTGT CCAAACCTCAT
2651 CAATGTATCT TATCATGTCT GGATCCTCTA CGCCGGACGC ATCGTGGCCG
2701 GCATCACCGG CGCCACAGGT GCGGTTGCTG GCGCCTATAT CGCCGACATC
2751 ACCGATGGGG AAGATCGGGC TCGCCACTTC GGGCTCATGA GCGCTTGTTT
2801 CGGCGTGGGT ATGGTGGCAG GCCCGTGGCC GGGGGACTGT TGGGCGCCAT
2851 CTCCTTGCAT GCACCATTC TGTGGCGGCG GGTGCTCAAC GGCCTCAACC
2901 TACTACTGGG CTGCTTCCTA ATGCAGGAGT CGCATAAGGG AGAGCGTCGA
2951 CCGATGCCCT TGAGAGCTT CAACCGAGTC AGCTCCTTCC GGTGGGCGCG
3001 GGGCATGACT ATCGTCGCGG CACTTATGAC TGTCTTCTTT ATCATGCAAC

89085519

66/93

FIG. 21 (cont'd)

3051 TCGTAGGACA GGTGCCGGCA GCGCTCTGGG TCATTTTCGG CGAGGACCGC
3101 TTTCGCTGGA GCGCGACGAT GATCGGCCCG TCGCTTGCGG TATTCGGAAT
3151 CTTGCACGCC CTCGCTCAAG CCTTCGTCAAC TGGTCCCGCC ACCAAACGTT
3201 TCGGCGAGAA GCAGGCCATT ATCGCCGGCA TGGCGGCCGA CGCGCTGGGC
3251 TACGTCTTGC TGGCGTTCGC GACGCGAGGC TGGATGGCCT TCCCCATTAT
3301 GATTCTTCTC GCTTCCGGCG GCATCGGGAT GCGCGCGTTG CAGGCCATGC
3351 TGTCCAGGCA GGTAGATGAC GACCATCAGG GACAGCTTCA AGGATCGCTC
3401 GCGGCTCTTA CCAGCCTAAC TTCGATCACT GGACCGCTGA TCGTCACGGC
3451 GATTTATGCC GCCTCGGCGA GCACATGGAA CGGGTTGGCA TGGATTGTAG
3501 GCGCCGCCCT ATACCTTGTC TGCCTCCCCG CGTTGCGTCG CGGTGCATGG
3551 AGCCGGGCCA CCTCGACCTG AATGGAAGCC GGCGGCACCT CGCTAACGGA
3601 TTCACCACTC CAAGAATTGG AGCCAATCAA TTCTTGCGGA GAACTGTGAA
3651 TGCGCAAACC AACCCTTGGC AGAACATATC CATCGCGTCC GCCATCTCCA
3701 GCAGCCGCAC GCGGCGCATC TCGGGCCGCG TTGCTGGCGT TTTTCCATAG
3751 GCTCCGCCCC CCTGACGAGC ATCACAAAAA TCGACGCTCA AGTCAGAGGT
3801 GCGGAAACCC GACAGGACTA TAAAGATACC AGGCGTTTCC CCCTGGAAGC
3851 TCCCTCGTGC GCTCTCCTGT TCCGACCCTG CCGCTTACCG GATACCTGTC
3901 CGCCTTTCTC CCTTCGGGAA GCGTGGCGCT TTCTCAATGC TCACGCTGTA
3951 GGTATCTCAG TTCGGTGTAG GTCGTTGCT CCAAGCTGGG CTGTGTGCAC
4001 GAACCCCCCG TTCAGCCCGA CCGCTGCGCC TTATCCGGTA ACTATCGTCT
4051 TGAGTCCAAC CCGGTAAGAC ACGACTTATC GCCACTGGCA GCAGCCACTG
4101 GTAACAGGAT TAGCAGAGCG AGGTATGTAG GCGGTGCTAC AGAGTTCTTG
4151 AAGTGGTGGC CTAACACGG CTACACTAGA AGGACAGTAT TTGGTATCTG
4201 CGCTCTGCTG AAGCCAGTTA CCTTCGGAAA AAGAGTTGGT AGCTCTTGAT
4251 CCGGCAAAEA AACCACCGCT GGTAGCGGTG GTTTTTTTGT TTGCAAGCAG
4301 CAGATTACGC GCAGAAAAAA AGGATCTCAA GAAGATCCTT TGATCTTTTC
4351 TACGGGGTCT GACGCTCAGT GGAACGAAAA CTCACGTAA GGGATTTTGG
4401 TCATGAGATT ATCAAAAAGG ATCTTCACCT AGATCCTTTT AAATTA AAAA
4451 TGAAGTTTTA AATCAATCTA AAGTATATAT GAGTAAACTT GGTCTGACAG
4501 TTACCAATGC TTAATCAGTG AGGCACCTAT CTCAGCGATC TGTCTATTTC
4551 GTTCATCCAT AGTTGCCTGA CTCCCCGTCG TGTAATAAC TACGATACGG
4601 GAGGGCTTAC CATCTGGCCC CAGTCTGCA ATGATACCGC GAGACCCACG
4651 CTCACCGGCT CCAGATTTAT CAGCAATAAA CCAGCCAGCC GGAAGGGCCG

67/93

FIG. 21 (cont'd)

4701 AGCGCAGAAG TGGTCCTGCA ACTTTATCCG CCTCCATCCA GTCTATTAAT
4751 TGTTGCCGGG AAGCTAGAGT AAGTAGTTCG CCAGTTAATA GTTTGCGCAA
4801 CGTTGTTGCC ATTGCTGCAG GCATCGTGTT GTCACGCTCG TCGTTTGGTA
4851 TGGCTTCATT CAGCTCCGGT TCCCAACGAT CAAGGCGAGT TACATGATCC
4901 CCCATGTTGT GCAAAAAAGC GGTTAGCTCC TTCGGTCCTC CGATCGTTGT
4951 CAGAAGTAAG TTGGCCGCAG TGTTATCACT CATGGTTATG GCAGCACTGC
5001 ATAATTCTCT TACTGTCATG CCATCCGTAA GATGCTTTTC TGTGACTGGT
5051 GAGTACTCAA CCAAGTCATT CTGAGAATAG TGTATGCGGC GACCGAGTTG
5101 CTCTTGCCCG GCGTCAACAC GGGATAATAC CGCGCCACAT AGCAGAACTT
5151 TAAAAGTGCT CATCATTGGA AAACGTTCTT CGGGGCGAAA ACTCTCAAGG
5201 ATCTTACCGC TGTTGAGATC CAGTTCGATG TAACCCACTC GTGCACCCAA
5251 CTGATCTTCA GCATCTTTTA CTTTCACCAG CGTTTCTGGG TGAGCAAAAA
5301 CAGGAAGGCA AAATGCCGCA AAAAAGGGAA TAAGGGCGAC ACGGAAATGT
5351 TGAATACTCA TACTCTTCCT TTTTCAATAT TATTGAAGCA TTTATCAGGG
5401 TTATTGTCTC ATGAGCGGAT ACATATTTGA ATGTATTTAG AAAAAATAAC
5451 AAATAGGGGT TCCGCGCACA TTTCCCCGAA AAGTGCCACC TGACGTCTAA
5501 GAAACCATTA TTATCATGAC ATTAACCTAT AAAAAATAGGC GTATCACGAG
5551 GCCCTTTCGT CTTCAA

89085519

68/93

FIG. 22

p86395
 :BG368 backbone
 :soluble T4#10
 :AA #3 = LVS
 :first 131 AA of T4

bg395.seq Length: 5413

```

1  GAATTAATTC CAGCTTGCTG TGAATGTGT GTCAGTTAGG GTGTGGAAAG
51  TCCCCAGGCT CCCCAGCAGG CAGAAAGTATG CAAAGCATGC ATCTCAATTA
101 GTCAGCAACC AGGTGTGGAA AGTCCCCAGG CTCCCCAGCA GGCAGAAGTA
151 TGCAAAGCAT GCATCTCAAT TAGTCAGCAA CCATAGTCCC GCCCCTAACT
201 CCGCCCATCC CGCCCTAAC TCCGCCAGT TCCGCCATT CTCCGCCCA
251 TGGCTGACTA ATTTTTTTTA TTTATGCAGA GGCCGAGGCC GCCTCGGCCT
301 CTGAGCTATT CCAGAAGTAG TGAGGAGGCT TTTTGGAGG GGTCTCTCTC
351 GTATAGAAAC TCGGACCACT CTGAGACGAA GGCTCGCGTC CAGGCCAGCA
401 CGAAGGAGGC TAAGTGGGAG GGGTAGCGGT CGTTGTCCAC TAGGGGGTCC
451 ACTCGCTCCA GGGTGTGAAG ACACATGTCG CCCTCTTCGG CATCAAGGAA
501 GGTGATTGGT TTATAGGTGT AGGCCACGTG ACCGGGTGTT CCTGAAGGGG
551 GGCTATAAAA GGGGGTGGGG GCGCGTTCGT CCTCACTCTC TTCCGCATCG
601 CTGTCTGCGA GGGCCAGCTG TTGGGCTCGC GGTTGAGGAC AAACCTCTTCG
651 CGGTCTTTCC AGTACTCTTG GATCGGAAAC CCGTCGGCCT CCGAACGGTA
701 CTCCGCCACC GAGGGACCTG AGCGAGTCCG CATCGACCGG ATCGGAAAAC
751 CTCTCGAGAA AGGCGTCTAA CCAGTCACAG TCGCAAGGTA GGCTGAGCAC
801 CGTGGCGGGC GGCAGCGGGT GGCGGTCGGG GTTGTCTCTG GCGGAGGTGC
851 TGCTGATGAT GTAATTAAG TAGGCGGTCT TGAGACGGCG GATGGTCGAG
901 GTGAGGTGTG GCAGGLTTGA GATCGATCTG GCCATACACT TGAGTGACAA
951 TGAGATCCAC TTGCTCTTC TCTCCACAGG TGTCCTCTCC CAGGTCCAA
1001 TGGATCCAAG CTTCGACTCG AGGAATTCCC CGAAGGAACA AAGCACCTC
1051 CCCACTGGGC TCCTGGTTCG AGAGCTCCAA GTCCTCACAC AGATACGCCT
1101 GTTTGAGAAG CAGCGGGCAA GAAAGACGCA AGCCCAGAGG CCCTGCCATT
1151 TCTGTGGGCT CAGGTCCCTA CTGGCTCAGG CCCTGCCTC CCTCGGCAAG
1201 GCCACAATGA ACCGGGGAGT CCCTTTTAGG CACTTGCTTC TGGTGCTGCA
1251 ACTGGCGCTC CTCCCAGCAG CCACTCAGGG AAAGAAAGTG GTGCTGGGCA
1301 AAAAAGGGGA TACAGTGGAA CTGACCTGTA CAGCTTCCCA GAAGAAGAGC
1351 ATACAATTCC ACTGGAAAAA CTCCAACCAG ATAAAGATTG TGGGAAATCA
1401 GGGCTCTTTC TTAAGTAAAG GTCCATCCAA GCTGAATGAT CGCGCTGACT
  
```

3824

89085519

SEQUENCE SHEET

69/93

FIG. 22(cont'd)

1451 CAAGAAGAAG CTTGTGGGAC CAAGGAAACT TTCCCCTGAT CATCAAGAAT
 1501 CTTAAGATAG AAGACTCAGA TACTTACATC TGTGAAGTGG AGGACCAGAA
 1551 GGAGGAGGTG CAATTGCTAG TGTTCGGATT GACTGCCAAC TCTGACACCC
 1601 ACCTGCTTCA GGGGCAGAGC CTGACCCTGA CCTTGGAGAG CCCCCGGGT
 1651 AGTAGCCCCT CAGTGCAATG AGATCTTTGT GAAGGAACCT TACTTCTGTG
 1701 GTGTGACATA ATTGGACAAA CTACCTACAG AGATTTAAAG CTCTAAGGTA
 1751 AATATAAAAT TTTTAAGTGT ATAATGTGTT AACTACTGA TTCTAATTGT
 1801 TTGTGTATTT TAGATTCCAA CCTATGGAAC TGATGAATGG GAGCAGTGGT
 1851 GGAATGCCTT TAATGAGGAA AACCTGTTTT GCTCAGAAGA AATGCCATCT
 1901 AGTGATGATG AGGCTACTGC TGA CTCTCAA CATTCTACTC CTCCAAAAA
 1951 GAAGAGAAAAG GTAGAAGACC CCAAGGACTT TCCTTCAGAA TTGCTAAGTT
 2001 TTTTGAGTCA TGCTGTGTTT AGTAATAGAA CTCTTGCTTG CTTTGCTATT
 2051 TACACCACAA AGGAAAAAGC TGC ACTGCTA TACAAGAAAA TTATGGAAAA
 2101 ATATTCTGTA ACCTTTATAA GTAGGCATAA CAGTTATAAT CATAACATAC
 2151 TGTTTTTTCT TACTCCACAC AGGCATAGAG TGTCTGCTAT TAATACTAT
 2201 GCTCAAAAAT TGTGTACCTT TAGCTTTTTA ATTTGTAAAG GGGTTAATAA
 2251 GGAATATTTG ATGTATAGTG CCTTGACTAG AGATCATAAT CAGCCATACC
 2301 ACATTTGTAG AGGTTTTACT TGCTTTAAAA AACCTCCCAC ACCTCCCCCT
 2351 GAACCTGAAA CATAAAATGA ATGCAATTGT TGTGTTAAC TTGTTTATTG
 2401 CAGCTTATAA TGGTTACAAA TAAAGCAATA GCATCACAAA TTTACAAAT
 2451 AAAGCATTTT TTTCACTGCA TTCTAGTTGT GGTGTGTTCA AACTCATCAA
 2501 TGTATCTTAT CATGTCTGGA TCCTCTACGC CGGACGCATC GTGGCCGGCA
 2551 TCACCGGCGC CACAGGTGCG GTTGCTGGCG CCTATATCGC CGACATCACC
 2601 GATGGGGAAG ATCGGGCTCG CCACTTCGGG CTCATGAGCG CTTGTTTCGG
 2651 CGTGGGTATG GTGGCAGGCC CGTGGCCGGG GGA CTGTTGG GCGCCATCTC
 2701 CTTGCATGCA CCATTCCTTG CGGCGGCGGT GCTCAACGGC CTCAACCTAC
 2751 TACTGGGCTG CTTCTAATG CAGGAGTCGC ATAAGGGAGA GCGTCGACCG
 2801 ATGCCCTTGA GAGCCTTCAA CCCAGTCAGC TCCTTCCGGT GGGCGCGGGG
 2851 CATGACTATC GTCGCCGCAC TTATGACTGT CTTCTTTATC ATGCAACTCG
 2901 TAGGACAGGT GCCGGCAGCG CTCTGGGTCA TTTTCGGCGA GGACCGCTTT
 2951 CGCTGGAGCG CGACGATGAT CGGCCTGTCT CTTGCGGTAT TCGGAATCTT
 3001 GCACGCGCTC GCTCAAGCCT TCGTCACTGG TCCCGCCACC AAACGTTTCG
 3051 GCGAGAAGCA GGCCTATTA C GCGGCAATGG CCGCCGACGC GCTGGGCTAC

3825

89085519

FIG. 22(cont'd)

3101 GTCTTGCTGG CGTTCGCGAC CCGAGGCTGG ATGGCCTTCC CCATTATGAT
 3151 TCTTCTCGCT TCCGGCGGCA TCGGGATGCC CGCGTTGCAG GCCATGCTGT
 3201 CCAGGCAGGT AGATGACGAC CATCAGGGAC AGCTTCAAGG ATCGCTCGCG
 3251 GCTCTTACCA GCCTAACTTC GATCACTGGA CCGCTGATCG TCACGGCGAT
 3301 TTATGCCGCE TCGGCGAGCA CATGGAACGG GTTGGCATGG ATTGTAGGCG
 3351 CCGCCCTATA CCTTGTCTGC CTCCCCGCGT TCGGTGCGGG TGCATGGAGC
 3401 CGGGCCACCT CGACCTGAAT GGAAGCCGGC GGCACCTCGC TAACGGATTG
 3451 ACCACTCCAA GAATTGGAGC CAATCAATTC TTGCGGAGAA CTGTGAATGC
 3501 GCAAACCAAC CCTTGGCAGA ACATATCCAT CGCGTCCGCC ATCTCCAGCA
 3551 GCCGCACGCG GCGCATCTCG GGCCGCGTTG CTGGCGTTTT TCCATAGGCT
 3601 CCGCCCCCCT GACGAGCATC ACAAAAATCG ACGCTCAAGT CAGAGGTGGC
 3651 GAAACCCGAC AGGACTATAA AGATACCAGG CGTTTCCCCC TGGAAAGCTCC
 3701 CTCGTGCGCT CTCCTGTTCC GACCCTGCCG CTTACCGGAT ACCTGTCCGC
 3751 CTTTCTCCCT TCGGGAAGCG TGGCGCTTTC TCAATGCTCA CGCTGTAGGT
 3801 ATCTCAGTTC GGTGTAGGTC GTTCGCTCCA AGCTGGGCTG TGTGCACGAA
 3851 CCCCCCGTTC AGCCCGACCG CTGCGCCTTA TCCGGTAACT ATCGTCTTGA
 3901 GTCCAACCCG GTAAGACACG ACTTATCGCC ACTGGCAGCA GCCACTGGTA
 3951 ACAGGATTAG CAGAGCGAGG TATGTAGGCG GTGCTACAGA GTTCTTGAAG
 4001 TGGTGGCCTA ACTACGGCTA CACTAGAAGG ACAGTATTTG GTATCTGCGC
 4051 TCTGCTGAAG CCAGTTACCT TCGGAAAAAG AGTTGGTAGC TCTTGATCCG
 4101 GCAAACAAAC CACCGCTGGT AGCGGTGGTT TTTTGTGTTG CAAGCAGCAG
 4151 ATTACGCGCA GAAAAAAGG ATCTCAAGAA GATCCTTTGA TCTTTTCTAC
 4201 GGGGTCTGAC GCTCAGTGGA ACGAAAACCTC ACGTTAAGGG ATTTTGGTCA
 4251 TGAGATTATC AAAAAGGATC TTCACCTAGA TCCTTTTAAA TAAAAAATGA
 4301 AGTTTAAAT CAATCTAAAG TATATATGAG TAAACTTGGT CTGACAGTTA
 4351 CCATGCTTA ATCAGTGAGG CACCTATCTC AGCGATCTGT CTATTTCTGT
 4401 CATCCATAGT TGCTGACTC CCGTCTGTGT AGATAACTAC GATACGGGAG
 4451 GCTTACCAT CTGGCCCCAG TGCTGCAATG ATACCGCGAG ACCCAGCTC
 4501 ACCGGCTCCA GATTTATCAG CAATAAACCA GCCAGCCGGA AGGGCCGAGC
 4551 GCAGAAGTGG TCCTGCAACT TTATCCGCCT CCATCCAGTC TATTAATTGT
 4601 TGCCGGGAAG CTAGAGTAAG TAGTTCGCCA GTTAATAGTT TGCACAACGT
 4651 TGTTCCTATT GCTGAGGCA TCGGGTCTC ACGCTGCTCG TTTGGTATGG
 4701 CTTCATTGAG CTCCCTTCC CAACGATCAA GGCGAGTTAC ATGATCCCCC

3826

89085519

SUBSTITUTE SHEET


71/93

FIG. 22(cont'd)


4751 ATGTTGTGCA AAAAAGCGGT TAGCTCCTTC GGTCTCCGA TCGTTGTCAG
4801 AAGTAAGTTG GCCGCAGTGT TATCACTCAT GGTTATGGCA GCACTGCATA
4851 ATTCTCTTAC TGTCATGCCA TCCGTAAGAT GCTTTTCTGT GACTGGTGAG
4901 TACTCAACCA AGTCATTCTG AGAATAGTGT ATGCGGCGAC CGAGTTGCTC
4951 TTGCCCCGGCG TCAACACGGG ATAATACCGC GCCACATAGC AGAACTTTAA
5001 AAGTGCTCAT CATTGGAAAA CGTTCTTCGG GCGGAAACT CTCAAGGATC
5051 TTACCGCTGT TGAGATCCAG TTCGATGTAA CCCACTCGTG CACCCAAGTG
5101 ATCTTCAGCA TCTTTTACTT TCACCAGCGT TTCTGGGTGA GCAAAAACAG
5151 GAAGGCAAAA TGCCGCAAAA AAGGGAATAA GGGCGACACG GAAATGTTGA
5201 ATACTCATAC TCTTCCTTTT TCAATATTAT TGAAGCATTT ATCAGGGTTA
5251 TTGTCTCATG AGCGGATACA TATTTGAATG TATTTAGAAA AATAAACAAA
5301 TAGGGGTTC GCGCACATTT CCCCAGAAAAG TGCCACCTGA CGTCTAAGAA
5351 ACCATTATTA TCATGACATT AACCTATAAA AATAGGCGTA TCACGAGGCC
5401 CTTTCGTCTT CAA

72/93

FIG. 23



73/93



3829

FIG. 24

20045510

74/93




FIG. 25

75/93





FIG. 26

3331


89085519

76/93


FIG. 27



Site directed mutagenesis
to change an Asn at amino
acid position #3 to a Lys,
using T4-66




Site directed mutagenesis
to delete Gln and Gly at
amino acid positions #1, #2,
using T4AID-87



77/93

FIG. 28



3833

78/93




FIG. 29A

3833^A

89085519

79/93




FIG. 29B

20/93






FIG. 31



3836

89085519

82/93




FIG. 32

3837

89085519

83/93




FIG.33

89085519

83/93





FIG.33

89085519


84/93

FIG. 34



8:/93


FIG. 35



3840

86/93

FIG. 36




3841

89085519


87/93

FIG. 37



88/93

FIG. 38
Competition ELISA for r574.113




3843

89085519

89/93

FIG. 39


P24 ANTIGEN RIA



3844


89085519

FIG. 40
C8/66 CELL FUSION ASSAY




5./93

FIG. 41




92/93

FIG. 42



3847

93/93

**FIG. 43**

MICROORGANISMS

Optional Sheet in connection with the microorganism referred to on page 95, lines 29-35 of the description:
96, lines 6-11

A. IDENTIFICATION OF DEPOSIT *

96, lines 19-21

Further deposits are identified on an additional sheet ☒ -- 3 additional sheets attached

Name of depositary institution *

In Vitro International, Inc.

Address of depositary institution (including postal code and country) *

611 (P) Hammocks
 Ferry Road, Linthicum, Maryland 21090
 United States of America

Date of deposit *

See attached additional sheets

Accession Number *

See attached additional sheets

B. ADDITIONAL INDICATIONS * (Leave blank if not applicable). This information is continued on a separate attached sheet ☐

In respect of those designations in which a European patent is sought samples of the deposited microorganisms will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).

C. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE * (If the indications are not for all designated States)

D. SEPARATE FURNISHING OF INDICATIONS * (Leave blank if not applicable)

The indications listed below will be submitted to the International Bureau later * (Specify the general nature of the indications e.g., "Accession Number of Deposit")

E. ☐ This sheet was received with the international application when filed (to be checked by the receiving Office)

(Authorized Officer)

☐ The date of receipt (from the applicant) by the International Bureau is:

was 13 JANUARY 1989
 (13. 01. 89)

(Authorized Officer)

(January 1985)

Additional Sheet 1 of 3 To Form
PCT/RO/134

Continuation Of Box A

IDENTIFICATION OF DEPOSITS

BG378: E.coli MC1061/pBG378
 199-7: E.coli MC1061/p199-7
 170-2: E.coli JA221/p170-2
 EC100: E.coli JM83/pEC100
 BG377: E.coli MC1061/pBG377
 BG380: E.coli MC1061/pBG380
 BG381: E.coli MC1061/pBG381

DATE OF DEPOSITS

2 September 1987

ACCESSION NUMBERS

IVI 10143
 IVI 10144
 IVI 10145
 IVI 10146
 IVI 10147
 IVI 10148
 IVI 10149

3850

89085519

Additional Sheet 2 of 2 To Form
PCT/RO/134

Continuation Of Box A

IDENTIFICATION OF DEPOSITS

BG-391:	<u>E.coli</u>	MC1061/pBG391
BG-392:	<u>E.coli</u>	MC1061/pBG392
BG-393:	<u>E.coli</u>	MC1061/pBG393
BG-394:	<u>E.coli</u>	MC1061/pBG394
BG-396:	<u>E.coli</u>	MC1061/pBG396
203-5:	<u>E.coli</u>	SG936/p203-5

DATE OF DEPOSITS

6 January 1988

ACCESSION NUMBERS

IVI 10151
IVI 10152
IVI 10153
IVI 10154
IVI 10155
IVI 10156

89085519

3851

Additional Sheet 3 of 3 To Form
PCT/RO/134

Continuation Of Box A

IDENTIFICATION OF DEPOSITS

211-11: E.coli A89/pBG211-11
214-10: E.coli A89/pBG214-10
215-7: E.coli A89/pBG215-7

DATE OF DEPOSITS

24 August 1988

ACCESSION NUMBERS

IVI 10183
IVI 10184
IVI 10185

3852

89085519

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US88/02940

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) *

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC(4): C07H 15/12; C12G 1/70, C12Q 1/02, see attachment.

U.S. CL.: 536/27; 435/5,29,39,68,91,170,172.3, see attachment.

II. FIELDS SEARCHED

Minimum Documentation Searched *

Classification System	Classification Symbols
U.S.	435/5,29,39,68,91,170,172.1,172.3,240,255,320; 530/350,412; 514/2; 424/85; 536/27; 935/6, 9, 11,12,15,22,23,24,59,60,65,66

Documentation Searched other than Minimum Documentation
to the extent that such documents are included in the fields searched *

Chemical Abstract Data Base (CAS) 1967-1988; Biosis Data Base 1969-1988 Keywords: CD4, T4, TCell, AIDS, HTLV, HTLVI, HTLVIII, see attachment.

III. DOCUMENTS CONSIDERED TO BE RELEVANT *

Category *	Citation of Document, with indication, where appropriate, of the relevant passages *	Relevant to Claim No. *
Y	<u>SCIENCE</u> , Volume 234, issued 1986 November, (Washington, DC., U.S.A.), (Q.J. SATTENTAU ET AL), "Epitopes of the CD4 Antigen and HIV Infection" See pages 1120-1123. See particularly page 1120	13-20, 29-33 and 48-52
Y	<u>SCIENCE</u> , Volume 234, issued 1986, November, (Washington, D.C. U.S.A) (J.A. HOXIE ET AL), "Alterations in T4 (CD4) Protein and mRNA synthesis in Cells Infected with HIV" see pages 1123-1127. See particularly page 1123.	13-20, 29-33, and 48-52
Y,P	<u>PROCEEDINGS NATIONAL ACADEMY OF SCIENCES</u> , U.S.A, Volume 84, issued 1987 December (Washington, D.C., U.S.A), (P.J. MADDON ET AL.), "Structure and Expression of the Human and Mouse T4 Genes", See pages 9155-9159, See particularly page 9155 and 9156.	1-4,25-27, 34-36 and 39-46

* Special categories of cited documents: *

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

Date of Mailing of this International Search Report

20 NOVEMBER 1988

03 FEB 1989

International Searching Authority

ISA/US

Signature of Authorizing Officer

Richard C. Pezz
RICHARD C. PEZZ

Attachment to PCT/ISA/210
I. Classification of Subject Matter

IPC: C12Q 1/06, C12P 21/00, C12P 19/34, C12P 1/04, C12N
15/00, C12N 7/00; C07K 13/00, C07K 3/00; A61K
37/68; A61K 39/00, A61K 45/02

US.CL.: 240, 320; 530/350, 412; 514/2; 424/85

II. Fields Searched

Keywords: ARC, Surface, receptor, therap?, purif?,
Immunoassay, Detection, Pharmaceutical Composition,
Lymphocyte, Igg, Polyvalent, Solub?, gene, Clon?, Protein,
Polypeptide, Fusion, Expression, Vector, Plasmid, Surface
Protein, Surface Antigen, Acquired Immune Deficiency
Syndrome, Retrovirus

3854

89085519

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	<u>PROCEEDINGS NATIONAL ACADEMY OF SCIENCES, U.S.A.</u> , Volume 84, issued 1987, June (Washington, D.C., U.S.A.), (T.C. CHANH ET AL.), "Monoclonal Anti-idiotypic Antibody Mimics the CD4 Receptor and Binds Human Immunodeficiency Virus" See pages 3891-3895. See particularly page 3891.	13-24, 29-33 and 48-52
X Y	<u>CELL</u> , Volume 47, issued 1986, November, (Cambridge, Mass., U.S.A) (P.J. MADDON ET AL), "The T4 Gene Encodes the AIDS Virus Receptor and is Expressed in the Immune System and the Brain", See pages 333-348, See particularly pages 333-335.	1,3-6 and 25-27 2,7-24 and 28-50
P, Y	<u>CHEMICAL ABSTRACTS</u> , Volume 107, no. 15, issued 1987 October 12 (Columbus, Ohio, U.S.A), T.L. LENTZ et al, "Rabies virus binding to cellular membranes measured by enzyme immunoassay" see page 359, column 1, the abstract no. 131853f, Muscle Nerve, 1985, 8(4), 336-345 (Eng).	16-18 32-33 and 50
Y	<u>CHEMICAL ABSTRACTS</u> , Volume 106, no. 21, issued 1987, May 25, (Columbus, Ohio, U.S.A), J.P. ZIMMER ET AL., 'Diphenylhydantoin (DPH) blocks HIV-receptor on T-lymphocyte surface', see page 123, column 1, the abstract no. 168522c, Biut, 1986, 53(6), 447-450 (Eng).	13-15, 19-20, 29-30, 48-49 and 51-52
Y, P	<u>BIOLOGICAL ABSTRACTS</u> , Volume 85, no. 4, issued 1988, April 15 (Philadelphia, PA, U.S.A), A.G. DALGLEISH ET AL., 'Neutralization of HIV isolates by anti-idiotypic antibodies which mimic the T4 (CD4) epitope: A potential AIDS vaccine' see page 222, abstract no. 37595, Lancet 2 (8567): 1047-1050 (Eng).	13-15, 19-20, 29-30, 48-49, and 51-52

3855

89085519